

BRAIDED HOPF ALGEBRAS

Shouchuan Zhang

Department of Mathematics, Hunan University, 410082
P.R.China

Hunan Normal University Press
First edition 1999
Second edition 2005
Third edition 2007
ISBN7-81031-812-8/0.036

Preface

The term “quantum group” was popularized by Drinfeld in his address to the International Congress of Mathematicians in Berkeley . However, the concepts of quantum groups and quasitriangular Hopf algebras are the same. Therefore, Hopf algebras have close connections with various areas of mathematics and physics.

The development of Hopf algebras can be divided into five stages .

The first stage is that integral and Maschke’s theorem are found. Maschke’s theorem gives a nice criterion of semisimplicity, which is due to M.E. Sweedler. The second stage is that the Lagrange’s theorem is proved, which is due to W.D. Nichols and M.B. Zoeller. The third stage is the research of the actions of Hopf algebras, which unifies actions of groups and Lie algebras. S. Montgomery’s “Hopf algebras and their actions on rings” contains the main results in this area. S. Montgomery and R.J. Blattner show the duality theorem. S. Montgomery, M. Cohen, H.J. Schneider, W. Chin, J.R. Fisher and the author study the relation between algebra R and the crossed product $R\#_{\sigma}H$ of the semisimplicity, semiprimeness, Morita equivalence and radical. S. Montgomery and M. Cohen ask whether $R\#_{\sigma}H$ is semiprime when R is semiprime for a finite-dimensional semiprime Hopf algebra H . This is a famous semiprime problem. The answer is sure when the action of H is inner or H is commutative or cocommutative, which is due to S. Montgomery, H.J. Schneider and the author. This is the best result in the study of the semiprime problem up to now. However, this problem is still open. Y. Doi and M. Takeachi show that the crossed product is a cleft extension.

The research of quantum groups is the fourth stage. The concepts of quantum groups and (co)quasitriangular Hopf algebras are the same. The Yang-Baxter equation first came up in a paper by Yang as factorization condition of the scattering S-matrix in the many-body problem in one dimension and in work of Baxter on exactly solvable models in statistical mechanics. It has been playing an important role in mathematics and physics (see [10], [76]). Attempts to find solutions of the Yang-Baxter equation (YBE) in a systematic way have led to the theory of quantum groups. In other words, we can obtain a solution of the Yang-Baxter equation by a (co)quasitriangular Hopf algebra. It is well-known that the universal enveloping algebras of Lie algebras are Hopf algebras. But they are not quasitriangular in general. Drinfeld obtained a quasitriangular Hopf algebra by means of constructing a quantum enveloping algebras of simple Lie algebra. Drinfeld also obtained a quasitriangular Hopf algebra $D(H)$, quantum double, for any Hopf algebra H . D.E. Radford and R.G. Larson study the algebraic structure of (co)quasitriangular Hopf algebras. Kassel’s “Quantum Group” contains the main results in

this area.

The research of braided Hopf algebras and classification of finite-dimensional Hopf algebras are the fifth stage. Supersymmetry has attracted a great deal of interest from physicists and mathematicians (see [65], [110], [88]). It finds numerous applications in particle physics and statistical mechanics (see [30]) . Of central importance in supersymmetric theories is the Z_2 -graded structure which permeates them. Superalgebras and super-Hopf algebras are most naturally very important.

Braided tensor categories were introduced by Joyal and Street in 1986 [61]. It is a generalization of super case. Majid, Joyal, Street and Lyubasheko have reached many interesting conclusions in braided tensor categories, for example, the braided reconstruction theorem, transmutation and bosonisation, integral, q -Fourier transform, q -Mikowski space, random walk and so on (see [76], [82], [83], [88], [73]).

The main results of my research consist of the following:

In braided tensor categories we show the Maschke's theorem and give the necessary and sufficient conditions for double cross biproducts and crossbiproducts and biproducts to be bialgebras. We obtain the factorization theorem for braided Hopf algebras;

In symmetric tensor categories, we show the duality theorem and construct quantum double;

In ordinary vector space category with ordinary twist, we obtain the relation between the global dimension, the weak dimension, the Jacobson radical, Baer radical of algebra R and its crossed product $R \#_\sigma H$. We also give the relation between the decompositions of comodules and coalgebras. We obtain all solutions of constant classical Yang-Baxter equation (CYBE) in Lie algebra L with $\dim L \leq 3$. We also give the sufficient and necessary conditions for $(L, [\], \Delta_r, r)$ to be a coboundary (or triangular) Lie bialgebra.

These are also the points of originality of this book.

This book is organized into three parts. In the three parts we study Hopf algebras in braided tensor categories, symmetric braided tensor categories and the category of vector spaces over field k with ordinary twist braiding respectively. Furthermore, we divide the three parts into twelve chapters and they are briefly described as follows.

In Chapter 1, we recall the basic concepts and conclusions of Hopf algebras living in braided tensor categories.

In Chapter 2, we obtain the fundamental theorem of Hopf modules living in braided tensor categories. In Yetter-Drinfeld category, we obtain that the antipode S of H is invertible and the integral $\int_H^l$ of H is a one-dimensional space for any finite-dimensional braided Hopf algebra H . We give Maschke's Theorem for YD Hopf algebras.

In Chapter 3, the double biproduct $D = A \bowtie^b H$ of two bialgebras A and H is constructed in a braided tensor category and the necessary and sufficient conditions for D to be a bialgebra are given. The universal property of double biproduct is obtained. In particular, the necessary and sufficient conditions for double biproducts, double products (or bicrossed products), double coproducts, Drinfeld double and smash products of two anyonic (or super) Hopf algebras A and H to be anyonic Hopf algebras (or super-Hopf algebras) are obtained.

In Chapter 4, we construct the bicrossproducts and biproducts in braided tensor category

$(\mathcal{C}, C)$ and give the necessary and sufficient conditions for them to be bialgebras. We obtain that for any left module (A, α) of bialgebra H the smash product $A \# H$ is a bialgebra iff A is an H -module bialgebra and H is cocommutative with respect to (A, α) .

In Chapter 5, We show that the braided group analogue of double cross (co)product is the double cross (co)product of braided group analogues. we give the factorization theorem of Hopf algebras living in braided tensor categories and the relation between Hopf algebra H and its factors.

In Chapter 6, braided m-Lie algebras induced by multiplication are introduced, which generalize Lie algebras, Lie color algebras and quantum Lie algebras. The necessary and sufficient conditions for the braided m-Lie algebras to be strict Jacobi braided Lie algebras are given. Two classes of braided m-Lie algebras are given, which are generalized matrix braided m-Lie algebras and braided m-Lie subalgebras of $End_F M$, where M is a Yetter-Drinfeld module over B with $\dim B < \infty$. In particular, generalized classical braided m-Lie algebras $sl_{q,f}(GM_G(A), F)$ and $osp_{q,t}(GM_G(A), M, F)$ of generalized matrix algebra $GM_G(A)$ are constructed and their connection with special generalized matrix Lie superalgebra $sl_{s,f}(GM_{\mathbf{Z}_2}(A^s), F)$ and orthosymplectic generalized matrix Lie super algebra $osp_{s,t}(GM_{\mathbf{Z}_2}(A^s), M^s, F)$ are established. The relationship between representations of braided m-Lie algebras and their associated algebras are established.

In Chapter 7, we construct the Drinfeld double in symmetric tensor category.

In Chapter 8, we give the duality theorem for Hopf algebras living in a symmetric braided tensor category. Using the result, we show that

$$(R \# H) \# H^{\hat{*}} \cong M_n(R) \quad \text{as algebras}$$

when H is a finite-dimensional Hopf algebra living in a symmetric braided vector category. In particular, the duality theorem holds for any finite-dimensional super-Hopf algebra H .

In chapter 9, we give the relation between the decomposition of coalgebras and comodules.

In Chapter 10, we obtain that the global dimensions of R and the crossed product $R \#_{\sigma} H$ are the same; in the meantime, their weak dimensions are also the same when H is finite-dimensional semisimple and cosemisimple Hopf algebra.

In Chapter 11, we obtain that the relation between H -radical of H -module algebra R and radical of $R \# H$.

In Chapter 12, we give that the relation between H -radical of twisted H -module algebra R and radical of crossed product $R \#_{\sigma} H$. We also obtain that if H is a finite-dimensional semisimple, cosemisimple, and either commutative or cocommutative Hopf algebra, then R is H -semiprime iff R is semiprime iff $R \#_{\sigma} H$ is semiprime.

In Chapter 13, all solutions of constant classical Yang-Baxter equation (CYBE) in Lie algebra L with $\dim L \leq 3$ are obtained and the sufficient and necessary conditions for $(L, [\quad], \Delta_r, r)$ to be a coboundary (or triangular) Lie bialgebra are given.

The strongly symmetric elements in $L \otimes L$ are found and they are all solutions of CYBE.

In Chapter 14, we classify quiver Hopf algebras. This leads to the classification of multiple Taft algebras as well as pointed Yetter-Drinfeld modules and their corresponding Nichols algebras. In particular, when the ground-field k is the complex field and G is a finite abelian group,

we classify quiver Hopf algebras over G , multiple Taft algebras over G and Nichols algebras in ${}^k_G\mathcal{YD}$. We show that the quantum enveloping algebra of a complex semisimple Lie algebra is a quotient of a semi-path Hopf algebra.

I would like to express my gratitude to my advisors Professor Wenting Tong and Professor Yonghua Xu.

I would like to express my gratitude to Professor Boxun Zhou for his support.

I am very grateful to Professor Nanqing Ding, Professor Fanggui Wang, Professor Jianlong Chen, Professor Huixiang Chen, Professor Guoqiang Hu, Professor Jinqing Li and Professor Minyi Wang for their help.

Finally, I wish to thank my parents, my wife and all of my friends for their support.

Contents

I	Hopf Algebras Living in Braided Tensor Categories	9
1	Preliminaries	10
2	The Maschke's Theorem for Braided Hopf Algebras	34
2.1	The integrals in braided tensor categories	35
2.2	The Integrals of Braided Hopf algebras in Yetter-Drinfeld Categories	40
3	The Double Bicrossproducts in Braided Tensor Categories	46
3.1	Double bicrossproduct	51
3.2	The universal property of double bicrossproduct	75
4	The Bicrossproducts in Braided Tensor Categories	85
4.1	Crossed Products	97
4.2	Bicrossproducts	103
4.3	Biproducts	115
5	The Factorization Theorem of Bialgebras in Braided Tensor Categories	125
5.1	the braided reconstruction theorem	125
5.2	Braided group analogues of double cross products	138
5.3	The factorization of ordinary bialgebras	148
6	Braided Lie Algebras	150
6.1	Braided m-Lie Algebras	151
6.2	Generalized Matrix Braided m-Lie Algebras	154
6.3	Braided m-Lie Algebras in the Yetter-Drinfeld Category and Their Representations	158
II	Hopf Algebras Living in Symmetric Tensor Categories	161
7	The Quasitriangular Hopf Algebras in Symmetric Braided Tensor Categories	162
7.1	Yang-Baxter equation	167
7.2	The antipode of quasitriangular Hopf algebra	168
7.3	Drinfeld double	169

7.4	(Co) quasitriangular Hopf algebras of double cross (co)products	194
8	The Duality Theorem for Braided Hopf Algebras	200
8.1	The duality theory for Hopf algebras living in symmetric tensor categories . . .	201
8.2	The duality theory for Yetter-Drinfeld Hopf algebras	219
III	Hopf Algebras Living in the Category of Vector spaces	220
9	The Relation Between Decomposition of Comodules and Coalgebras	221
9.1	The relation between the decomposition of comodules and coalgebras	222
9.2	The coradicals of comodules	233
10	The Homological Dimensions of Crossed Products	236
10.1	The homological dimensions of modules over crossed products	236
10.2	The global dimensions and weak dimensions of crossed products	242
11	The Radicals of Hopf Module Algebras	244
11.1	The H -special radicals for H -module algebras	245
11.2	H -Baer radical	247
11.3	The H -module theoretical characterization of H -special radicals	249
12	The H-Radicals of Twisted H-Module Algebras	259
12.1	The Baer radical of twisted H -module algebras	260
12.2	The Baer and Jacobson radicals of crossed products	261
12.3	The general theory of H -radicals for twisted H -module algebras	269
12.4	The relations among radical of R , radical of $R\#_{\sigma}H$, and H -radical of R	270
12.5	The H -Von Neumann regular radical	275
12.6	About J.R. Fisher's question	278
13	Classical Yang-Baxter Equation And Low Dimensional Triangular Lie Algebras	280
13.1	The solutions of CYBE	282
13.2	Coboundary Lie bialgebras	290
13.3	Strongly symmetric elements	293
13.4	The solutions of CYBE in the case of $\text{char } k = 2$	295
13.5	Coboundary Lie bialgebras in the case of $\text{char } k = 2$	300
13.6	The solutions of CYBE with $\text{char } k \neq 2$	305
13.7	The solutions of CYBE with $\text{char } k = 2$	307
13.8	Coboundary Lie bialgebras	308
14	Classification of PM Quiver Hopf Algebras	310
14.1	PM quiver Hopf algebras	313
14.2	Multiple Taft algebras	321

14.3	The relation between quiver Hopf algebras and quotients of free algebras	327
14.3.1	The structure of multiple Taft algebras and semi-path Hopf algebras . . .	327
14.3.2	A characterization of multiple Taft algebras	333
14.3.3	The relation between semi-path Hopf algebras and quantum enveloping algebras	335
14.4	Classification of ramification systems for the symmetric group	340
14.4.1	Ramification Systems and Isomorphisms	342
14.4.2	Action of Group	344
14.4.3	The structure of centralizer in S_n	344
14.4.4	Characters of Z_σ	347
14.4.5	Fixed Point Set $F_{(\phi_{g_C}, \phi_C)_C}$	349
14.4.6	The Formula Computing the Number of Isomorphic Classes	350
14.4.7	Representative	353
15	Appendix	355

Part I

Hopf Algebras Living in Braided Tensor Categories

Chapter 1

Preliminaries

We begin with the tensor category. We define the product $\mathcal{C} \times \mathcal{D}$ of two category $\mathcal{C}$ and $\mathcal{D}$ whose objects are pairs of objects $(U, V) \in (ob\mathcal{C}, ob\mathcal{D})$ and whose morphisms are given by

$$Hom_{\mathcal{C} \times \mathcal{D}}((V, W)(V', W')) = Hom_{\mathcal{C}}(V, V') \times Hom_{\mathcal{D}}(W, W').$$

Let $\mathcal{C}$ be a category and $\otimes$ be a functor from $\mathcal{C} \times \mathcal{C}$ to $\mathcal{C}$. This means

- (i) we have object $V \otimes W$ for any $V, W \in ob\mathcal{C}$;
- (ii) we have morphism $f \otimes g$ from $U \otimes V$ to $X \otimes Y$ for any morphisms f and g from U to X and from V to Y ;
- (iii) we have

$$(f \otimes g)(f' \otimes g') = (ff') \otimes (gg')$$

for any morphisms $f : U \rightarrow X, g : V \rightarrow Y, f' : U' \rightarrow V$ and $g' : V' \rightarrow V$;

- (iv) $id_{U \otimes V} = id_U \otimes id_V$.

Let $\otimes\tau$ denote the functor from $\mathcal{C} \times \mathcal{C}$ to $\mathcal{C}$ such that $(\otimes\tau)(U, V) = (V \otimes U)$ and $(\otimes\tau)(f, g) = g \otimes f$, for any objects U, V, X, Y in $\mathcal{C}$, and for any morphisms $f : U \rightarrow X$ and $g : V \rightarrow Y$.

An associativity constraint a for tensor $\otimes$ is a natural isomorphism

$$a : \otimes(\otimes \times id) \rightarrow \otimes(id \times \otimes).$$

This means that, for any triple (U, V, W) of objects of $\mathcal{C}$, there is a morphism $a_{U,V,W} : (U \otimes V) \otimes W \rightarrow U \otimes (V \otimes W)$ such that

$$a_{U',V',W'}((f \otimes g) \otimes h) = (f \otimes (g \otimes h))a_{U,V,W}$$

for any morphisms f, g and h from U to U' , from V to V' and from W to W' respectively.

Let I be an object of $\mathcal{C}$. If there exist natural isomorphisms

$$l : \otimes(I \times id) \rightarrow id \quad \text{and} \quad r : \otimes(id \times I) \rightarrow id ,$$

then I is called the unit object of $\mathcal{C}$ with left unit constraint l and right unit constraint r .

Definition 1.0.1. $(\mathcal{C}, \otimes, I, a, l, r)$ is called a tensor category if $\mathcal{C}$ is equipped with a tensor product $\otimes$, with a unit object I , an associativity constraint a , a left unit constraint l and a right unit constraint r such that the Pentagon Axiom and the Triangle Axiom are satisfied, i.e.

$$(id_U \otimes a_{V,W,X})a_{U,V \otimes W,X}(a_{U,V,W} \otimes id_X) = a_{U,V,W \otimes X}a_{U \otimes V,W,X}$$

and

$$(id_V \otimes l_W)a_{V,I,W} = r_V \otimes id_W$$

for any $U, V, W, X \in ob\mathcal{C}$.

Furthermore, if there exists a natural isomorphism

$$C : \otimes \rightarrow \otimes \tau$$

such that the Hexagon Axiom holds, i.e.

$$a_{V,W,U}C_{U,V \otimes W}a_{U,V,W} = (id_V \otimes C_{U,W})a_{V,U,W}(C_{U,V} \otimes id_W)$$

and

$$a_{W,U,V}^{-1}C_{U \otimes V,W}a_{U,V,W}^{-1} = (C_{U,W} \otimes id_V)a_{U,W,V}^{-1}(id_U \otimes C_{V,W}),$$

for any $U, V, W \in ob\mathcal{C}$, then $(\mathcal{C}, \otimes, I, a, l, r, C)$ is called a braided tensor category. In this case, C is called a braiding of $\mathcal{C}$. If $C_{U,V} = C_{V,U}^{-1}$ for any $U, V \in ob\mathcal{C}$, then $(\mathcal{C}, C)$ is called a symmetric braided tensor category or a symmetric tensor category. Here functor $\tau : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C} \times \mathcal{C}$ is the flip functor defined by $\tau(U \times V) = V \times U$ and $\tau(f \times g) = g \times f$, for any $U, V \in \mathcal{C}$, and morphisms f and g . Note that we denote the braiding C in braided tensor category $(\mathcal{C}, \otimes, I, a, l, r, C)$ by ${}^C C$ sometimes.

A tensor category $(\mathcal{C}, \otimes, I, a, l, r)$ is said to be strict if a, r, l are all identities in the category.

By Definition, the following Lemmas are clear

Lemma 1.0.2. Let $\otimes$ be a functor from $\mathcal{C} \times \mathcal{C}$ to $\mathcal{C}$ and I an object in $\mathcal{C}$. Then $(\mathcal{C}, \otimes, I, a = id, l = id, r = id)$ is a strict tensor category iff

(ST1): $(U \otimes V) \otimes W = U \otimes (V \otimes W)$ for any $U, V, W \in ob\mathcal{C}$;

(ST2): $I \otimes U = U \otimes I = U$ for any $U \in ob\mathcal{C}$.

Lemma 1.0.3. Let $\otimes$ be a functor from $\mathcal{C} \times \mathcal{C}$ to $\mathcal{C}$ and I an object in $\mathcal{C}$. Assume that there exists a natural isomorphism $C : \otimes \rightarrow \otimes \tau$. Then $(\mathcal{C}, \otimes, I, a = id, l = id, r = id, C)$ is a strict braided tensor category iff $(\mathcal{C}, \otimes, I, a = id, l = id, r = id)$ is a strict tensor category and

(SBT3): $C_{U,V \otimes W} = (id_V \otimes C_{U,W})(C_{U,V} \otimes id_W)$ for any $U, V, W \in ob\mathcal{C}$;

(SBT4): $C_{U \otimes V,W} = (C_{U,W} \otimes id_V)(id_U \otimes C_{V,W})$ for any $U, V, W \in ob\mathcal{C}$.

Let $(\mathcal{C}, \otimes, I, a, l, r)$ and $(\mathcal{D}, \otimes, I, a, l, r)$ be two tensor categories. A tensor functor from $\mathcal{C}$ to $\mathcal{D}$ is a triple (F, μ_0, μ) , where F is a functor from $\mathcal{C}$ to $\mathcal{D}$, μ_0 is an isomorphism from I to $F(I)$, and

$$\mu(U, V) : F(U) \otimes F(V) \rightarrow F(U \otimes V)$$

is a family of natural isomorphisms indexed by all couples (U, V) of objects of $\mathcal{C}$ such that

$$\begin{aligned} & \mu(U, V \otimes W)(id_{F(U)} \otimes \mu(V, W))a_{F(U), F(V), F(W)} \\ &= F(a_{U, V, W})\mu(U \otimes V, W)(\mu(U, V) \otimes id_{F(W)}), \\ & l_{F(U)} = F(l_U)\mu(I, U)(\varphi_0 \otimes id_{F(U)}) \end{aligned}$$

and

$$r_{F(U)} = F(r_U)\mu(U, I)(id_{F(U)} \otimes \mu_0)$$

for any $U, V, W \in ob\mathcal{C}$. Furthermore, if $(\mathcal{C}, C)$ and $(\mathcal{D}, C')$ be two braided tensor categories, and

$$F(C_{V, V'})\mu(F(V), F(V')) = \mu(F(V'), F(V))C_{F(V), F(V')}$$

for any $V, V' \in ob\mathcal{C}$, then (F, μ_0, μ) is called a braided tensor functor.

We now construct a strict tensor category $(\mathcal{C}^{str}, \underline{\otimes})$ by a general braided tensor category $(\mathcal{C}, \otimes, I, a, l, r, C)$ as follows. The objects of $\mathcal{C}^{str}$ consist of all finite sequences $S = (V_1, V_2, \dots, V_n)$ of $\mathcal{C}$, including the empty sequence $\emptyset$. The integer n is called the length of the sequence $S = (V_1, V_2, \dots, V_n)$. The length of empty sequence is zero. As in [64, P 288 – 291], let

$$\begin{aligned} F((V_1, V_2, \dots, V_n)) &= ((\dots (V_1 \otimes V_2) \otimes \dots) \otimes V_{n-1}) \otimes V_n, & F(\emptyset) &= I, \\ Hom_{\mathcal{C}^{str}}(S'', S''') &= Hom_{\mathcal{C}}(F(S''), F(S''')), \\ \mu(\emptyset, \emptyset) &= l_I, & \mu(S, \emptyset) &= r_{F(S)} & \mu(\emptyset, S) &= l_{F(S)}, \\ \mu(S, (V)) &= id_{F(S) \otimes V}, \\ \mu(S, S' \underline{\otimes} (V)) &= (\mu(S, S') \otimes id_V)a_{F(S), F(S'), V}^{-1} \end{aligned} \tag{1.1}$$

for $S, S', S'', S''' \in ob(\mathcal{C}^{str})$ with $S \neq \emptyset$ and $S' \neq \emptyset$. It is shown that the two tensor categories $\mathcal{C}^{str}$ and $\mathcal{C}$ are equivalent in [64, Proposition XI.5.1]. If $S = (V)$, then we write $S = V$ in short.

If, for any pair of objects (S, S') in $\mathcal{C}^{str}$, we define

$$F(\overline{C}_{S, S'}) = \mu(S', S)C_{F(S), F(S')}\mu(S, S')^{-1},$$

we shall show that $(\mathcal{C}^{str}, \overline{C})$ is a strict braided tensor category.

Definition 1.0.4. *let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a functor. Then F is an equivalence of categories if there exist a functor $G : \mathcal{D} \rightarrow \mathcal{C}$ and natural isomorphisms*

$$\eta : id_{\mathcal{D}} \rightarrow FG \quad \text{and} \quad \theta : GF \rightarrow id_{\mathcal{C}}.$$

we now give a useful criterion for a functor $F : \mathcal{C} \rightarrow \mathcal{D}$ to be equivalence of categories. let us first say that a functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is *essentially surjective* if, for any object W of $\mathcal{D}$, there exists an object V of $\mathcal{C}$ such that $F(V) \cong W$ in $\mathcal{D}$. It is said to be *faithful* (resp. fully faithful) if, for any couple (V, V') of object of $\mathcal{C}$, the map

$$F : Hom(V, V') \rightarrow Hom(F(V), F(V'))$$

on morphisms is injective (resp. bijective).

Proposition 1.0.5. *A functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is an equivalence of categories if and only if F is essentially surjective and fully faithful.*

Proof. (a) Suppose that F is an equivalence. Then there exist a functor $G : \mathcal{D} \rightarrow \mathcal{C}$ and natural isomorphisms $\eta : id_{\mathcal{D}} \rightarrow FG$ and $\theta : GF \rightarrow id_{\mathcal{C}}$. The first isomorphism shows that $W \cong F(G(W))$ for any object W of $\mathcal{C}$. In other words, F is essentially surjective. Now consider a morphism $f : V \rightarrow V'$ in $\mathcal{C}$. The square

$$\begin{array}{ccc} & \theta(V) & \\ GF(V) & \longrightarrow & V \\ \downarrow GF(f) & & \downarrow f \\ GF(V') & \longrightarrow & V' \\ & \theta(V') & \end{array}$$

commutes. It results that if $F(f) = F(f')$, hence $GF(f) = GF(f')$, then we have $f = f'$. Therefore, the functor F is faithful. Using the natural isomorphism η in a similar way, we prove that G is faithful too. Now consider a morphism $g : F(V) \rightarrow F(V')$. Let us show that $g = F(f)$ where $f = \theta(V') \circ G(g) \circ \theta(V)^{-1}$. Indeed,

$$\theta(V') \circ GF(g) \circ \theta(V)^{-1} = f = \theta(V') \circ G(g) \circ \theta(V)^{-1}.$$

Therefore $GF(f) = G(g)$. As G is faithful, we get $g = F(f)$. This proves that F is fully faithful.

(b) Let F is essentially surjective and fully faithful functor. For any object W in $\mathcal{D}$, we choose an object $G(W)$ of $\mathcal{C}$ and an isomorphism $\eta(W) : W \rightarrow FG(W)$ in $\mathcal{D}$. If $g : W \rightarrow W'$ is a morphism of $\mathcal{D}$. We may consider

$$\eta(W') \circ g \circ \eta(W)^{-1} : FG(W) \rightarrow FG(W').$$

Since F is fully faithful, there exists a unique morphism $G(g)$ from $G(W)$ to $G(W')$ such that

$$FG(g) = \eta(W') \circ g \circ \eta(W)^{-1} : FG(W) \rightarrow FG(W').$$

One checks easily that this define a functor G from $\mathcal{D}$ into $\mathcal{C}$ and that $\eta : id_{\mathcal{D}} \rightarrow FG$ is a natural isomorphism. In order to show that F and G are equivalences of categories, we have only to find a natural isomorphism $\theta : GF \rightarrow id_{\mathcal{C}}$. We define $\theta(V) : GF(V) \rightarrow V$ for any object $V \in Ob(\mathcal{C})$ as the unique morphism such that $F(\theta(V)) = \eta(F(V))^{-1}$. It is easily checked that this formula defines a natural isomorphism. $\square$

Proposition 1.0.6. *The categories $\mathcal{C}^{str}$ and $\mathcal{C}$ are equivalent.*

Proof. The map F defined above on objects of $\mathcal{C}^{str}$ extends to a functor $F : \mathcal{C}^{str} \rightarrow \mathcal{C}$ which is the identity on morphisms, hence fully faithful. As any object in $\mathcal{C}$ is clearly isomorphic to

the image under F of a sequence of length one, we see that F is essentially surjective. This proves the proposition in view of Proposition 1.0.5. Observe that $G(V) = V$ defines a functor $G : \mathcal{C} \rightarrow \mathcal{C}^{str}$ which is the inverse equivalence to F . Indeed, we have $FG = id_{\mathcal{C}}$. $\theta : GF \rightarrow id_{\mathcal{C}^{str}}$ via the natural isomorphism

$$\theta(S) = id_{F(S)} : GF(S) \rightarrow S. \square$$

Lemma 1.0.7. *If S, S', S'' are objects on $\mathcal{C}^{str}$, we have*

$$\mu(S, S' \underline{\otimes} S'') \circ (id_S \otimes \mu(S', S'')) \circ a_{F(S), F(S'), F(S'')} = \mu(S \underline{\otimes} S', S'') \circ (\mu(S, S') \otimes id_{S''}).$$

Proof. We proceed by induction on the length of S'' . If $S'' = \emptyset$, we have

$$\begin{aligned} \mu(S, S')(id_S \otimes \mu(S', \emptyset))a_{F(S), F(S'), I} &= \mu(S, S')(id_{F(S)} \otimes r_{F(S'')})a_{F(S), F(S'), I} \\ &= \mu(S, S')r_{F(S) \otimes F(S')} \\ &= r_{F(S \underline{\otimes} S')}(\mu(S, S') \otimes id_I) \\ &= \mu(S \underline{\otimes} S', \emptyset)(\mu(S, S') \otimes id_I). \end{aligned}$$

The first and last equalities are by definition, the second one by [64, Lemma XI.2.2] and the third one by naturality of r .

Let V be an object of the category. Let us prove that the equality of [64, Lemma XI. 5.2] for the triple (S, S', S'') implies the equality for $(S, S', S'' \underline{\otimes} (V))$. We have

$$\begin{aligned} &\mu(S, S' \underline{\otimes} S'' \underline{\otimes} (V))(id_S \otimes \mu(S', S'' \underline{\otimes} (V)))a_{F(S), F(S'), F(S'' \underline{\otimes} (V))} \\ &= (\mu(S, S' \underline{\otimes} S'') \otimes id_V)a_{F(S), F(S' \underline{\otimes} S''), V}^{-1}(id_S \otimes \mu(S', S'') \otimes id_V) \\ &\quad (id_S \otimes a_{F(S'), F(S''), V}^{-1})a_{F(S), F(S'), F(S'' \underline{\otimes} (V))} \\ &= (\mu(S, S' \underline{\otimes} S'') \otimes id_V)(id_S \otimes \mu(S', S'' \otimes id_V))a_{F(S), F(S') \underline{\otimes} F(S''), V}^{-1} \\ &\quad (id_S \otimes a_{F(S'), F(S''), V}^{-1})a_{F(S), F(S'), F(S'' \underline{\otimes} (V))} \\ &= (\mu(S, S' \underline{\otimes} S'') \otimes id_V)(id_S \otimes \mu(S', S'' \otimes id_V)) \\ &\quad (a_{F(S), F(S'), F(S'')} \otimes id_V)a_{F(S) \otimes F(S'), F(S''), V}^{-1} \\ &= (\mu(S \underline{\otimes} S', S'') \otimes id_V)(\mu(S, S') \otimes id_{S''} \otimes id_V)a_{F(S) \otimes F(S'), F(S''), V}^{-1} \\ &= (\mu(S \underline{\otimes} S', S'') \otimes id_V)a_{F(S) \otimes F(S \underline{\otimes} S'), F(S''), V}^{-1}(\mu(S, S') \otimes id_{S''} \otimes id_V) \\ &= (\mu(S \underline{\otimes} S', S'' \underline{\otimes} (V))(\mu(S, S') \otimes id_{S'' \underline{\otimes} (V)})). \end{aligned}$$

The first and last equalities follow from (1.1), the second and fifth ones from the naturality of the associativity constraint, the third from the Pentagon Axiom, and the fourth one from the induction hypothesis. $\square$

Theorem 1.0.8. *Equipped with this tensor product $\mathcal{C}^{str}$ is a strict tensor category. The categories $\mathcal{C}$ and $\mathcal{C}^{str}$ are tensor equivalent.*

Proof. It is easy to check that $\underline{\otimes}$ is a functor. This functor is strictly associative by construction. Therefore $\mathcal{C}^{str}$ is a strict tensor category.

In order to prove that it is tensor equivalent to $\mathcal{C}$, we have to exhibit tensor functors and natural isomorphisms. We first claim that the triple (F, id_I, μ) is a tensor functor from $\mathcal{C}^{str}$ to $\mathcal{C}$ where μ is the natural isomorphism defined above. By [64, Lemma XI.5.2] and [64, Proposition XI.5.2], the functor G of the proof of is a tensor functor. Finally, the natural isomorphism θ is a natural tensor isomorphism. $\square$

Theorem 1.0.9. *If $(\mathcal{C}, C)$ is a braided tensor category, then so is $(\mathcal{C}^{str}, \bar{C})$, and the two braided tensor categories are equivalent.*

Proof. We first show that

$$\bar{C}_{S, S' \underline{\otimes} S''} = (id_{S'} \underline{\otimes} \bar{C}_{S, S''})(\bar{C}_{S, S'} \underline{\otimes} id_{S''})$$

and

$$\bar{C}_{S \underline{\otimes} S', S''} = (\bar{C}_{S, S''} \underline{\otimes} id_{S'})(id_S \underline{\otimes} \bar{C}_{S', S''}).$$

We only check the first relation. The second one can similarly be shown.

We see that

$$\begin{aligned} F & \quad (\text{the right side of the first relation}) \\ &= \mu(S', S'' \underline{\otimes} S)(id_{F(S')} \otimes F(\bar{C}_{S, S''}))\mu(S', S \underline{\otimes} S'')^{-1} \\ & \quad \mu(S' \underline{\otimes} S, S'')(F(\bar{C}_{S, S'}) \otimes id_{F(S'')})\mu(S \underline{\otimes} S', S'')^{-1} \\ &= \mu(S', S'' \underline{\otimes} S)(id_{F(S')} \otimes \mu(S'', S)C_{F(S), F(S'')})\mu(S, S'')^{-1}\mu(S', S \underline{\otimes} S'')^{-1}\mu(S' \underline{\otimes} S, S'') \\ & \quad (\mu(S', S)C_{F(S), F(S')}\mu(S, S')^{-1} \otimes id_{F(S'')})\mu(S \underline{\otimes} S', S'')^{-1} \\ &= \mu(S', S'' \underline{\otimes} S)(id_{F(S')} \otimes \mu(S'', S))(id_{F(S')} \otimes C_{F(S), F(S'')})(id_{F(S')} \otimes \mu(S, S'')^{-1}) \\ & \quad \mu(S', S \underline{\otimes} S'')^{-1}\mu(S' \underline{\otimes} S, S'') \\ & \quad (\mu(S', S) \otimes id_{F(S'')})(C_{F(S), F(S')} \otimes id_{F(S'')})(\mu(S, S')^{-1} \otimes id_{F(S'')})\mu(S \underline{\otimes} S', S'')^{-1} \\ &= \mu(S' \underline{\otimes} S'', S)(\mu(S', S'') \otimes id_{F(S)})a_{F(S'), F(S''), F(S)}^{-1}(id_{F(S')} \otimes C_{F(S), F(S'')})a_{F(S'), F(S), F(S'')} \\ & \quad (C_{F(S), F(S')} \otimes id_{F(S'')})a_{F(S), F(S'), F(S'')}^{-1}(id_{F(S)} \otimes \mu(S', S'')^{-1})\mu(S, S' \underline{\otimes} S'')^{-1} \\ & \quad \text{by [64, lemma XI.5.2]} \\ &= \mu(S' \underline{\otimes} S'', S)(\mu(S', S'') \otimes id_{F(S)})C_{F(S), F(S') \otimes F(S'')}(id_{F(S)} \otimes \mu(S', S'')^{-1})\mu(S, S' \underline{\otimes} S'')^{-1} \\ &= \mu(S' \underline{\otimes} S'', S)C_{F(S), F(S' \underline{\otimes} S'')}\mu(S, S' \underline{\otimes} S'')^{-1} \\ &= F(\text{the left side of the first relation.}) \end{aligned}$$

Therefore the first relation holds.

Next we can obtain that $\bar{C}$ is natural by the diagram below:

$$\begin{array}{ccccc}
& & C_{F(S),F(S')} & & \\
& & \longrightarrow & & \\
F(S) \otimes F(S') & & & & F(S') \otimes F(S) \\
& \searrow \mu(S, S') & & \swarrow \mu(S', S) & \\
& F(S \underline{\otimes} S') & \xrightarrow{F(\bar{C}_{S,S'})} & F(S' \underline{\otimes} S) & \\
f \otimes g \downarrow & \downarrow f \underline{\otimes} g & & \downarrow g \underline{\otimes} f & \\
& F(T \underline{\otimes} T') & \xrightarrow{F(\bar{C}_{T,T'})} & F(T' \underline{\otimes} T) & \\
& \nearrow \mu(T, T') & & \nwarrow \mu(T', T) & \\
F(T) \otimes F(T') & & \longrightarrow & & F(T') \otimes F(T) \\
& & C_{F(T),F(T')} & &
\end{array}$$

Finally, we can easily check that F is a braided tensor functor and $(\mathcal{C}^{str}, \bar{C})$ and $(\mathcal{C}, C)$ are braided tensor equivalent categories. $\square$

Therefore, we can view every braided tensor category as a strict braided tensor category and use braiding diagrams freely (we can also see [64, exercise 5 P337]).

Example 1.0.10. (*The tensor category of vector spaces*) The most fundamental example of a tensor category is given by the category $\mathcal{C} = \text{Vect}(k)$ of vector spaces over field k . $\text{Vect}(k)$ is equipped with tensor structure for which $\otimes$ is the tensor product of the vector spaces over k , the unit object I is the ground field k itself, and the associativity constraint and unit constraint are the natural isomorphisms

$$a_{U,V,W}((u \otimes v) \otimes w) = u \otimes (v \otimes w) \quad \text{and} \quad l_V(1 \otimes v) = v = r_V(v \otimes 1)$$

for any vector space U, V, W and $u \in U, v \in V, w \in W$.

Furthermore, the most fundamental example of a braided tensor category is given by the tensor category $\text{Vect}(k)$, whose braiding is usual twist map from $U \otimes V$ to $V \otimes U$ defined by sending $a \otimes b$ to $b \otimes a$ for any $a \in U, b \in V$.

Now we define some notations. If f is a morphism from U to V and g is a morphism from V to W , we denote the composition gf by

$$\begin{array}{c} U \\ | \\ \textcircled{gf} \\ | \\ W \end{array} = \begin{array}{c} U \\ | \\ \textcircled{f} \\ | \\ \textcircled{g} \\ | \\ W \end{array} .$$

We usually omit I and the morphism id_I in any diagrams. In particular, If f is a morphism from I to V , g is a morphism from V to I , we denote f and g by

$$\begin{array}{c} \textcircled{f} \\ | \\ V \end{array} , \begin{array}{c} V \\ | \\ \textcircled{g} \end{array} .$$

If f is a morphism from $U \otimes V$ to P , g is a morphism from $U \otimes V$ to I and ζ is a morphism from $U \otimes V$ to $V \otimes U$, we denote f , g and ζ by

$$\begin{array}{c} U \quad V \\ | \quad | \\ \textcircled{f} \\ | \\ P \end{array}, \quad \begin{array}{c} U \quad V \\ | \quad | \\ \textcircled{g} \\ | \\ I \end{array}, \quad \begin{array}{c} U \quad V \\ \diagdown \quad \diagup \\ \textcircled{\zeta} \\ \diagup \quad \diagdown \\ V \quad U \end{array}.$$

In particular, we denote the braiding $C_{U,V}$ and its inverse $C_{U,V}^{-1}$ by

$$\begin{array}{c} U \quad V \\ \diagdown \quad \diagup \\ V \quad U \end{array}, \quad \begin{array}{c} V \quad U \\ \diagdown \quad \diagup \\ U \quad V \end{array}.$$

ξ is called an R -matrix of $\mathcal{C}$ if ξ is a natural isomorphism from $\otimes$ to $\otimes \tau$ and for any $U, V, W \in \text{ob}\mathcal{C}$, the Yang-Baxter equation of $\mathcal{C}$:

(YBE):

$$\begin{array}{c} U \quad V \quad W \\ \diagdown \quad \diagup \quad | \\ \textcircled{\xi} \\ | \quad | \quad | \\ \textcircled{\xi} \\ | \quad | \quad | \\ \textcircled{\xi} \\ \diagup \quad \diagdown \quad | \\ W \quad V \quad U \end{array} = \begin{array}{c} U \quad V \quad W \\ | \quad | \quad \diagdown \quad \diagup \\ \textcircled{\xi} \\ | \quad | \quad | \\ \textcircled{\xi} \\ | \quad | \quad | \\ \textcircled{\xi} \\ \diagup \quad \diagdown \quad | \\ W \quad V \quad U \end{array}$$

holds. In particular, the above equation is called the Yang-Baxter equation on V when $U = V = W$.

Lemma 1.0.11. (i) *The braiding C of braided tensor category $(\mathcal{C}, C)$ is an R -matrix of $\mathcal{C}$, i.e.*

$$\begin{array}{c} U \quad V \quad W \\ \diagdown \quad \diagup \quad | \\ | \quad | \quad | \\ \diagup \quad \diagdown \quad | \\ W \quad V \quad U \end{array} = \begin{array}{c} U \quad V \quad W \\ | \quad | \quad \diagdown \quad \diagup \\ | \quad | \quad | \\ \diagup \quad \diagdown \quad | \\ W \quad V \quad U \end{array}$$

holds.

(ii)

$$\begin{array}{c} U \quad V \quad W \\ | \quad | \quad \diagdown \quad \diagup \\ | \quad | \quad | \\ \textcircled{f} \\ | \\ P \end{array} = \begin{array}{c} U \quad VW \\ | \quad | \\ \textcircled{f} \\ | \\ W \quad P \end{array}, \quad \begin{array}{c} U \quad W \quad V \\ | \quad | \quad \diagdown \quad \diagup \\ | \quad | \quad | \\ \textcircled{f} \\ | \\ P \end{array} = \begin{array}{c} U \quad WV \\ | \quad | \\ \textcircled{f} \\ | \\ W \quad P \end{array}.$$

Proof. (i)

$$\begin{array}{c}
\begin{array}{c} U & V & W \\ \diagdown & & \diagup \\ W & V & U \end{array} = \begin{array}{c} U & V & W \\ | & | & | \\ \textcircled{C_{V,W}} & & | \\ | & | & | \\ \textcircled{C_{U,W}} & & | \\ | & | & | \\ W & V & U \\ | & | & | \\ U \otimes V & & W \\ | & | & | \\ \textcircled{C_{U \otimes V, W}} & & | \\ | & | & | \\ W & V \otimes U \end{array} \begin{array}{c} \text{by Hexagon Axiom} \\ \hline \end{array} \begin{array}{c} U \otimes V & W \\ | & | \\ \textcircled{C_{V,W}} & \\ | & | \\ \textcircled{C_{V \otimes U, W}} & \\ | & | \\ W & V \otimes U \end{array} \\
\begin{array}{c} \text{by naturality} \\ \hline \end{array} \begin{array}{c} U \otimes V & W \\ | & | \\ \textcircled{C_{U \otimes V, W}} & \\ | & | \\ \textcircled{C_{U,V}} & \\ | & | \\ W & V \otimes U \end{array} \begin{array}{c} \text{by Hexagon Axiom} \\ \hline \end{array} \begin{array}{c} U & V & W \\ | & | & | \\ \diagdown & & \diagup \\ W & V & U \end{array} .
\end{array}$$

(ii) We have

$$\begin{array}{c}
\begin{array}{c} U & VW \\ | & | \\ \textcircled{f} & \\ | & | \\ W & P \end{array} = \begin{array}{c} U \otimes VW \\ | & | \\ \textcircled{f} & \\ | & | \\ W & P \end{array} \begin{array}{c} \text{by naturality} \\ \hline \end{array} \begin{array}{c} U \otimes VW \\ | & | \\ \textcircled{f} & \\ | & | \\ W & P \end{array} \\
\begin{array}{c} \text{by Hexagon Axiom} \\ \hline \end{array} \begin{array}{c} U & V & W \\ | & | & | \\ \diagdown & & \diagup \\ W & P & \end{array} .
\end{array}$$

Similarly, we can show the second equation. $\square$

Now we give some concepts as follows: Assume that $H, A \in ob \mathcal{C}$, and

$$\begin{aligned}
\alpha : H \otimes A &\rightarrow A & , & & \beta : H \otimes A &\rightarrow H, \\
\phi : A &\rightarrow H \otimes A & , & & \psi : H &\rightarrow H \otimes A, \\
m_H : H \otimes H &\rightarrow H & , & & m_A : A \otimes A &\rightarrow A, \\
\Delta_H : H &\rightarrow H \otimes H & , & & \Delta_A : A &\rightarrow A \otimes A, \\
\eta_H : I &\rightarrow H & , & & \eta_A : I &\rightarrow A, \\
\epsilon_H : H &\rightarrow I & , & & \epsilon_A : A &\rightarrow I.
\end{aligned}$$

are morphisms in $\mathcal{C}$.

(A, m_A, η_A) is called an algebra living in $\mathcal{C}$, if the following conditions are satisfied:

$$\begin{array}{c} A & A & A \\ | & | & | \\ \textcircled{m} & & \\ | & & \\ \textcircled{m} & & \\ | & & \\ A & & \end{array} =
 \begin{array}{c} A & A & A \\ | & | & | \\ \textcircled{m} & & \\ | & & \\ \textcircled{m} & & \\ | & & \\ A & & \end{array}, \quad
 \begin{array}{c} & & A \\ & & | \\ \textcircled{\eta_A} & & \\ | & & \\ \textcircled{m} & & \\ | & & \\ A & & \end{array} =
 \begin{array}{c} A & & \textcircled{\eta_A} \\ | & & | \\ & & \textcircled{m} \\ & & | \\ & & A \end{array} =
 \begin{array}{c} A \\ | \\ A \end{array}.$$

In this case, η_A and m_A are called unit and multiplication of A respectively.

$(H, \Delta_H, \epsilon_H)$ is called a coalgebra living in $\mathcal{C}$, if the following conditions are satisfied:

$$\begin{array}{c} H \\ \downarrow \\ \text{---} \Delta \text{---} \\ | \quad | \\ H H \quad H \end{array} = \begin{array}{c} H \\ \downarrow \\ \text{---} \Delta \text{---} \\ | \quad | \\ H \quad H H \end{array}, \quad \begin{array}{c} H \\ \downarrow \\ \text{---} \Delta \text{---} \\ | \quad | \\ \varepsilon_H \quad H \end{array} = \begin{array}{c} H \\ \downarrow \\ \text{---} \Delta \text{---} \\ | \quad | \\ H \quad \varepsilon_H \end{array} = \begin{array}{c} H \\ | \\ H \end{array}.$$

In this case, ϵ_H and Δ_H are called counit and comultiplication of H respectively.

If A is an algebra and H is a coalgebra, then $Hom_{\mathcal{C}}(H, A)$ becomes an algebra under the convolution product

and its unit element $\eta = \eta_A \epsilon_H$. If S is the inverse of id_H in $Hom_{\mathcal{C}}(H, H)$, then S is called antipode of H .

If (H, m_H, η_H) is an algebra, and $(H, \Delta_H, \epsilon_H)$ is a coalgebra living in $\mathcal{C}$, and the following condition is satisfied:

$$\begin{array}{c} H & H \\ \diagup & \diagdown \\ \text{---} m \text{---} \\ \diagdown & \diagup \\ H & H \end{array} = \begin{array}{c} H & H \\ \diagup & \diagdown \\ \Delta & \Delta \\ \diagdown & \diagup \\ m & m \\ \diagup & \diagdown \\ H & H \end{array}, \quad \begin{array}{c} H & H \\ \diagup & \diagdown \\ \text{---} m \text{---} \\ \diagdown & \diagup \\ \varepsilon & \varepsilon \end{array} = \begin{array}{c} H & H \\ | & | \\ \varepsilon & \varepsilon \end{array}, \quad \begin{array}{c} \eta \\ | \\ \Delta \end{array} = \begin{array}{c} \eta \\ | \\ \eta \end{array}, \quad \begin{array}{c} \eta \\ | \\ \varepsilon \end{array} = \begin{array}{c} I \\ | \\ I \end{array}.$$

then H is called a bialgebra living in $\mathcal{C}$. If H is a bialgebra and there is an inverse S of id_H under convolution product in $Hom_{\mathcal{C}}(H, H)$, then H is called a Hopf algebra living in $\mathcal{C}$, or a braided Hopf algebra.

For $V \in \text{object } \mathcal{C}$, if there exists an object U and morphisms :

$$d_V : U \otimes V \rightarrow I \quad \text{and} \quad b_V : I \rightarrow V \otimes U$$

in $\mathcal{C}$ such that

$$(d_V \otimes id_U)(id_U \otimes b_V) = id_U \quad \text{and} \quad (id_V \otimes d_V)(b_V \otimes id_V) = id_V,$$


then U is called a left duality of V , written as V^* . In this case, d_V and b_V are called evaluation morphism and coevaluation morphism of V respectively.

If U and V have left duality U^* and V^* , respectively, then $V^* \otimes U^*$ is a left duality of $U \otimes V$. Its evaluation and coevaluation are

$$d_{U \otimes V} = d_V(id_{V^*} \otimes d_U \otimes id_V), \quad b_{U \otimes V} = (id_U \otimes b_V \otimes id_{U^*})b_U,$$


respectively.


The multiplication, comultiplication, evaluation morphism and coevaluation are usually denoted by




in short, respectively.


Let us first define the transpose $f^* : V^* \rightarrow U^*$ of morphism $f : U \rightarrow V$ by $f^* = (d \otimes id_{U^*})(id_{V^*} \otimes f \otimes id_{U^*})(id_{V^*} \otimes b)$.

Lemma 1.0.12. Let $\xi =$  be a morphism from $P \otimes Q$ to $Y \otimes Z$ and all objects

concerned have left duals. Then $\xi^* =$ 

Proof. We see that





Proposition 1.0.13. (cf. [82, Lemma 2.3, Pro. 2.4]) (i) If H is a Hopf algebra living in $\mathcal{C}$, then $Sm = m_{C_{H,H}}(S \otimes S)$ and $\Delta S = C_{H,H}(S \otimes S)\Delta$.

(ii) If $(H, \Delta_H, \epsilon_H, m_H, \eta_H)$ is a bialgebra or a Hopf algebra living in $\mathcal{C}$ and has a left duality H^* in $\mathcal{C}$, then $(H^*, (\Delta_H)^*, (\epsilon_H)^*, (m_H)^*, (\eta_H)^*)$ is a bialgebra or a Hopf algebra in $\mathcal{C}$.

Proof. (i)



$$\begin{aligned}
Sm &= \begin{array}{c} \begin{array}{cc} H & H \\ | & | \\ \text{---} \diagdown & \diagup \text{---} \\ | & | \\ \text{---} \diagup & \diagdown \text{---} \\ | & | \\ S & S \\ | & | \\ H & H \end{array} \\ = & \begin{array}{c} \begin{array}{cc} H & H \\ | & | \\ \text{---} \diagdown & \diagup \text{---} \\ | & | \\ \text{---} \diagup & \diagdown \text{---} \\ | & | \\ S & S \\ | & | \\ H & H \end{array} \\ = & \begin{array}{c} \begin{array}{cc} H & H \\ | & | \\ \text{---} \diagdown & \diagup \text{---} \\ | & | \\ \text{---} \diagup & \diagdown \text{---} \\ | & | \\ S & S \\ | & | \\ H & H \end{array} \\ = & mC_{H,H}(S \otimes S). \end{aligned}$$


Similarly, we can show $\Delta S = C_{H,H}(S \otimes S)\Delta$.

(ii) It is clear that H^* is not only an algebra but also a coalgebra if we set $\Delta_{H^*} = (m_H)^*$, $\epsilon_{H^*} = (\eta_H)^*$ and $m_{H^*} = (\Delta_H)^*$.

Now we show that

$$\begin{array}{c} \begin{array}{cc} H^* & H^* \\ | & | \\ \text{---} \diagdown & \diagup \text{---} \\ | & | \\ \text{---} \diagup & \diagdown \text{---} \\ | & | \\ H^* & H^* \end{array} \\ = & \begin{array}{c} \begin{array}{cc} H^* & H^* \\ | & | \\ \text{---} \diagdown & \diagup \text{---} \\ | & | \\ \text{---} \diagup & \diagdown \text{---} \\ | & | \\ H^* & H^* \end{array} \\ = & \begin{array}{c} \begin{array}{cc} H^* & H^* \\ | & | \\ \text{---} \diagdown & \diagup \text{---} \\ | & | \\ \text{---} \diagup & \diagdown \text{---} \\ | & | \\ H^* & H^* \end{array} \end{array}$$

The left side =  = 

=  by Lem.1.0.12 = the right side .

Thus H^* is a bialgebra living in $\mathcal{C}$. If H is a Hopf algebra, then H^* is also a Hopf algebra with antipode $(S_H)^*$. $\square$

If H is an algebra and the following conditions are satisfied:

$$\begin{array}{c} HH \quad A \\ \downarrow \quad \downarrow \\ \alpha \quad \alpha \\ \downarrow \quad \downarrow \\ H \end{array} = \begin{array}{c} H \quad HA \\ \downarrow \quad \downarrow \\ m \quad \alpha \\ \downarrow \quad \downarrow \\ H \end{array}, \quad \begin{array}{c} A \\ \downarrow \\ \eta_H \\ \downarrow \\ \alpha \\ \downarrow \\ H \end{array} = \begin{array}{c} A \\ \downarrow \\ A \end{array}$$

then (A, α) is called a braided H -module or an H -module living in $\mathcal{C}$.

If H is a coalgebra and the following conditions are satisfied:

$$\begin{array}{c} A \\ \downarrow \\ \phi \\ \downarrow \\ HH \quad A \end{array} = \begin{array}{c} A \\ \downarrow \\ \phi \\ \downarrow \\ H \quad HA \end{array}, \quad \begin{array}{c} A \\ \downarrow \\ \phi \\ \downarrow \\ \varepsilon_H \\ \downarrow \\ A \end{array} = \begin{array}{c} A \\ \downarrow \\ A \end{array},$$

then (A, ϕ) is called a braided H -comodule or an H -comodule living in $\mathcal{C}$. Let ${}_H\mathcal{M}(\mathcal{C})$ and ${}^H\mathcal{M}(\mathcal{C})$ denote the categories of H -modules in $\mathcal{C}$ and H -comodules in $\mathcal{C}$, called a braided H -module category and a braided H -comodule category, respectively. Let ${}_H\mathcal{M}(\mathcal{C}) = {}_H\mathcal{M}$ and ${}^H\mathcal{M}(\mathcal{C}) = {}^H\mathcal{M}$ when $\mathcal{C} = \mathcal{V}ect(k)$ in Example 1.0.10.

Similarly, we can define the concepts of algebra morphism, coalgebra morphism and H -module morphism. An algebra (A, α) is called an H -module algebra, if (A, α) is an H -module

and its unit η_A and multiplication m_A are H -module morphisms. Similarly, we can define the concepts of H -module coalgebra, H -comodule algebra and H -comodule coalgebra.

Assume that $\mathcal{C}$ is a concrete braided tensor category. If H is a braided bialgebra and M is an H -module in $\mathcal{C}$, then

$$M^H := \{x \in M \mid hx = \epsilon(h)x, \forall h \in H\}$$

is called the invariants of M . If M is a regular H -module (i.e. the module operation is m), then M^H is called the integral of H and is written as $\int_H^l$.

A bialgebra $(H, m, \eta, \Delta, \epsilon)$ with convolution-invertible R in $\text{Hom}_{\mathcal{C}}(I, H \otimes H)$ is called a quasitriangular bialgebra living in braided tensor category $\mathcal{C}$ if there is a morphism $\bar{\Delta} : H \rightarrow H \otimes H$ such that $(H, \bar{\Delta}, \epsilon)$ is a coalgebra and

(QT1):

(QT2):

(QT3):

In this case, we also say that $(H, R, \bar{\Delta})$ is a braided quasitriangular bialgebra in $\mathcal{C}$.

A bialgebra $(H, m, \eta, \Delta, \epsilon)$ with convolution-invertible r in $\text{Hom}_{\mathcal{C}}(H \otimes H, I)$ is called a coquasitriangular bialgebra living in braided tensor category $\mathcal{C}$ if there exists a morphism $\bar{m} : H \otimes H \rightarrow H$ such that $(H, \bar{m}, \eta)$ is an algebra and

(CQT1):

(CQT2):

(CQT3):

In this case, we also say that $(H, r, \bar{m})$ is a braided coquasitriangular bialgebra in $\mathcal{C}$.

A braided Hopf algebra H is called a braided group if there exists a braided quasitriangular structure $(H, R, \bar{\Delta})$ with $\bar{\Delta} = \Delta_H$ or there exists a coquasitriangular structure $(H, r, \bar{m})$ with $m = \bar{m}$.

Let m^{op} and Δ^{cop} denote $m_H C_{H,H}^{-1}$ and $C_{H,H}^{-1} \Delta$ respectively.

In particular, we say that (H, R) is quasitriangular if (H, R, Δ^{cop}) is quasitriangular. Dually, we say that (H, r) is quasitriangular if (H, r, m^{op}) is coquasitriangular.

Let $(H, R, \bar{\Delta})$ be a quasitriangular bialgebra living in braided tensor category $\mathcal{C}$. (V, α) is said to be symmetric with respect to $(H, R, \bar{\Delta})$, if (V, α) is a left H -module and satisfies the following condition:

(SWRT) :

Define

$$O(H, \bar{\Delta}) = \{V \in {}_H\mathcal{M} \mid (V, \alpha) \text{ is symmetric with respect to } (H, R, \bar{\Delta})\}.$$

Let $(H, r, \bar{m})$ be a coquasitriangular bialgebra living in braided tensor category. (V, ϕ) is said to be symmetric with respect to $(H, r, \bar{m})$ if (V, ϕ) is a left H -comodule and satisfies the following condition:

(SWRT) :

Define

$$O(H, \bar{m}) = \{V \in {}^H\mathcal{M} \mid (V, \phi) \text{ is symmetric with respect to } (H, r, \bar{m})\}.$$

Proposition 1.0.14. (i) If $(H, R, \bar{\Delta})$ is a quasitriangular bialgebra living in braided tensor category $(\mathcal{C}, \otimes, I, a, l, r, C)$, then $(O(H, \bar{\Delta}), \otimes, I, a, l, r, C^R)$ is a braided tensor category, where

$$C_{V,W}^R = \begin{array}{c} \begin{array}{cc} & V & W \\ & | & | \\ \textcircled{R} & & \\ & \diagdown & \diagup \\ \textcircled{\alpha_V} & & \textcircled{\alpha_W} \\ & \diagup & \diagdown \\ & W & V \end{array} \end{array} \quad \text{and} \quad \alpha_{V \otimes W} = \begin{array}{c} \begin{array}{ccc} & H & V & W \\ & | & | & | \\ & \text{---} & & \\ & \text{---} & & \\ & \text{---} & & \\ \textcircled{\alpha_V} & & \textcircled{\alpha_W} \\ & | & | & | \\ V & & W \end{array} \end{array}$$

for any $(V, \alpha_V), (W, \alpha_W) \in O(H, \bar{\Delta})$

(ii) If $(H, r, \bar{m})$ is a coquasitriangular bialgebra living in braided tensor category $(\mathcal{C}, \otimes, I, a, l, r)$, then $(O(H, \bar{m}), \otimes, I, a, l, r, C^r)$ is a braided tensor category, where

$$C_{V,W}^r = \begin{array}{c} \begin{array}{cc} & V & W \\ & | & | \\ & \diagdown & \diagup \\ \textcircled{\phi_W} & & \textcircled{\phi_V} \\ & \diagup & \diagdown \\ \textcircled{r} & & \\ & | & | \\ W & & V \end{array} \end{array} \quad \text{and} \quad \phi_{V \otimes W} = \begin{array}{c} \begin{array}{ccc} & V & W \\ & | & | \\ \textcircled{\phi_V} & & \textcircled{\phi_W} \\ & \diagdown & \diagup \\ & H & V & W \\ & | & | & | \\ & \text{---} & & \\ & \text{---} & & \\ & \text{---} & & \end{array} \end{array}$$

for any $(V, \phi_V), (W, \phi_W) \in O(H, \bar{m})$.

Proof. We only show part (i) since part (ii) can be shown similarly. We first show that $O(H, \bar{\Delta})$ is a tensor category. That is, we need show that $V \otimes W \in O(H, \bar{\Delta})$ for any $V, W \in O(H, \bar{\Delta})$. In fact,

$$\begin{array}{c} \begin{array}{ccc} & H & V \otimes W \\ & | & | \\ & \text{---} & \\ & \text{---} & \\ & \text{---} & \\ H & & V \otimes W \\ & | & | \\ & \text{---} & \\ & \text{---} & \\ & \text{---} & \end{array} \end{array} = \begin{array}{c} \begin{array}{ccc} & H & V & W \\ & | & | & | \\ & \text{---} & & \\ & \text{---} & & \\ & \text{---} & & \\ H & & \textcircled{\alpha} & \textcircled{\alpha} \\ & | & | & | \\ H & & V & W \end{array} \end{array} = \begin{array}{c} \begin{array}{ccc} & H & V & W \\ & | & | & | \\ & \text{---} & & \\ & \text{---} & & \\ & \text{---} & & \\ H & & \textcircled{\alpha} & \textcircled{\alpha} \\ & | & | & | \\ H & & V & W \end{array} \end{array}$$

$$\begin{array}{c}
\begin{array}{ccc} H & V & W \\ \downarrow & & \\ \text{Diagram 1} \\ \downarrow & & \\ \text{Diagram 2} \\ \downarrow & & \\ H & V & W \end{array} & \stackrel{by(SWRT)}{=} & \begin{array}{ccc} H & V & W \\ \downarrow & & \\ \text{Diagram 3} \\ \downarrow & & \\ \text{Diagram 4} \\ \downarrow & & \\ H & V & W \end{array} \\
\\
\begin{array}{ccc} H & & \\ \downarrow & & \\ \text{Diagram 5} \\ \downarrow & & \\ \text{Diagram 6} \\ \downarrow & & \\ H & V & W \end{array} & \stackrel{by(SWRT)}{=} & \begin{array}{ccc} H & V \otimes W & \\ \downarrow & & \\ \text{Diagram 7} \\ \downarrow & & \\ \text{Diagram 8} \\ \downarrow & & \\ HV \otimes W & & \end{array}
\end{array}$$

Therefore $O(H, \bar{\Delta})$ is a tensor subcategory of $\mathcal{C}$. Next we show that C^R is a braiding in $O(H, \bar{\Delta})$.

It is clear that the inverse of $C_{V,W}^R$ is

$$\begin{array}{ccc} & W & V \\ & \downarrow & \downarrow \\ \text{Diagram 9} \\ \downarrow & & \\ V & & W \end{array}$$


We see that

$$\begin{array}{c}
 \begin{array}{c} H \quad V \otimes W \\ | \quad | \\ \text{---} \alpha \text{---} \\ | \\ W \otimes V \end{array} = \begin{array}{c} H \quad V \quad W \\ | \quad | \quad | \\ \text{---} \alpha \text{---} \\ | \quad | \\ W \quad V \end{array} \stackrel{\text{by}(SWRT)}{=} \begin{array}{c} H \quad VW \\ | \quad | \\ \text{---} \alpha \text{---} \\ | \quad | \\ W \quad V \end{array} \\
 \\
 \begin{array}{c} H \\ | \\ \text{---} \alpha \text{---} \\ | \quad | \\ W \quad V \end{array} \stackrel{\text{by}(QT3)}{=} \begin{array}{c} H \\ | \\ \text{---} \alpha \text{---} \\ | \quad | \\ W \quad V \end{array} = \begin{array}{c} H \quad V \otimes W \\ | \quad | \\ \text{---} \alpha \text{---} \\ | \\ W \otimes V \end{array} .
 \end{array}$$

Thus $C_{V,W}^R$ is an H -module morphism.

We also see that

$$\begin{array}{c}
 C_{V \otimes W, X}^R = \begin{array}{c} \begin{array}{c} R \\ | \\ \text{---} \alpha \text{---} \\ | \quad | \quad | \\ X \quad V \quad W \end{array} \end{array} = \begin{array}{c} \begin{array}{c} R \\ | \\ \text{---} \alpha \text{---} \\ | \quad | \quad | \\ X \quad V \quad W \end{array} \end{array}
 \end{array}$$



$= (C_{V,X}^R \otimes id_W)(id_V \otimes C_{W,X}^R)$. Thus

$$C_{V \otimes W, X}^R = (C_{V,X}^R \otimes id_W)(id_V \otimes C_{W,X}^R).$$

Similarly, we have

$$C_{V,W \otimes X} = (id_W \otimes C_{V,X}^R)(C_{V,W}^R \otimes id_X).$$

Therefore $O(H, \bar{\Delta})$ is a braided tensor category. $\square$

It is clear that $O(H, \Delta_H^{cop}) = {}_H\mathcal{M}(\mathcal{C})$ and $O(H, m_H^{op}) = {}^H\mathcal{M}(\mathcal{C})$ when $(\mathcal{C}, C)$ is a symmetric braided tensor category.

Corollary 1.0.15. *Let $(\mathcal{C}, \otimes, I, a, l, r, C)$ be a symmetric braided tensor category.*

(i) *If (H, R) is a quasitriangular bialgebra in $\mathcal{C}$, then $({}_H\mathcal{M}, \otimes, I, a, l, r, C^R)$ is a braided tensor category.*

(ii) *If (H, r) is a coquasitriangular bialgebra in $\mathcal{C}$, then $({}^H\mathcal{M}, \otimes, I, a, l, r, C^r)$ is a braided tensor category.*

Let H be a bialgebra algebra in braided tensor category $(\mathcal{C}, C)$, and (M, α, ϕ) a left H -module and left H -comodule in $(\mathcal{C}, C)$. If

$$(YD) : \quad \begin{array}{c} \begin{array}{c} H \\ \downarrow \\ \text{cap} \\ \downarrow \\ \alpha \\ \downarrow \\ \phi \\ \downarrow \\ \text{cup} \\ \downarrow \\ H \end{array} \quad \begin{array}{c} A \\ \downarrow \\ \text{cap} \\ \downarrow \\ \phi \\ \downarrow \\ \alpha \\ \downarrow \\ \text{cup} \\ \downarrow \\ A \end{array} \\ \downarrow \\ \begin{array}{c} H \\ \downarrow \\ \text{cap} \\ \downarrow \\ \phi \\ \downarrow \\ \alpha \\ \downarrow \\ \text{cup} \\ \downarrow \\ H \end{array} \quad \begin{array}{c} A \\ \downarrow \\ \text{cap} \\ \downarrow \\ \phi \\ \downarrow \\ \alpha \\ \downarrow \\ \text{cup} \\ \downarrow \\ A \end{array} \end{array} = \begin{array}{c} \begin{array}{c} H \\ \downarrow \\ \text{cap} \\ \downarrow \\ H \end{array} \quad \begin{array}{c} A \\ \downarrow \\ \phi \\ \downarrow \\ \alpha \\ \downarrow \\ A \end{array} \\ \downarrow \\ \begin{array}{c} H \\ \downarrow \\ \text{cap} \\ \downarrow \\ H \end{array} \quad \begin{array}{c} A \\ \downarrow \\ \phi \\ \downarrow \\ \alpha \\ \downarrow \\ A \end{array} \end{array} .$$

then (M, α, ϕ) is called a Yetter-Drinfeld H -module in $\mathcal{C}$, written as YD H -module in short. When H is a Hopf algebra algebra in braided tensor category $(\mathcal{C}, C)$, then the condition above is equivalent to

(YD):

$$\phi\alpha = \begin{array}{c} \begin{array}{c} H \\ \downarrow \\ \text{cap} \\ \downarrow \\ \phi \\ \downarrow \\ \alpha \\ \downarrow \\ \text{cup} \\ \downarrow \\ H \end{array} \quad \begin{array}{c} M \\ \downarrow \\ \phi \\ \downarrow \\ \alpha \\ \downarrow \\ M \end{array} \\ \downarrow \\ \begin{array}{c} H \\ \downarrow \\ \text{cap} \\ \downarrow \\ H \end{array} \quad \begin{array}{c} M \\ \downarrow \\ \phi \\ \downarrow \\ M \end{array} \end{array} .$$

Let ${}^H_H\mathcal{YD}(\mathcal{C})$ denote the category of all Yetter-Drinfeld H -modules in $\mathcal{C}$. If $(\mathcal{C}, C) = \mathcal{Vect}(k)$, we write ${}^H_H\mathcal{YD}(\mathcal{C}) = {}^H_H\mathcal{YD}$, called Yetter-Drinfeld category. It follows from [107] and [11, Theorem 4.1.1] that ${}^H_H\mathcal{YD}(\mathcal{C})$ is a braided tensor category with ${}^{YD}C_{U,V} = (\alpha_V \otimes id_U)(id_H \otimes C_{U,V})(\phi_U \otimes id_V)$ for any two Yetter-Drinfeld modules (U, ϕ_U, α_U) and (V, ϕ_V, α_V) when H has an invertible

antipode. In this case, ${}^{YD}C_{U,V}^{-1} = (id_V \otimes \alpha_U)(C_{H,V}^{-1} \otimes id_U)(S^{-1} \otimes C_{U,V}^{-1})(C_{U,H}^{-1} \otimes id_V)(id_U \otimes \phi_V)$. Algebras, coalgebras and Hopf algebras and so on in ${}^H_H\mathcal{YD}$ are called Yetter-Drinfeld ones or YD ones in short.

The structure of a Nichols algebra appeared first in the paper [91] and was rediscovered later by several authors.

Let V be a YD H -module in $\mathcal{Vect}(k)$. A graded braided Hopf algebra $R = \bigoplus_{n \geq 0} R(n)$ in ${}^H_H\mathcal{YD}$ is called a *Nichols algebra* of V if the following are satisfied:

- $k \cong R(0)$,
- $V \cong R(1)$ in ${}^H_H\mathcal{YD}$,
- $R(1) = P(R)$, the vector space of the primitive elements of R ,
- R is generated as an algebra by $R(1)$.


Remark ¹

Example 1.0.16. Let V be a finite-dimensional vector space over field k and $V^* = \text{Hom}_k(V, k)$. Define maps $b_V : k \rightarrow V \otimes V^*$ and $d_V : V^* \otimes V \rightarrow k$ by

$$b_V(1) = \sum_i v_i \otimes v^i \quad \text{and} \quad \sum_{i,j} d_V(v^i \otimes v_j) = v^i(v_j)$$

where $\{v_i \mid i = 1, 2, \dots, n\}$ is any basis of V and $\{v^i \mid i = 1, 2, \dots, n\}$ is the dual basis in V^* . It is clear that b_V and d_V are evaluation and coevaluation of V in $\mathcal{Vect}(k)$, respectively.

Proposition 1.0.17. (i) Let $(\mathcal{C}, C)$ be a braided tensor category and H be an algebra living in $\mathcal{C}$. If (V, α) is a left H -module and has a left duality V^* in $\mathcal{C}$, then V^* is a left duality of V in ${}_H\mathcal{M}(\mathcal{C})$. Here the module operation α_{V^*} of V^* is defined as follows:



(ii) Let $(\mathcal{C}, C)$ be a braided tensor category and H be a coalgebra living in $(\mathcal{C}, C)$. If (V, ψ) is a right H -comodule and has a left duality V^* in $(\mathcal{C}, C)$, then V^* is a left duality of V in $\mathcal{M}(\mathcal{C})^H$.


¹ For any YD H -module V , it follows from [7, Proposition 2.2] that there exists a unique Nichols algebra $\mathcal{B}(V)$ of V up to graded Hopf algebra isomorphism in ${}^H_H\mathcal{YD}$. The structure of $\mathcal{B}(V)$ can be described as follows:

Let V be a YD H -module. Then the tensor algebra $T(V) = \bigoplus_{n \geq 0} T(V)(n)$ of the vector space V admits a natural structure of a YD H -module, since ${}^H_H\mathcal{YD}$ is a braided monoidal category. It is then an algebra in ${}^H_H\mathcal{YD}$. Let $T(V) \underline{\otimes} T(V)$ be the tensor product algebra of $T(V)$ with itself in ${}^H_H\mathcal{YD}$. Then there is a unique algebra map $\Delta : T(V) \rightarrow T(V) \underline{\otimes} T(V)$ such that $\Delta(v) = v \otimes 1 + 1 \otimes v$ for all $v \in V$. With this structure, $T(V)$ is a graded braided Hopf algebra in ${}^H_H\mathcal{YD}$. Let $\tilde{I}(V)$ be the sum of all YD H -submodules I of $T(V)$ such that


- I is a graded ideal generated by homogeneous elements of degree ≥ 2 ,
- I is also a coideal, i.e., $\Delta(I) \subset I \otimes T(V) + T(V) \otimes I$.

Then $\mathcal{B}(V) = T(V)/\tilde{I}(V)$.

Here the comodule operation ψ_{V^*} of V^* is defined as follows:




Proof . (i) We easily check that (V^*, α_{V^*}) is a H -module. Now we show that d_V and b_V are H -module homomorphisms. We see that



Thus d_V is an H -module homomorphism.

We also see that



The diagram illustrates the Frobenius property for the counit. On the left, a box labeled S (representing the counit) has an input H and an output V . This is connected to a box labeled α (representing the Frobenius element), which has an input V and an output V^* . On the right, a box labeled ϵ (representing the counit) has an input H and an output V . This is connected to a box labeled V^* (representing the Frobenius element), which has an input V and an output V^* . The two diagrams are shown to be equal, representing the Frobenius property.

Thus b_V is an H -module homomorphism.

(ii) is the dual case of part (i). $\square$

Every algebra (coalgebra, Hopf algebra, quasitriangular Hopf algebra) living in $\mathcal{Vect}(k)$ is called an algebra (coalgebra, Hopf algebra, quasitriangular Hopf algebra) or an ordinary algebra (ordinary coalgebra, ordinary Hopf algebra, ordinary quasitriangular Hopf algebra). We usually use ϕ and δ^- to denote left comodule operations; ψ and δ^+ to denote right comodule operations; α and α^- to denote left module operations; β and α^+ to denote right module operations. We use The Sweedler's and Montgomery's sigma notations (see [114] [87]) for coalgebras and comodules are $\Delta(x) = \sum x_1 \otimes x_2$, $\phi(x) = \sum x^{(-1)} \otimes x^{(0)}$, $\psi(x) = \sum x^{(0)} \otimes x^{(1)}$; or $\Delta(x) = \sum x_{(1)} \otimes x_{(2)}$, $\phi(x) = \sum x_{(-1)} \otimes x_{(0)}$, $\psi(x) = \sum x_{(0)} \otimes x_{(1)}$.

In fact, if H is an ordinary algebra and V is a finite-dimensional left H -module, then V has a left duality H^* in category ${}_H\mathcal{M}$.

Chapter 2

The Maschke's Theorem for Braided Hopf Algebras

In this chapter, we obtain the fundamental theorem of Hopf modules living in braided tensor categories. In Yetter-Drinfeld category, we obtain that the antipode S of H is invertible and the integral $\int_H^l$ of H is a one-dimensional space for any finite-dimensional braided Hopf algebra H . We give Maschke's Theorem for YD Hopf algebras.

Supersymmetry has attracted a great deal of interest from physicists and mathematicians (see [65], [110], [88]). It finds numerous applications in particle physics and statistical mechanics (see [30]). Of central importance in supersymmetric theories is the Z_2 -graded structure which permeates them. Superalgebras and super-Hopf algebras are most naturally very important.

Braided tensor categories were introduced by Joyal and Street in 1986 [61]. It is a generalization of super case. Majid, Joyal, Street and Lyubasheko have obtained many interesting conclusions in braided tensor categories, for example, the braided reconstruction theorem, transmutation and bosonisation, integral, q -Fourier transform, q -Mikowski space, random walk and so on (see [76], [82], [83], [88], [73]).

In many applications one needs to know the integrals of braided Hopf algebras and whether a braided Hopf algebra is semisimple or finite-dimensional. A systematic study of braided Hopf algebras, especially of those braided Hopf algebras that play a role in physics (as, e.g. braided line algebra $\mathbf{C}[x]$ in [76] and anyonic line algebra $\mathbf{C}[\xi] \cong \mathbf{C}\{x\}/\langle x^n \rangle$ in [88]) is very useful.

V. Lyubasheko in [73] introduced the integral for braided Hopf algebras and gave a formula about integral. S. Majid in [88] obtained the integral of anyonic line algebra. In this chapter, we obtain that the antipode S of H is invertible and the integral $\int_H^l$ of H is a one-dimensional space for any finite-dimensional (super) anyonic or q -statistical Hopf algebra H . In particular, if H is a finite-dimensional (super) anyonic Hopf algebra, then $\int_H^l$ is a homogeneous space. We show that a finite-dimensional (super) anyonic or q -statistical Hopf algebra H is semisimple iff $\epsilon(\int_H^l) \neq 0$. Furthermore, if H is a semisimple (super) anyonic Hopf algebra, then H is finite-dimensional.

We begin by recalling some definitions and properties. Attempts to find solutions of the Yang-Baxter equation in a systematic way have led to the theory of quantum groups, in which the concept of (co)quasitriangular Hopf algebra is introduced.

A pair (H, r) is called coquasitriangular Hopf algebra if H is a Hopf algebra over field k and there exists a convolution- invertible k -bilinear form $r: H \otimes H \rightarrow k$ such that the following conditions hold: for any $a, b, c \in H$,

$$(CQT1) \quad r(a, bc) = \sum r(a_1, c)r(a_2, b);$$

$$(CQT2) \quad r(ab, c) = \sum r(a, c_1)r(b, c_2);$$

$$(CTQ3) \quad \sum r(a_1, b_1)a_2b_2 = \sum b_1a_1r(a_2, b_2).$$

Furthermore, if $r^{-1}(a, b) = r(b, a)$ for any $a, b \in H$, then (H, r) is called cotriangular.

For coquasitriangular Hopf algebra (H, r) , we can define a braiding C^r in the category ${}^H\mathcal{M}$ of H -comodules as follows:

$$C_{U,V}^r : U \otimes V \longrightarrow V \otimes U$$

sending $u \otimes v$ to $\sum r(v^{(-1)}, u^{(-1)}) \otimes v^{(0)}u^{(0)}$ for any $u \in U, v \in V$, where (U, ϕ_U) and (V, ϕ_V) are H -comodule and $\phi_V(v) = \sum v^{(-1)} \otimes v^{(0)}$ (see [64, Proposition VIII.5.2]). This braided tensor category is called one determined by coquasitriangular structure. Dually, we have the concept of quasitriangular Hopf algebra. We also have that $({}^H\mathcal{M}, C^R)$ is a braided tensor category, which is called one determined by quasitriangular structure.

For example, let $H = \mathbf{C}\mathbf{Z}_n$ and $r(a, b) = (e^{\frac{2\pi i}{n}})^{ab}$ for any $a, b \in \mathbf{Z}_n$, where $\mathbf{C}$ is the complex field. It is clear that $(\mathbf{C}\mathbf{Z}_n, r)$ is a coquasitriangular Hopf algebra. Thus $(\mathbf{C}\mathbf{Z}_n\mathcal{M}, C^r)$ is a braided tensor category, usually written as $\mathcal{C}_n$. Every algebra or Hopf algebra living in $\mathcal{C}_n$ is called an anyonic algebra or anyonic Hopf algebra (see [83, Example 9.2.4]). Every algebra or Hopf algebra living in $\mathcal{C}_2$ is called a superalgebra or super-Hopf algebra.

If $H = \mathbf{C}\mathbf{Z}$ and $r(a, b) = q^{ab}$ for any $a, b \in \mathbf{Z}$, then $(\mathbf{C}\mathbf{Z}\mathcal{M}, C^r)$ is a braided tensor category, where $0 \neq q \in \mathbf{C}$. Every algebra or Hopf algebra living in it is called a q-statistical algebra or q-statistical Hopf algebra (see [83, Definition 10.1.9]).

For example, let (H, r) be a cocommutative cotriangular Hopf algebra. If (L, ϕ) is a left H -comodule with an H -comodule homomorphism $[\ , \]$ from $L \otimes L$ to L in $({}^H\mathcal{M}, C^r)$ such that the following conditions hold:

$$(i) \quad [\ , \](id + C_{L,L}^r) = 0;$$

$$(ii) \quad [\ , \]([\ , \] \otimes id)(id + (id_L \otimes C_{L,L}^r)(C_{L,L}^r \otimes id_L) + (C_{L,L}^r \otimes id_L)(id_L \otimes C_{L,L}^r)) = 0,$$

then $(L, [\ , \])$ is called a Lie H -algebra or a Lie algebra living in $({}^H\mathcal{M}, C_{L,L}^r)$ (see [14, Definition 1.5]). If $H = \mathbf{C}\mathbf{Z}_2$ and $r(a, b) = (-1)^{ab}$ for any $a, b \in \mathbf{Z}_2$, then every Lie H -algebra is called a Lie superalgebra (see [65]). If $H = kG$ and r is a bicharacter of abelian group G with $r^{-1}(a, b) = r(b, a)$ for any $a, b \in G$, then every Lie H -algebra is called a Lie colour algebra (see [87]) or ϵ Lie algebra (see [110, Definition 2.1]). If $(L, [\ , \])$ is a Lie colour algebra, then the universal enveloping algebra $U(L)$ of L is an algebra living in $({}^H\mathcal{M}, C^r)$ (see [110]). Thus the universal enveloping algebras of Lie superalgebras are superalgebras.

2.1 The integrals in braided tensor categories

In this section, we obtain the fundamental theorem of Hopf modules living in braided tensor categories.

An object E in category $\mathcal{C}$ is called a final object if for every object V in $\mathcal{C}$, there exists a unique morphism f from V to E . For two morphisms $f_1, f_2 : A \rightarrow B$ in $\mathcal{C}$, set $target(A, f_1, f_2) =: \{(D, g) \mid g \text{ is a morphism from } D \text{ to } A \text{ such that } f_1 g = f_2 g\}$ and $source(B, f_1, f_2) =: \{(D, g) \mid g \text{ is a morphism from } B \text{ to } D \text{ such that } g f_1 = g f_2\}$, the final object of full subcategory $target(A, f_1, f_2)$ and the initial object of full subcategory $source(B, f_1, f_2)$ are called equaliser and coequaliser of f_1 and f_2 , written as $equaliser(f_1, f_2)$ and $coequaliser(f_1, f_2)$, respectively. If (M, ϕ) is a left comodule over bialgebra H in $\mathcal{C}$, then $equaliser(\phi, \eta_H \otimes id_M)$ is called the coinvariants of M , written (M^{coH}, i) . That is, for every H -comodule N in $\mathcal{M}$ and a morphism f from N to M in ${}^H\mathcal{M}(\mathcal{C})$, then there exists a unique a morphism $\bar{f}$ from N to M^{coH} in ${}^H\mathcal{M}(\mathcal{C})$ such that $f = i \bar{f}$.

Definition 2.1.1. Let H be a bialgebra, (M, α) be a left H -module and (M, ϕ) a left H -comodule. If α is an H -comodule morphism and ϕ is an H -module morphism, then (M, α, ϕ) is called an H -Hopf module.

We easily know that (M, α, ϕ) is an H -Hopf module iff

Theorem 2.1.2. (The fundamental theorem of Hopf modules) Let $(\mathcal{C}, C)$ be a braided tensor category. If (M, α, ϕ) is an H -Hopf module in braided tensor category $(\mathcal{C}, C)$ and there exists the equaliser $(\phi, \eta_H \otimes id_M)$, then

$$H \otimes M^{coH} \cong M \quad (\text{as } H\text{-Hopf modules.})$$

Proof. Let $u =$. We have that $\phi u =$ $=$ by Pro.1.0.13(i)

$=$. Thus there exists an H -comodule homomorphism $\bar{u}$ from M to

M^{coH} such that $u = i\bar{u}$.

Let $\xi = \begin{array}{c} HM^{coH} \\ | \\ \alpha \\ | \\ M \end{array}$ and $\psi = \begin{array}{c} M \\ | \\ \phi \\ | \\ H \quad M \\ | \quad | \\ \bar{u} \end{array}$. We see that $\xi\psi = \begin{array}{c} M \\ | \\ \phi \\ | \\ \alpha \\ | \\ M \end{array} = \begin{array}{c} M \\ | \\ \phi \\ | \\ S \\ | \\ \alpha \\ | \\ M \end{array} = \begin{array}{c} M \\ | \\ \phi \\ | \\ S \\ | \\ \alpha \\ | \\ M \end{array} = \begin{array}{c} M \\ | \\ \phi \\ | \\ S \\ | \\ \alpha \\ | \\ M \end{array}$ and

$$\psi\xi = \begin{array}{c} HM^{coH} \\ | \\ \alpha \\ | \\ \phi \\ | \\ S \\ | \\ \alpha \\ | \\ M \quad M^{coH} \end{array} = \begin{array}{c} H \quad M \\ | \quad | \\ \phi \\ | \\ \alpha \\ | \\ S \\ | \\ \alpha \\ | \\ M^{coH} \end{array} = \begin{array}{c} H \quad M^{coH} \\ | \quad | \\ \phi \\ | \\ \alpha \\ | \\ S \\ | \\ \alpha \\ | \\ M^{coH} \end{array} = \begin{array}{c} H \quad M^{coH} \\ | \quad | \\ S \\ | \\ \alpha \\ | \\ M^{coH} \end{array} = \begin{array}{c} HM^{coH} \\ | \\ HM^{coH} \end{array}.$$

Thus ξ is an isomorphism.

Since $\begin{array}{c} H \otimes M^{coH} \\ | \\ \xi \\ | \\ \phi \\ | \\ H \quad M^{coH} \end{array} = \begin{array}{c} H \quad M^{coH} \\ | \quad | \\ \phi \\ | \\ \alpha \\ | \\ H \quad M^{coH} \end{array} = \begin{array}{c} H \otimes M^{coH} \\ | \\ \phi \\ | \\ H \quad M^{coH} \end{array}$ and

$$\begin{array}{c} H \\ | \\ S \\ | \\ \alpha \\ | \\ M \end{array} = \begin{array}{c} H \\ | \\ S \\ | \\ \alpha \\ | \\ M \end{array} = \begin{array}{c} H \quad H \quad M^{coH} \\ | \quad | \quad | \\ \alpha \\ | \\ M \end{array} = \begin{array}{c} H \quad H \otimes M^{coH} \\ | \quad | \\ \alpha \\ | \\ M \end{array}.$$

Consequently, ξ is an H -Hopf module isomorphism. $\square$

Lemma 2.1.3. *If H is a Hopf algebra with left duality, then (H^*, α, ϕ) is a left H -Hopf*

module, where the operations of module and comodule are defined as follows:

$$\begin{array}{c} H \quad H^* \\ \downarrow \quad \downarrow \\ \alpha \\ \downarrow \\ H^* \end{array} = \begin{array}{c} H \quad H^* \\ \downarrow \quad \downarrow \\ \neg \\ \downarrow \\ H^* \end{array} = \begin{array}{c} H \quad H^* \\ \downarrow \quad \downarrow \\ S \\ \downarrow \quad \downarrow \\ H^* \end{array}, \quad \begin{array}{c} H^* \\ \downarrow \\ \phi \\ \downarrow \\ H \quad H^* \end{array} = \begin{array}{c} H^* \\ \downarrow \\ H \quad H^* \end{array}.$$

Proof. We see that

$$\begin{array}{c} H \quad H \quad H^* \\ \downarrow \quad \downarrow \quad \downarrow \\ \alpha \\ \downarrow \\ H^* \end{array} = \begin{array}{c} H \quad H \quad H^* \\ \downarrow \quad \downarrow \quad \downarrow \\ S \\ \downarrow \quad \downarrow \quad \downarrow \\ H^* \end{array} \stackrel{\text{by Pro.1.0.13(i)}}{=} \begin{array}{c} H \quad H \quad H^* \\ \downarrow \quad \downarrow \quad \downarrow \\ S \quad S \\ \downarrow \quad \downarrow \quad \downarrow \\ H^* \end{array} \\
 \stackrel{\text{by Pro.1.0.13(ii)}}{=} \begin{array}{c} H \quad H \quad H^* \\ \downarrow \quad \downarrow \quad \downarrow \\ S \quad S \\ \downarrow \quad \downarrow \quad \downarrow \\ H^* \end{array} = \begin{array}{c} H \quad H \quad H^* \\ \downarrow \quad \downarrow \quad \downarrow \\ S \quad S \\ \downarrow \quad \downarrow \quad \downarrow \\ H^* \end{array} = \begin{array}{c} H \quad H \quad H^* \\ \downarrow \quad \downarrow \quad \downarrow \\ \alpha \\ \downarrow \\ H^* \end{array} \\
 \begin{array}{c} \eta_H \\ \downarrow \\ \alpha \\ \downarrow \\ H^* \end{array} = \begin{array}{c} H^* \\ \downarrow \\ \eta_H \\ \downarrow \\ H^* \end{array} = \begin{array}{c} H^* \\ \downarrow \\ H^* \end{array}.$$

Thus (H^*, α) is an H -module.

We see that

by Pro 1.0.13(i)

and

The diagram shows a sequence of equalities between string diagrams. The first diagram on the left has two strands labeled H and H^* entering a node α , and two strands labeled H and H^* exiting a node ϕ . This is equal to a diagram where the H strands are connected by a cup and the H^* strands by a cap, with a node S on the H strand. This is further equal to a diagram with a crossing and a node S . The final diagram shows the strands rearranged with a node S and a combined strand labeled HH^* .

Thus

The diagram shows an equality between two string diagrams. On the left, there are two strands labeled H and H^* . The top strand has a node α and the bottom strand has a node ϕ . On the right, the strands are crossed, and the nodes α and ϕ are now on the opposite strands.

Consequently, (H^*, α, ϕ) is an H -Hopf module. $\square$

2.2 The Integrals of Braided Hopf algebras in Yetter-Drinfeld Categories

In this section, using the fundamental theorem of Hopf modules in braided tensor categories, we give the relation between the integrals and semisimplicity of Braided Hopf algebras in Yetter-Drinfeld Categories.

Assume that H is a YD Hopf algebra and (M, ϕ) is a H -comodule in ${}^B_B\mathcal{YD}$. Obviously,

$$\{x \in M \mid \phi(x) = \eta_H \otimes x\} = \ker(\phi - \eta_H \otimes id_M).$$

It is a straightforward verification that (M^{coH}, i) is the *equivaiser* $(\phi, \eta_H \otimes id_M)$ with

$$M^{coH} := \{x \in M \mid \phi(x) = \eta_H \otimes x\}$$

and inclusion map i . Since $\phi - \eta_H \otimes id_M$ is a morphism in ${}^B_B\mathcal{YD}$, $\ker(\phi - \eta_H \otimes id_M)$ is object in ${}^B_B\mathcal{YD}$. Let $\int_H^l := \{x \in H \mid hx = \epsilon(h)x, \text{ for any } h \in H\}$, called the integral space of H .

If H is a finite-dimensional Hopf algebra living in a Yetter-Drinfeld category, then H has a left duality H^* (see [64, Example 1 P347] or Lemma 1.0.17) and $\int_{H^*}^l = (H^*)^{coH}$, where the H -comodule operation of H^* is defined in Lemma 2.1.3.

Theorem 2.2.1. *(The Maschke's theorem) Assume B is an ordinary Hopf algebra over field k with invertible antipode. If H is a finite-dimensional Hopf algebra living in Yetter-Drinfeld category ${}^B_B\mathcal{YD}$, then*

- (1) $H \otimes \int_{H^*}^l \cong H^*$ (as H -Hopf modules in ${}^B_B\mathcal{YD}$);
- (2) $\int_H^l$ is a one dimensional space;
- (3) the antipode S of H is bijective;
- (4) H is semisimple iff $\epsilon(\int_H^l) \neq 0$;
- (5) Every semisimple Hopf algebra H living in ${}^B_B\mathcal{YD}$ is finite-dimensional.

Proof. (1) It follows from the above discussion.

(2) It follows from part (1).

(3) According to the proof of Lemma 2.1.2 and Lemma 2.1.3, we have that

$$\xi = (id_{H^*} \otimes d_H)(id_{H^*} \otimes C_{H,H^*})(C_{H,H^*} \otimes id_{H^*})(id_H \otimes \Delta)(S \otimes id_{\int_{H^*}^l})$$

is an isomorphism from $H \otimes \int_{H^*}^l$ to H^* , which implies that S is injective. Considering H is a finite-dimensional vector space, we obtain that S is bijective.

(4) If H is semisimple then there is a left ideal I such that

$$H = I \oplus \ker \epsilon.$$

For any $y \in I, h \in H$, we see that

$$\begin{aligned} hy &= ((h - \epsilon(h)1_H) + \epsilon(h)1_H)y \\ &= (h - \epsilon(h)1_H)y + \epsilon(h)y \\ &= \epsilon(h)y \quad \text{since } (h - \epsilon(h)1_H)y \in (\ker \epsilon)I = 0. \end{aligned}$$


Thus $y \in \int_H^l$, and so $I \subseteq \int_H^l$, which implies $\epsilon(\int_H^l) \neq 0$.

Conversely, if $\epsilon(\int_H^l) \neq 0$, let $z \in \int_H^l$ with $\epsilon(z) = 1$. Say M is a left H -module and N is an H -submodule of M . Assume that ξ is a k -linear projection from M to N . We define

$$\mu(m) = \sum z_1 \cdot \xi(S(z_2) \cdot m)$$

for any $m \in M$. It is sufficient to show that μ is an H -module projection from M to N . Obviously,

μ is a k -linear projection. Now we only need to show that it is an H -module map.



Thus, μ is an H -module morphism. $\square$

(5) The smash product $H \# B$ is an ordinary semisimple Hopf algebra by bosonisation [79, Theorem 4.1] and by [87, Theorem 7.4.2]. Thus $H \# B$ is finite-dimensional, which implies that H is finite dimensional. $\square$

Lemma 2.2.2. *Let B be an ordinary finite-dimensional Hopf algebra with left duality B^* . If (H, ϕ) is a left B -comodule algebra and B -comodule coalgebra, then (H, α) is a left $B^{* \text{ cop}}$ -module*

algebra and B^* -module coalgebra, where

$$\begin{array}{c} B^* \\ | \\ \text{---} \alpha \text{---} \\ | \\ H \end{array} = \begin{array}{c} B^* \\ | \\ \text{---} \phi \text{---} \\ | \\ H \end{array}.$$

Proof. It is trivial. $\square$

Proposition 2.2.3. *Let $\text{char } k = 0$.*

(i) *If H is not only a kG -module algebra but also a kG -module coalgebra, then the action of G on $\int_H^l$ is stable;*

(ii) *If H is a Hopf algebra living in $(\mathcal{M}^{kG}, C^r)$ and G is a commutative finite group with a bicharacter r , then $\int_H^l$ is a homogeneous space of G -graded space H .*

In particular, $\int_H^l$ is an (super) anyonic space for every finite-dimensional (super) anyonic Hopf algebra H .

Proof. (i) For any $g \in G, x \in H, t \in \int_H^l$, we see that

$$\begin{aligned} x(g \cdot t) &= (g \cdot y)(g \cdot t) \quad \text{since } H \text{ is a } kG\text{-module algebra, where } y = g^{-1} \cdot x \\ &= g \cdot (yt) \\ &= \epsilon(y)(g \cdot t) \\ &= \epsilon(x)(g \cdot t) \quad \text{since } H \text{ is a } kG\text{-module coalgebra.} \end{aligned}$$

Therefore action of G on $\int_H^l$ is stable.

(ii) Since H is a left kG -comodule algebra and kG -comodule coalgebra, we have that H is a left $(kG)^*$ -module algebra and $(kG)^*$ -module coalgebra by Lemma 2.2.2. However, $(kG)^* = k\hat{G}$ (as Hopf algebra) by [83, Corollary 1.5.6], where $\hat{G}$ denotes the character group of G . Thus H is a left $k\hat{G}$ -module algebra and a $k\hat{G}$ -module coalgebra, which implies that the action of $\hat{G}$ on $\int_H^l$ is stable by part (1). That is, $\int_H^l$ is a sub-module of $(kG)^*$ -module H , which implies that $\int_H^l$ is a homogeneous space of G -graded space H . $\square$

Example 2.2.4. *Let H denote the anyonic line algebra, i.e. $H = \mathbf{C}\{x\} / \langle x^n \rangle$, where $\mathbf{C}\{x\}$ is a free algebra over complex field $\mathbf{C}$ and $\langle x^n \rangle$ is an ideal generated by x^n of $\mathbf{C}\{x\}$. Set $\mathbf{C}\{x\} / \langle x^n \rangle = \mathbf{C}[\xi]$ with $\xi^n = 0$. Its comultiplication and counit and antipode are*

$$\Delta(\xi) = \xi \otimes 1 + 1 \otimes \xi \quad \text{and} \quad \epsilon(\xi) = 0 \quad \text{and} \quad S(\xi) = -\xi$$

respectively. It is straightforward to check that H is an anyonic Hopf algebra (see [88]). Obviously, the dimension of H is n . It is clear that $\int_H^l = \mathbf{C}\xi^{n-1}$. Since $\epsilon(\int_H^l) = 0$, H is not semisimple.

Example 2.2.5. (see [83, P510]) *Let $H = \mathbf{C}[x]$ denote the braided line algebra. It is just the usual algebra $\mathbf{C}[x]$ of polynomials in x over complex field $\mathbf{C}$, but we regard it as a q -statistical Hopf algebra with*

$$\Delta(x) = x \otimes 1 + 1 \otimes x, \quad \epsilon(x) = 0, \quad S(x) = -x, \quad |x^n| = n$$

and

$$C^r(x^n, x^m) = q^{nm}(x^m \otimes x^n).$$

If $y = \sum_0^n a_i x^i \in \int_H^l$, then $a_i = 0$ for $i = 1, 2, \dots, n$ since $xy = \epsilon(x)y = 0$. Thus $\int_H^l = 0$. It follows from the proof of Theorem 2.2.1 (5) that H is not semisimple.

By the way, we give some properties of group graded (co)algebras.

An algebra A is called a G -graded algebra if $A = \sum_{g \in G} \oplus A_g$, $A_g A_h \subseteq A_{gh}$ and $1_A \in A_e$, where e is the unity element of G . A coalgebra D is called a G -graded coalgebra if $D = \sum_{g \in G} \oplus D_g$, $\Delta(D_g) \subseteq \sum_{x, y \in G, xy=g} D_x \otimes D_y$ and $\epsilon(D_h) = 0$ for any $h, g \in G$ and $h \neq e$.

Proposition 2.2.6. (1) A is a right kG -comodule algebra iff A is a G -graded algebra;
(2) D is a right kG -comodule coalgebra iff D is a G -graded coalgebra.

Proof. (1) It was obtained in [29, Proposition 1.3].

(2) If (D, ψ) is a kG -comodule coalgebra, then $D = \sum_{g \in G} \oplus D_g$ is a G -graded space by [20, Lemma 4.8]. For any homogeneous $0 \neq d \in D_g$, let

$$\Delta(d) = \sum_{i=1,2,\dots,m} (d_{x_i} \otimes d_{y_i})$$

where $0 \neq d_{x_i} \in D_{x_i}$, $0 \neq d_{y_i} \in D_{y_i}$, $x_i, y_i \in G$ for $i = 1, 2, \dots, m$. Since Δ is a kG -comodule homomorphism we have that

$$\sum_{i=1,2,\dots,m} (d_{x_i} \otimes d_{y_i} \otimes g) = \sum_{i=1,2,\dots,m} (d_{x_i} \otimes d_{y_i} \otimes x_i y_i).$$

Thus

$$x_i y_i = g$$

for $i = 1, 2, \dots, m$, which implies that

$$\Delta(D_g) \subseteq \sum_{x, y \in G, xy=g} D_x \otimes D_y.$$

Since ϵ is a kG -comodule homomorphism, we have that

$$\epsilon(d)g = \epsilon(d)e$$

for any $g \in G, d \in D_g$, which implies that $\epsilon(D_g) = 0$ for any $e \neq g \in G$. Consequently, D is a G -graded coalgebra.

Conversely, if $D = \sum_{g \in G} \oplus D_g$ is a G -graded coalgebra, then (D, ψ) is a right kG -comodule by [20, Lemma 4.8]. It is sufficient to show that Δ and ϵ are kG -comodule maps. For any $d \in D_g$, we assume that

$$\Delta(d) = \sum_{x, y \in G, xy=g} d_x \otimes d_y$$

where $d_x \in D_x, d_y \in D_y$. We see that

$$\begin{aligned}
(\Delta \otimes id_D)\psi(d) &= (\Delta \otimes id_D)(d \otimes g) \\
&= \sum_{x,y \in G, xy=g} d_x \otimes d_y \otimes g \\
&= (id_{D \otimes D} \otimes m_{kG})(id_D \otimes \tau \otimes id_{kG})(\psi \otimes \psi)\Delta(d)
\end{aligned}$$

where τ is an ordinary twist map.

Thus Δ is a kG -comodule map. Since

$$(\epsilon \otimes id_{kG})\psi(d) = \epsilon(d)g = \epsilon(d)e = \epsilon(d)1_{kG}$$

for any $d \in D_g$, we have that ϵ is a kG -comodule map. $\square$

Chapter 3

The Double Bicrossproducts in Braided Tensor Categories

In this chapter, we construct the double bicrossproduct $D = A_\alpha^\phi \bowtie_\beta^\psi H$ of two bialgebras A and H in a braided tensor category and give the necessary and sufficient conditions for D to be a bialgebra. We study the universal property of double bicrossproduct.

Throughout this chapter, we assume that H and A are two bialgebras in braided tensor categories and

$$\begin{aligned} \alpha : H \otimes A &\rightarrow A, & \beta : H \otimes A &\rightarrow H, \\ \phi : A &\rightarrow H \otimes A, & \psi : H &\rightarrow H \otimes A \end{aligned}$$

such that (A, α) is a left H -module coalgebra, (H, β) is a right A -module coalgebra, (A, ϕ) is a left H -comodule algebra, and (H, ψ) is a right A -comodule algebra.

We define the multiplication m_D , unit η_D , comultiplication Δ_D and counit ϵ_D in $A \otimes H$ as follows:

and $\epsilon_D = \epsilon_A \otimes \epsilon_H$, $\eta_D = \eta_A \otimes \eta_H$. We denote $(A \otimes H, m_D, \eta_D, \Delta_D, \epsilon_D)$ by

$$A_\alpha^\phi \bowtie_\beta^\psi H,$$

which is called the double bicrossproduct of A and H . We denote it by $A \overset{b}{\bowtie} H$ in short.

If one of $\alpha, \beta, \phi, \psi$ is trivial, we omit the trivial one in the notation $A_\alpha^\phi \bowtie_\beta^\psi H$. For example, when ϕ and ψ are trivial, we denote $A_\alpha^\phi \bowtie_\beta^\psi H$ by $A_\alpha \bowtie_\beta H$. When α and β are trivial, we denote $A_\alpha^\phi \bowtie_\beta^\psi H$ by $A^\phi \bowtie^\psi H$. We call $A_\alpha \bowtie_\beta H$ a double cross product and denote it by $A \bowtie H$ in

short. We call $A^\phi \bowtie^\psi H$ a double cross coproduct and denote it by $A \overset{c}{\bowtie} H$ in short. We call $A_\alpha^\phi \bowtie H$ ($A \bowtie_\beta^\psi H$) and $A_\alpha \bowtie^\psi H$ ($A^\phi \bowtie_\beta H$) a biproduct and a bicrossproduct respectively. Note that S. Majid uses different notations.

We give some notation as follows.

$$\begin{aligned}
(M1): \quad & \begin{array}{c} H \\ | \\ \text{---} \circlearrowleft \eta_A \\ | \\ \alpha \\ | \\ A \end{array} = \begin{array}{c} H \\ | \\ \epsilon_H \\ | \\ A \end{array}, \quad \begin{array}{c} A \\ | \\ \text{---} \circlearrowleft \eta_H \\ | \\ \beta \\ | \\ H \end{array} = \begin{array}{c} A \\ | \\ \epsilon_A \\ | \\ H \end{array}; \\
(M2): \quad & \begin{array}{c} H \quad A \quad A \\ | \quad | \quad | \\ \text{---} \circlearrowleft \alpha \\ | \\ A \end{array} = \begin{array}{c} H \quad A \quad A \\ | \quad | \quad | \\ \text{---} \circlearrowleft \alpha \quad \text{---} \circlearrowleft \beta \\ | \quad | \quad | \\ \text{---} \circlearrowleft \alpha \\ | \\ A \end{array}; \\
(M3): \quad & \begin{array}{c} H \quad H \quad A \\ | \quad | \quad | \\ \text{---} \circlearrowleft \beta \\ | \\ H \end{array} = \begin{array}{c} H \quad H \quad A \\ | \quad | \quad | \\ \text{---} \circlearrowleft \alpha \quad \text{---} \circlearrowleft \beta \\ | \quad | \quad | \\ \text{---} \circlearrowleft \beta \\ | \\ H \end{array}; \\
(M4): \quad & \begin{array}{c} H \quad A \\ | \quad | \\ \text{---} \circlearrowleft \beta \quad \text{---} \circlearrowleft \alpha \\ | \quad | \\ H \quad A \end{array} = \begin{array}{c} H \quad A \\ | \quad | \\ \text{---} \circlearrowleft \alpha \quad \text{---} \circlearrowleft \beta \\ | \quad | \\ H \quad A \end{array}; \\
(CM1): \quad & \begin{array}{c} A \\ | \\ \text{---} \circlearrowleft \phi \\ | \\ H \end{array} = \begin{array}{c} A \\ | \\ \epsilon_A \\ | \\ H \end{array}, \quad \begin{array}{c} H \\ | \\ \text{---} \circlearrowleft \psi \\ | \\ A \end{array} = \begin{array}{c} H \\ | \\ \epsilon_H \\ | \\ A \end{array};
\end{aligned}$$

(CM2) :

Diagrammatic equation (CM2) showing a node ϕ with input H and output A , and another input A , equal to a more complex structure involving nodes ϕ and ψ with inputs H , A , and A .

(CM3) :

Diagrammatic equation (CM3) showing a node ψ with input H and output H , and another input HA , equal to a more complex structure involving nodes ψ and ϕ with inputs H , H , and A .

(CM4) :

Diagrammatic equation (CM4) showing two nodes ψ and ϕ with inputs H and A , and outputs H and A , equal to a similar structure with nodes ϕ and ψ and inputs H and A .

(B1) :

Diagrammatic equation (B1) showing a node α with input A and output A , and another input A , equal to a more complex structure involving nodes ϕ and α with inputs A and A .

(B2) :

Diagrammatic equation (B2) showing a node α with input H and output A , and another input A , equal to a more complex structure involving nodes ψ and α with inputs H , H , A , and A .

$$(B3): \quad \begin{array}{c} H \quad H \\ \text{---} \text{---} \\ | \\ \text{---} \text{---} \\ H \quad H \end{array} = \begin{array}{c} H \quad H \\ | \quad | \\ \text{---} \text{---} \\ | \quad | \\ \text{---} \text{---} \\ H \quad H \end{array} \quad ;$$

$$(B4): \quad \begin{array}{c} H \quad A \\ \text{---} \text{---} \\ | \\ \text{---} \text{---} \\ H \quad H \end{array} = \begin{array}{c} H \quad A \\ | \quad | \\ \text{---} \text{---} \\ | \quad | \\ \text{---} \text{---} \\ H \quad H \end{array} \quad ;$$

$$(B5): \quad \begin{array}{c} A \quad H \quad A \quad H \\ | \quad | \quad | \quad | \\ \text{---} \text{---} \text{---} \text{---} \\ | \quad | \quad | \quad | \\ \text{---} \text{---} \text{---} \text{---} \\ H \quad A \quad H \quad A \end{array} = \begin{array}{c} A \quad H \quad A \quad H \\ | \quad | \quad | \quad | \\ \text{---} \text{---} \text{---} \text{---} \\ | \quad | \quad | \quad | \\ \text{---} \text{---} \text{---} \text{---} \\ H \quad A \quad H \quad A \end{array} \quad ;$$

$$(B6): \quad \begin{array}{c} H \quad A \\ | \quad | \\ \text{---} \text{---} \\ | \quad | \\ \text{---} \text{---} \\ H \quad A \end{array} = \begin{array}{c} H \quad A \\ | \quad | \\ \text{---} \text{---} \\ | \quad | \\ \text{---} \text{---} \\ H \quad A \end{array} \quad ;$$

$$(B6') : \quad \begin{array}{c} H \quad A \\ | \quad | \\ \diagdown \quad \diagup \\ | \quad | \\ \diagup \quad \diagdown \\ | \quad | \\ H \quad A \end{array} \quad \begin{array}{c} \beta \\ \psi \end{array} = \begin{array}{c} H \quad A \\ | \quad | \\ \psi \quad \beta \\ | \quad | \\ H \quad A \end{array} ;$$

$$(CB2) : \quad \begin{array}{c} A \quad A \\ | \quad | \\ \phi \\ | \quad | \\ H \quad A \end{array} = \begin{array}{c} A \quad A \\ | \quad | \\ \phi \quad \phi \\ | \quad | \\ \beta \quad \phi \\ | \quad | \\ H \quad A \end{array} ;$$

$$(CB4) : \quad \begin{array}{c} H \quad H \\ | \quad | \\ \psi \\ | \quad | \\ H \quad A \end{array} = \begin{array}{c} H \quad H \\ | \quad | \\ \psi \quad \psi \\ | \quad | \\ \psi \quad \alpha \\ | \quad | \\ H \quad A \end{array} ;$$

$$(B7) : \quad \begin{array}{c} H \quad A \\ | \quad | \\ \alpha \quad \psi \\ | \quad | \\ H \quad A \end{array} = \begin{array}{c} H \quad A \\ | \quad | \\ \psi \quad \alpha \\ | \quad | \\ H \quad A \end{array} ;$$

$$(B7') : \quad \begin{array}{c} H \quad A \\ | \quad | \\ \phi \quad \beta \\ | \quad | \\ H \quad A \end{array} = \begin{array}{c} H \quad A \\ | \quad | \\ \beta \quad \phi \\ | \quad | \\ H \quad A \end{array} .$$

Let

$$\begin{array}{c} H \\ \downarrow \\ \text{---} \circlearrowleft i_H \text{---} \\ \uparrow \\ A \end{array} = \begin{array}{c} H \\ \downarrow \\ \text{---} \circlearrowleft \eta_A \text{---} \\ \uparrow \\ A \end{array} ; \quad \begin{array}{c} A \\ \downarrow \\ \text{---} \circlearrowleft i_A \text{---} \\ \uparrow \\ H \end{array} = \begin{array}{c} A \\ \downarrow \\ \text{---} \circlearrowleft \eta_H \text{---} \\ \uparrow \\ H \end{array} ; \quad \begin{array}{c} A \quad H \\ \downarrow \\ \text{---} \circlearrowleft \pi_H \text{---} \\ \uparrow \\ H \end{array} = \begin{array}{c} A \quad H \\ \downarrow \\ \text{---} \circlearrowleft \epsilon_A \text{---} \\ \uparrow \\ H \end{array} ; \quad \begin{array}{c} A \quad H \\ \downarrow \\ \text{---} \circlearrowleft \pi_A \text{---} \\ \uparrow \\ A \end{array} = \begin{array}{c} A \quad H \\ \downarrow \\ \text{---} \circlearrowleft \epsilon_A \text{---} \\ \uparrow \\ A \end{array}.$$

π_A and π_H are called trivial actions; i_A and i_H are called trivial coactions.

3.1 Double bicrossproduct

In this section, we give the sufficient and necessary conditions for double bicrossproduct $A_\alpha^\phi \bowtie_\beta^\psi H$ to be a bialgebra.

Lemma 3.1.1. (i) If A and H are bialgebras and H acts on A trivially, then A is an H -module algebra and an H -module coalgebra.

(ii) If A and H are bialgebras and H coacts on A trivially, then A is an H -comodule algebra and an H -comodule coalgebra.

(iii) $C_{I,I} = id_I$, $C_{I,I \otimes I} = a_{I,I,I}^{-1}$, $C_{I \otimes I, I} = a_{I,I,I}$;

(iv) If we define:


$$\Delta_I = r_I^{-1}, \quad m_I = r_I, \quad \epsilon_I = id_I = \eta_I = S,$$

then I is a Hopf algebra.


(v) If H is a bialgebra, then I is an H -module algebra, H -module coalgebra, H -comodule algebra, H -comodule coalgebra. Meantime, H is an I -module algebra, I -module coalgebra, I -comodule algebra, I -comodule coalgebra.


(vi) If B is a coalgebra and A is an algebra, then $Hom_{\mathcal{C}}(B, A)$ is a monoid, i.e. $Hom_{\mathcal{C}}(B, A)$ is a semigroup with unit.

(vii) If (B1) and (B2) hold, then




(viii) If (B3) and (B4) hold, then






Similarly, we can check that the right hand is equal to the diagram above. Therefore, the equation holds.

Next we see that




Thus D is an algebra.

Conversely, if D is an algebra, we show that $(M1)$, $(M2)$ and $(M3)$ hold. By



We have that



Similarly, we have that

$$\begin{array}{c} A \\ | \\ \eta_H \\ | \\ \beta \\ | \\ H \end{array} = \begin{array}{c} A \\ | \\ \epsilon_A \\ | \\ \eta_H \\ | \\ H \end{array}.$$

Thus (M1) holds.

By associative law, we have that

Thus (2) holds. Similarly, we can show that (M3) holds. $\square$

Lemma 3.1.3. $D = A^\phi \bowtie^\psi H$ is a coalgebra iff (CM1)–(CM3) hold.



Proof. It follows by turning the diagram in proof of Theorem 3.1.2 upside down. $\square$

Theorem 3.1.4. Assume that H and A are two bialgebras in braided tensor category $\mathcal{C}$ and (A, α) is a left H -module coalgebra, (H, β) is a right A -module coalgebra, (A, ϕ) is a left H -comodule algebra, and (H, ψ) is a right A -comodule algebra in $\mathcal{C}$. Then $D = A^\phi_\alpha \bowtie^\psi_\beta H$ is a bialgebra in $\mathcal{C}$ iff (M1)–(M3), (CM1)–(CM3), (B1)–(B5) hold.


Proof. It is sufficient to show that (B1)–(B5) hold iff


$$\begin{array}{c} D \quad D \\ | \quad | \\ \text{---} \\ | \quad | \\ D \quad D \end{array} = \begin{array}{c} D \quad D \\ | \quad | \\ \text{---} \\ | \quad | \\ D \quad D \end{array} \cdot \dots\dots (1)$$


If (B1) – (B5) hold, see that





by Lemma 3.1.1(vii)(viii)















Conversely, if (1) holds, we have that




and obtain (B1) by computation.


Similarly, we can obtain (B2) by applying $\eta_A \otimes id \otimes id \otimes \eta_H$ to the top of relation (1) and $id \otimes \epsilon_H \otimes id \otimes \epsilon_H$ to the bottom of relation (1). We can also obtain (B3) and (B4) by relation (1), $\eta_A \otimes id \otimes \eta_A \otimes id$, $\epsilon_A \otimes id \otimes \epsilon_A \otimes id$, $\eta_A \otimes id \otimes id \otimes \eta_H$ and $\epsilon_A \otimes id \otimes \epsilon_A \otimes id$ respectively. By applying $\epsilon_A \otimes id \otimes id \otimes \epsilon_A$ to relation (1), we obtain (B5). $\square$


Theorem 3.1.5. *Let $D = A_\alpha^\phi \bowtie_\beta^\psi H$ be a bialgebra. If A and H are Hopf algebras with antipodes S_A and S_H respectively, then D is a Hopf algebra with an antipode*

$$S_D = \begin{array}{c} \begin{array}{cc} A & H \\ \downarrow & \downarrow \\ \phi & \phi \\ \downarrow & \downarrow \\ \psi & \psi \\ \downarrow & \downarrow \\ \alpha & \beta \\ \downarrow & \downarrow \\ A & H \end{array} \\ S_D = \begin{array}{cc} \downarrow & \downarrow \\ S_H & S_A \\ \downarrow & \downarrow \\ \alpha & \beta \\ \downarrow & \downarrow \\ A & H \end{array} \end{array} .$$


Proof.









by Proposition $\underline{\underline{1.0.13(i)}}$







Similarly, we can show that $id_D * S_D = \epsilon_D \eta_D$. $\square$

Proposition 3.1.6. *Let $D = A \bowtie H$ be a bialgebra, and let A and H be Hopf algebras with antipodes S_A and S_H respectively. Then*

Proof.

We can easily check Lemma 3.1.7, Corollary 3.1.8 and Corollary 3.1.9 by Theorem 3.1.4 and by simple computation.

Lemma 3.1.7. (i) *If ϕ and ψ are trivial, then (B1)–(B5) hold iff (M4) holds;*
(ii) *If α and β are trivial, then (B1)–(B5) hold iff (CM4) holds.*

Corollary 3.1.8. (i) *$A \bowtie H$ is a bialgebra iff (M1)–(M4) hold;*
(ii) *$A^\phi \bowtie^\psi H$ is a bialgebra iff (CM1)–(CM4) hold.*

In fact, [80] already contains the essence of Corollary 3.1.8.




Corollary 3.1.9. (i) *If β and ψ are trivial, then biproduct $A_\alpha^\phi \bowtie H$ is a bialgebra iff (A, α) is an H -module algebra and (A, ϕ) is an H -comodule coalgebra, (B1) and (B6) hold;*

(ii) *If α and ϕ are trivial, then biproduct $A \bowtie_\beta^\psi H$ is a bialgebra iff (H, β) is an A -module algebra and (H, ψ) is an A -comodule coalgebra, (B3) and (B6') hold.*

Corollary 3.1.10. (i) *If β and ϕ are trivial, then bicrossproduct $A_\alpha \bowtie^\psi H$ is a bialgebra iff (A, α) is an H -module algebra and (H, ψ) is an A -comodule coalgebra, (B2), (B7) and (CB4) hold;*

(ii) *If α and ψ are trivial then bicrossproduct $A^\phi \bowtie_\beta H$ is a bialgebra iff (H, β) is an A -module algebra and (A, ϕ) is an H -comodule coalgebra, (B4), (B7') and (CB2) hold.*

Proof. (i) The necessity is clear. For the sufficiency, we only show that (B5) holds since the others can similarly be shown.



Thus (B5) holds.

(ii) It is a dual case of Part (i). $\square$

Lemma 3.1.11. *Let H be a Hopf algebra, and A and H have left duality A^* and H^* respectively.*


- (i) *If (A, α) is a left H -module coalgebra, then (A^*, α^*) is a right H^* -comodule algebra;*
- (ii) *If (A, ϕ) is a left H -comodule algebra, then (A^*, ϕ^*) is a right H^* -module coalgebra.*

Proof. We only show part (i), since part (ii) is the dual case of part (i).

The left side of (1) =

=

and


The right side of (1) = 


since (A, α) is an H -module the left hand of (1). Thus (1) holds.


It is clear that

$$\begin{array}{c} A^* \\ \downarrow \\ \alpha^* \\ \downarrow \\ A^* \end{array} \quad \begin{array}{c} \epsilon \\ \downarrow \\ A^* \end{array} = \begin{array}{c} A^* \\ \downarrow \\ A^* \end{array}.$$

Thus (A^*, α^*) is a right H^* -comodule.

Now we show that (2)

The left side of (2) = 

=  by Lem.1.0.12 = the right side of (2) .

Thus (2) holds. Obviously, η_{A^*} is a comodule morphism.

Consequently, (A^*, α^*) is a right H^* -comodule algebra. $\square$

Proposition 3.1.12. *Let A and H have left dualities in braided tensor category $(\mathcal{C}, C)$. If $D = A \varphi_\alpha \bowtie_\beta^\psi H$ is a bialgebra or Hopf algebra, then*

$$(A \varphi_\alpha \bowtie_\beta^\psi H)^* \cong H^* \beta^* \bowtie_{\varphi^*}^{\alpha^*} A^* \quad \text{as bialgebras or Hopf algebras}.$$

Proof. By [82, Proposition 2.4] or Proposition 1.0.13 (ii), D^* , H^* and A^* are Hopf algebras. By Lemma 3.1.11, we have that (A^*, α^*) is a right H^* -comodule algebra, (H^*, β^*) is a left A^* -comodule algebra, (A^*, ϕ^*) is a right H^* -module coalgebra and (H^*, ψ^*) is a left H^* -module coalgebra. Let $\bar{D} = H^* \beta^* \bowtie_{\varphi^*}^{\alpha^*} A^*$. Obviously, the unit and counit of $\bar{D}$ are the same of D^* .

Now we show that their multiplication are the same.

The diagram shows the equality of two expressions for the multiplication of D^* and D^* . On the left, a small diagram shows two vertical lines labeled D^* entering a circle labeled m_D , with a single vertical line labeled D^* exiting. This is equal to a larger diagram with four vertical lines at the top labeled $D^* D^*$. Two lines cross and then merge into two lines that enter a circle labeled m_D , with a single line labeled D^* exiting. This is equal to a complex diagram with four vertical lines at the top labeled $H^* A^* H^* A^*$. These lines enter a series of circles labeled ϕ and φ , which are connected by horizontal lines. The final output is two vertical lines labeled $H^* A^*$.

and

The diagram shows the equality of two expressions for the multiplication of $\bar{D}$ and $\bar{D}$. On the left, a small diagram shows two vertical lines labeled $\bar{D}$ entering a circle labeled $m_{\bar{D}}$, with a single vertical line labeled $\bar{D}$ exiting. This is equal to a diagram with four vertical lines at the top labeled $H^* A^* H^* A^*$. These lines enter a series of circles labeled Δ_A , Δ_H , φ^* , and ϕ^* , which are connected by horizontal lines. The final output is two vertical lines labeled H^* and A^* . This is equal to a complex diagram with four vertical lines at the top labeled $H^* A^* H^*$. These lines enter a series of circles labeled ϕ and φ , which are connected by horizontal lines. The final output is two vertical lines labeled H^* and A^* .

Thus the two multiplications are the same. Dually, we have that their comultiplications are the same. Similarly, we can show that their antipodes are the same. Consequently, D^* and $\bar{D}$ are Hopf algebra isomorphic. $\square$

We can immediately get the following by Proposition 3.1.12

Corollary 3.1.13. *Let A and H have left dualities in braided tensor category $(\mathcal{C}, C)$.*

(i) *If $D = A_\alpha \bowtie_\beta H$ is a bialgebra or Hopf algebra, then*

$$(A_\alpha \bowtie_\beta H)^* \cong H^{*\beta*} \bowtie^{\alpha*} A^* \quad \text{as bialgebras or Hopf algebras ;}$$

(ii) *If $D = A^\varphi \bowtie^\psi H$ is a bialgebra or Hopf algebra, then*

$$(A^\varphi \bowtie^\psi H)^* \cong H^* \bowtie_{\psi^* \bowtie_{\varphi^*}} A^* \quad \text{as bialgebras or Hopf algebras .}$$

Now, we give an example of double cross coproduct in a strictly braided tensor category.

Example 3.1.14. *In this example, we work in the braided tensor category $\text{Vect}(\mathbf{C})$ of vector spaces over complex field $\mathbf{C}$, where its tensor is the usual tensor, and its braiding is the usual twist map. Let $\mathbf{Z}_n$ be a finite cycle group of order n , generated by element g , and let $\mathbf{CZ}_n$ be its group algebra. We have that*

$$R = \frac{1}{n} \sum_{a,b=0}^{n-1} e^{-\frac{2\pi i ab}{n}} g^a \otimes g^b$$

is a quasitriangular structure of $\mathbf{CZ}_n$ and its module category $\mathbf{CZ}_n \mathcal{M}$ becomes a braided tensor category . Furthermore, the category is strictly braided for $n > 2$ (see [83, Example 2.1.6 and 9.2.5]). Assume $n > 2$ in the following. By [26, Theorem 2.2 and 2.5], there is a quasitriangular structure $\bar{R}$ of $\mathbf{CZ}_n \bowtie^R \mathbf{CZ}_n$ and $\bar{R}$ is not triangular. Let $D = \mathbf{CZ}_n \bowtie^R \mathbf{CZ}_n$. Thus $({}_D \mathcal{M}, C^R)$ is a strictly braided tensor category by [64, Proposition XIII.1.4]. It follows from [83, Example 9.4.9] (transmutation) that there is a Hopf algebra $\underline{D}$ living in the strictly braided tensor category ${}_D \mathcal{M}$, where $\underline{D}$ is given by the same algebra, unit and counit as D , and comultiplication

$$\underline{\Delta}(b) = \sum b_1 S_D(\bar{R}^{(2)}) \otimes (\bar{R}^{(1)} \triangleright b_2)$$

for any $b \in \underline{D}$; notice that “ $\triangleright$ ” is the quantum action. Since $\underline{D}$ is commutative we have that $\bar{R}^{(1)} \cdot b_2 = \epsilon(R^{(1)})b_2$ and

$$\underline{\Delta}(b) = \sum b_1 S_D(\bar{R}^{(2)}) \otimes (\bar{R}^{(1)} \triangleright b_2) = \sum b_1 \otimes b_2 = \Delta(b).$$

That is, $\underline{D} = D$ as Hopf algebra. Therefore there are non-trivial ϕ and non-trivial ψ such that $\mathbf{CZ}_n \bowtie^R \mathbf{CZ}_n = \mathbf{CZ}_n^\phi \bowtie^\psi \mathbf{CZ}_n$. Furthermore $\mathbf{CZ}_n^\phi \bowtie^\psi \mathbf{CZ}_n$ is a double cross coproduct in the strictly braided tensor category ${}_D\mathcal{M}$ by [26, Lemma 1.3].

3.2 The universal property of double bicrossproduct

In this section, we give the universal property of double bicrossproduct. Throughout this section, H and A are two bialgebras in braided tensor categories, (A, α) is a left H -module coalgebra, (H, β) is a right A -module coalgebra, (A, ϕ) is a left H -comodule algebra and (H, ψ) is a right H -comodule algebra.

Lemma 3.2.1. *Let B and E be bialgebras, p be a coalgebra morphism from B to D , j be an algebra morphism from D to E , where $D = A_\alpha^\phi \bowtie_\beta^\psi H$ is a bialgebra. Then*

- (i) $(\pi_A \otimes \pi_H)\Delta_D = id_D$, where π_A and π_H are trivial actions;
- (ii) *there are two coalgebra morphisms p_A from B to A and p_H from B to H such that*

$$p = (p_A \otimes p_H)\Delta_B.$$

Furthermore, if there are two coalgebra morphisms f from B to A and g from B to H such that

$$p = (f \otimes g)\Delta_B,$$

then $f = p_A = \pi_A p$ and $g = p_H = \pi_H p$.

- (iii) *If two coalgebra morphisms f and g from B to D satisfy*

$$\pi_A f = \pi_A g, \quad \pi_H f = \pi_H g,$$

then $f = g$;

- (iv) $m_D(i_A \otimes i_H) = id_D$;

- (v) *there are two algebra morphisms j_A from A to E and j_H from H to E such that*

$$j = m_E(j_A \otimes j_H).$$

Furthermore, if there are two algebra morphisms f from A to E and g from H to E such that

$$j = m_E(f \otimes g),$$


then $f = j_A = ji_A$ and $g = j_H = ji_H$.

- (vi) *If two algebra morphisms f and g from D to E satisfy*

$$fi_A = gi_A, \quad fi_H = gi_H,$$

then $f = g$.

Proof. (i)



(ii) Let $p_A = \pi_A p$ and $p_H = \pi_H p$, then p_A and p_H are coalgebra morphisms and

$$p = (\pi_A \otimes \pi_H) \Delta_D p = (\pi_A \otimes \pi_H)(p \otimes p) \Delta_B = (p_A \otimes p_H) \Delta_B$$

by part (i). On the other hand, if

$$p = (f \otimes g) \Delta_B,$$

then $p_A = \pi_A p = (id_A \otimes \epsilon_H)(f \otimes g) \Delta_B = f$ and $p_H = g$.


(iii) We see that


$$f = (\pi_A \otimes \pi_H) \Delta_D f = (\pi_A \otimes \pi_H)(f \otimes f) \Delta_B = (\pi_A \otimes \pi_H)(g \otimes g) \Delta_B = (\pi_A \otimes \pi_H) \Delta_B g = g.$$


Therefore $f = g$.


(iv)–(vi) can dually be shown. $\square$


Assume that p_A, p_H, j_A and j_H are morphisms from B to A , from B to H , from A to B and from H to B , respectively. Set



[u3]:  ;

[u4]:  ;


[u5]:  ;

[u6]:  .

Lemma 3.2.2. Let $D = A_\alpha^\phi \bowtie_\beta^\psi H$, B be a bialgebra and $p = (p_A \otimes p_H)\Delta_B$, where p_A and p_H are coalgebra morphisms from B to A and from B to H respectively. Then

- (i) p is a coalgebra morphism iff (u1) holds;
- (ii) p is an algebra morphism iff (u2) and (u3) hold;
- (iii) p is a bialgebra morphism iff (u1)–(u3) hold.

Proof. (i) We see that

(1)

If p is a coalgebra morphism, then by relation (1) and (2), we have that

i.e.

Thus (u1) holds.

Conversely, if (u1) holds, by relation (1) and (2), we have that

The diagrammatic equations are as follows:

$$\begin{array}{c} B \\ \circlearrowleft p \\ | \\ \text{---} \circlearrowleft \text{---} \\ D \quad D \end{array} = \begin{array}{c} B \\ | \quad | \\ \circlearrowleft p \quad \circlearrowleft p \\ | \quad | \\ D \quad D \end{array} \quad \text{and} \quad \begin{array}{c} B \\ \circlearrowleft p \\ | \\ \text{---} \circlearrowleft \text{---} \\ \epsilon_D \end{array} = \begin{array}{c} B \\ | \\ \text{---} \circlearrowleft \text{---} \\ \epsilon_B \end{array}$$

Thus p is a coalgebra morphism.

(ii) We first see that

and

and

If p is an algebra morphism, by the above proof, we have that

and

In this way, we can obtain (u2) and (u3).

Conversely, if (u1) and (u2) hold, we can similarly show that p is an algebra morphism.

(iii) It follows from part (i) and part (ii). $\square$

Dually, we have

Lemma 3.2.3. *Let B be a bialgebra and $j = m_B(j_A \otimes j_H)$, where j_A and j_H are algebra morphisms from A to B and from H to B respectively. Then*

- (i) j is an algebra morphism iff (u4) holds;
- (ii) j is a coalgebra morphism iff (u5) and (u6) hold;
- (iii) j is a bialgebra morphism iff (u4)–(u6) hold.

Definition 3.2.4. (UT1): Let B be a bialgebra and two coalgebra morphisms

$$p_A : B \longrightarrow A, \quad p_H : B \longrightarrow H$$

satisfy conditions (u1)–(u3). (B, p_A, p_H) is said to satisfy condition (UT1), if for any bialgebra E and any two coalgebra morphisms

$$q_A : E \rightarrow A, q_H : E \rightarrow H,$$

which satisfy conditions (u1)–(u3), there exists a unique bialgebra morphism q' from E to B such that

$$p_A q' = q_A, \quad p_H q' = q_H,$$

i.e. the diagram

$$\begin{array}{ccc} & A & \\ q_A \nearrow & & \nwarrow p_A \\ E & \xrightarrow{q'} & B \\ q_H \searrow & & \swarrow p_H \\ & H & \end{array}$$

commutes.

(UT2): Let B and $D = A_\alpha^\phi \bowtie_\beta^\psi H$ be bialgebras and p a bialgebra morphism from B to D . (B, p) is said to satisfy condition (UT2) if for any bialgebra E and any bialgebra morphism q from E to D , there exists a unique bialgebra morphism q' from E to B such that the diagram

$$\begin{array}{ccc} & D & \\ q \nearrow & & \nwarrow p \\ E & \xrightarrow{q'} & B \end{array}$$

commutes.

(UT3): Let B be coalgebra and two coalgebra morphisms

$$p_A : B \longrightarrow A, \quad p_H : B \longrightarrow H$$

satisfy condition (u1). (B, p_A, p_H) is said to satisfy condition (UT3), if for any coalgebra E and any two coalgebra morphisms

$$q_A : E \rightarrow A, q_H : E \rightarrow H,$$

which satisfy condition (u1), there exists a unique coalgebra morphism q' from E to B such that

$$p_A q' = q_A, \quad p_H q' = q_H,$$

i.e. the diagram

$$\begin{array}{ccc} & A & \\ q_A \nearrow & & \nwarrow p_A \\ E & \xrightarrow{q'} & B \\ q_H \searrow & & \swarrow p_H \\ & H & \end{array}$$

commutes.

(UT4): Let B and $D = A \overset{c}{\bowtie} H$ be coalgebras and p a coalgebra morphism from B to D . (B, p) is said to satisfy condition (UT4) if for any coalgebra E and any coalgebra morphism q from E to D , there exists a unique coalgebra morphism q' from E to B such that the diagram

$$\begin{array}{ccc} & D & \\ q \nearrow & & \nwarrow p \\ E & \cdots \xrightarrow{q'} & B \end{array}$$

commutes.

(UT1'): Let B be a bialgebra and two algebra morphisms

$$j_A : A \longrightarrow B, \quad j_H : H \longrightarrow B$$

satisfy conditions (u4)–(u6). (B, j_A, j_H) is said to satisfy condition (UT1'), if for any bialgebra E and any two algebra morphisms

$$q_A : A \rightarrow E, q_H : H \rightarrow E,$$

which satisfy conditions (u4)–(u6), there exists a unique bialgebra morphism q' from B to E such that

$$q'j_A = q_A, \quad q'j_H = q_H,$$

i.e. the diagram

$$\begin{array}{ccccc} & & A & & \\ q_A \swarrow & & & & \searrow j_A \\ E & & & & B \\ q_H \swarrow & \cdots \xrightarrow{q'} & & & \searrow j_H \\ & & H & & \end{array}$$

commutes.

(UT2'): Let B and $D = A \overset{\phi}{\bowtie}_{\alpha} \overset{\psi}{\bowtie}_{\beta} H$ be bialgebras and j a bialgebra morphism from D to B . (B, j) is said to satisfy condition (UT2') if for any bialgebra E and any bialgebra morphism q from D to E , there exists a unique bialgebra morphism q' from B to E such that the diagram $q'j = q$.

(UT3'): Let B be algebra and two algebra morphisms

$$j_A : A \longrightarrow B, \quad j_H : H \longrightarrow B$$

satisfy condition (u1). (B, j_A, j_H) is said to satisfy condition (UT3'), if for any algebra E and any two algebra morphisms

$$q_A : A \rightarrow E, q_H : H \rightarrow E,$$

which satisfy condition (u4), there exists a unique algebra morphism q' from B to E such that

$$q'j_A = q_A, \quad q'j_H = q_H.$$

(UT4'): Let B and $D = A \bowtie H$ be algebras and j an algebra morphism from D to B . (B, j) is said to satisfy condition (UT4'), if for any algebra E and any algebra morphism q from D to E , there exists a unique algebra morphism q' from B to E such that the diagram $q = q'j$.

Theorem 3.2.5. Let B and $D = A_{\alpha}^{\phi} \bowtie_{\beta}^{\psi} H$ be bialgebras.

- (I) (i) (D, π_A, π_H) satisfies condition (UT1);
- (ii) (D, id_D) satisfies condition (UT2);
- (iii) (D, i_A, i_H) satisfies condition (UT1');
- (iv) (D, id_D) satisfies condition (UT2').
- (II) The following statements are equivalent;
- (i) (B, p_A, p_H) satisfies condition (UT1);
- (ii) There exists a (necessarily unique) bialgebra isomorphism p' from B to D such that

$$\pi_A p' = p_A, \quad \pi_H p' = p_H.$$

- (iii) p is a bialgebra isomorphism from B to D and $p = (p_A \otimes p_H) \Delta_B$;
- (iv) (B, p) satisfies condition (UT2) and $p = (p_A \otimes p_H) \Delta_B$.
- (III) The following statements are equivalent.
- (i) (B, j_A, j_H) satisfies condition (UT1');
- (ii) There exists a (necessarily unique) bialgebra isomorphism j' from D to B such that

$$j' i_A = j_A, \quad j' i_H = j_H;$$

- (iii) j is a bialgebra isomorphism from D to B and $j = m_B(j_A \otimes j_H)$;
- (iv) (B, j) satisfies condition (UT2') and $j = m_B(j_A \otimes j_H)$.

Proof. (I) (i) For any bialgebra E and two coalgebra morphisms

$$q_A : E \longrightarrow A, \quad q_H : E \longrightarrow H,$$

which satisfy conditions (u1)–(u3), set

$$q' = (q_A \otimes q_H) \Delta_E.$$

It is clear that

$$\pi_A q' = q_A, \quad \pi_H q' = q_H$$

The uniqueness of q' can be obtained by Lemma 3.2.1 (iii). Otherwise, $id_D = (\pi_A \otimes \pi_H) \Delta_D$ by Lemma 3.2.1 (i) and π_A and π_H satisfy conditions (u1)–(u3) by Lemma 3.2.2 (iii). Thus (D, π_A, π_H) satisfies condition (TU1).

(ii) For any bialgebra E and any bialgebra morphism q from E to D , let $q' = q$. Obviously, $id_D q' = q$ and we have the uniqueness of q' . Thus (D, id_D) satisfies condition (UT2).

(iii) and (iv) can dually be checked.

(II) (i) $\Rightarrow$ (ii) Since (D, π_A, π_H) and (B, p_A, p_H) satisfy condition (UT1), we have that there exist two bialgebra morphisms

$$f : B \rightarrow D \quad \text{and} \quad g : D \rightarrow B$$

such that

$$p_A g = \pi_A, p_H g = \pi_H, \pi_A f = p_A, \pi_H f = p_H.$$

Now we see that

$$\pi_A f g = \pi_A, \pi_H f g = \pi_H, p_A g f = p_A, p_H g f = p_H.$$

Therefore, by uniqueness,

$$f g = id_D \quad \text{and} \quad g f = id_B.$$

That is, f is a bialgebra isomorphism from B to D .

(ii) $\Rightarrow$ (iii) Let $p = (p_A \otimes p_H) \Delta_B$. Since

$$\pi_A p' = p_A, \quad \pi_H p' = p_H, \quad \pi_A p = p_A, \quad \pi_H p = p_H,$$

considering Lemma 3.2.1 (iii), we have that

$$p = p'.$$

Therefore p is a bialgebra isomorphism.

(iii) $\Rightarrow$ (iv) For any bialgebra E and any bialgebra morphism q from E to D , let

$$q' = p^{-1} q.$$

Obviously, $p q' = q$ and we have the uniqueness of q' . Thus (B, p) satisfies condition (UT2).

(iv) $\Rightarrow$ (i) For any bialgebra E and two coalgebra morphisms

$$q_A : E \rightarrow A, q_H : E \rightarrow H,$$

which satisfy conditions (u1)–(u3), set $q = (q_A \otimes q_H) \Delta_E$. By Lemma 3.2 (iii), we have that q is a bialgebra morphism from E to D . Thus there exists a unique bialgebra morphism q' from E to B such that

$$p q' = q$$

since (B, p) satisfies condition (UT2). It is clear that

$$p_A q' = q_A, \quad p_H q' = q_H.$$

Otherwise, p_A and p_H satisfy conditions (u1)–(u3) by Lemma 3.2.2 (iii). Therefore, (B, p_A, p_H) satisfies condition (UT1).

(III) It can dually be obtained.

We can easily find the following two theorems from the proof of the previous theorem.

Theorem 3.2.6. *Let B and $D = A \overset{c}{\bowtie} H$ be coalgebras.*

(I) (i) (D, π_A, π_H) satisfies condition (UT3);

(ii) (D, id_D) satisfies condition (UT4);

(II) The following statements are equivalent

(i) (B, p_A, p_H) satisfies condition (UT3).

(ii) There exists a (necessarily unique) coalgebra isomorphism p' from B to D such that

$$\pi_A p' = p_A, \quad \pi_H p' = p_H;$$

(iii) p is a coalgebra isomorphism from B to D and $p = (p_A \otimes p_H)\Delta_B$.

(iv) (B, p) satisfies condition (UT4) and $p = (p_A \otimes p_H)\Delta_B$.

Theorem 3.2.7. Let B and $D = A \bowtie H$ be algebras.

(I) (i) (D, i_A, i_H) satisfies condition (UT3');

(ii) (D, id_D) satisfies condition (UT4');

(II) The following statements are equivalent.

(i) (B, j_A, j_H) satisfies condition (UT3').

(ii) There exists an (necessarily unique) algebra isomorphism j' from D to B such that

$$j' i_A = j_A, \quad j' i_H = j_H;$$

(iii) j is an algebra isomorphism from D to B and $j = m_B(j_A \otimes j_H)$.

(iv) (B, j) satisfies condition (UT4') and $j = m_B(j_A \otimes j_H)$.

Chapter 4

The Bicrossproducts in Braided Tensor Categories

In the category of usual vector spaces with usual twist braiding, S. Majid introduced bicrossproducts and gave the necessary and sufficient conditions for them to become bialgebras in [79, Theorem 2.9]. D.E. Radford introduced biproducts and gave the necessary and sufficient conditions for them to become bialgebras in [99, Theorem 1]. In braided tensor category, S. Majid introduced the smash product and showed that if (A, α) is an H -module bialgebra and H is cocommutative with respect to (H, α) , then the smash product $A_\alpha \# H$ is a bialgebra in [80, Theorem 2.4].

In this Chapter, we construct the bicrossproducts and biproducts in braided tensor category $(\mathcal{C}, C)$ and give the necessary and sufficient conditions for them to be bialgebras. We obtain that for any left module (A, α) of bialgebra H the smash product $A \# H$ is a bialgebra iff A is an H -module bialgebra and H is cocommutative with respect to (A, α) .

Now we give some concepts as follows: assume that $H, A \in \text{ob } \mathcal{C}$, and that

$$\begin{aligned}
 \alpha : H \otimes A &\rightarrow A \quad , \quad \beta : H \otimes A \rightarrow H, \\
 \phi : A &\rightarrow H \otimes A \quad , \quad \psi : H \rightarrow H \otimes A, \\
 m_H : H \otimes H &\rightarrow H \quad , \quad m_A : A \otimes A \rightarrow A, \\
 \Delta_H : H &\rightarrow H \otimes H \quad , \quad \Delta_A : A \rightarrow A \otimes A, \\
 \eta_H : I &\rightarrow H \quad , \quad \eta_A : I \rightarrow A, \\
 \epsilon_H : H &\rightarrow I \quad , \quad \epsilon_A : A \rightarrow I. \\
 \sigma : H \otimes H &\rightarrow A \quad , \quad \mu : A \otimes A \rightarrow H, \\
 P : A &\rightarrow H \otimes H \quad , \quad Q : H \rightarrow A \otimes A
 \end{aligned}$$

are morphisms in $\mathcal{C}$.

(H, α) is said to act weakly on A if the following conditions are satisfied:

$$(WA) : \begin{array}{c} H \quad A \quad A \\ | \quad | \quad | \\ \text{---} \alpha \text{---} \\ | \\ A \end{array} = \begin{array}{c} H \quad A \quad A \\ | \quad | \quad | \\ \text{---} \alpha \text{---} \quad \text{---} \alpha \text{---} \\ | \quad \quad | \\ A \quad \quad A \end{array}, \quad \begin{array}{c} H \\ | \\ \text{---} \eta_A \text{---} \\ | \\ A \end{array} = \begin{array}{c} H \\ | \\ \epsilon \\ | \\ A \end{array}, \quad \begin{array}{c} A \\ | \\ \text{---} \eta_H \text{---} \\ | \\ A \end{array} = \begin{array}{c} A \\ | \\ A \end{array}.$$

(A, β) is said to act weakly on H if the following conditions are satisfied:

$$(WA) : \begin{array}{c} H \quad H \quad A \\ | \quad | \quad | \\ \text{---} m \text{---} \\ | \\ H \end{array} = \begin{array}{c} H \quad H \quad A \\ | \quad | \quad | \\ \text{---} \beta \text{---} \quad \text{---} \beta \text{---} \\ | \quad \quad | \\ H \quad \quad A \end{array}, \quad \begin{array}{c} A \\ | \\ \text{---} \eta_H \text{---} \\ | \\ H \end{array} = \begin{array}{c} A \\ | \\ \epsilon \\ | \\ H \end{array}, \quad \begin{array}{c} H \\ | \\ \text{---} \eta_A \text{---} \\ | \\ H \end{array} = \begin{array}{c} A \\ | \\ H \end{array}.$$

(H, ϕ) is said to coact weakly on A if the following conditions are satisfied:

$$(WCA) : \begin{array}{c} A \\ | \\ \text{---} \phi \text{---} \\ | \\ H \quad A \quad A \end{array} = \begin{array}{c} A \\ | \\ \text{---} \phi \text{---} \quad \text{---} \phi \text{---} \\ | \quad \quad | \\ H \quad \quad A \quad \quad A \end{array}, \quad \begin{array}{c} A \\ | \\ \text{---} \phi \text{---} \\ | \\ H \end{array} = \begin{array}{c} A \\ | \\ \epsilon \\ | \\ H \end{array}, \quad \begin{array}{c} A \\ | \\ \text{---} \phi \text{---} \\ | \\ A \end{array} = \begin{array}{c} A \\ | \\ A \end{array}.$$

(A, ψ) is said to coact weakly on H if the following conditions are satisfied:

$$(WCA) : \begin{array}{c} H \\ | \\ \text{---} \psi \text{---} \\ | \\ H \quad H \quad A \end{array} = \begin{array}{c} H \\ | \\ \text{---} \psi \text{---} \quad \text{---} \psi \text{---} \\ | \quad \quad | \\ H \quad \quad H \quad \quad A \end{array}, \quad \begin{array}{c} H \\ | \\ \text{---} \psi \text{---} \\ | \\ A \end{array} = \begin{array}{c} H \\ | \\ \epsilon \\ | \\ A \end{array}, \quad \begin{array}{c} H \\ | \\ \text{---} \psi \text{---} \\ | \\ H \end{array} = \begin{array}{c} H \\ | \\ H \end{array}.$$

σ is called a 2-cocycle from $H \otimes H$ to A if the following conditions are satisfied:

$$\begin{aligned}
 (2-COC) : \quad & \begin{array}{c} \text{Diagram 1: Three vertical lines from } H \text{ at the top. The left and middle lines cross at a point. Below the crossing, the left line goes to a box labeled } \alpha, \text{ and the middle line goes to a box labeled } \sigma. \text{ The right line goes to a box labeled } \sigma. \text{ All three boxes } \alpha, \sigma, \sigma \text{ are connected to a single line labeled } A \text{ at the bottom.} \end{array} \\
 & = \begin{array}{c} \text{Diagram 2: Three vertical lines from } H \text{ at the top. The left and middle lines cross. Below the crossing, the left line goes to a box labeled } \sigma, \text{ and the middle line goes to a box labeled } \sigma. \text{ The right line goes to a box labeled } \sigma. \text{ All three boxes } \sigma, \sigma, \sigma \text{ are connected to a single line labeled } A \text{ at the bottom.} \end{array}, \\
 \\
 & \begin{array}{c} \text{Diagram 3: A vertical line from } H \text{ at the top goes to a box labeled } \eta_H, \text{ then to a box labeled } \sigma, \text{ and finally to } A. \end{array} \\
 & = \begin{array}{c} \text{Diagram 4: A vertical line from } H \text{ at the top goes to a box labeled } \eta_H, \text{ then to a box labeled } \sigma, \text{ and finally to } A. \end{array} \\
 & = \begin{array}{c} \text{Diagram 5: A vertical line from } H \text{ at the top goes to a box labeled } \epsilon, \text{ then to a box labeled } \eta_A, \text{ and finally to } A. \end{array}.
 \end{aligned}$$

μ is called a 2-cocycle from $A \otimes A$ to H if the following conditions are satisfied:

$$\begin{aligned}
 (2-COC) : \quad & \begin{array}{c} \text{Diagram 1: Three vertical lines from } A \text{ at the top. The left and middle lines cross. Below the crossing, the left line goes to a box labeled } \mu, \text{ and the middle line goes to a box labeled } \mu. \text{ The right line goes to a box labeled } \mu. \text{ All three boxes } \mu, \mu, \mu \text{ are connected to a single line labeled } H \text{ at the bottom.} \end{array} \\
 & = \begin{array}{c} \text{Diagram 2: Three vertical lines from } A \text{ at the top. The left and middle lines cross. Below the crossing, the left line goes to a box labeled } \mu, \text{ and the middle line goes to a box labeled } \mu. \text{ The right line goes to a box labeled } \beta. \text{ All three boxes } \mu, \mu, \beta \text{ are connected to a single line labeled } H \text{ at the bottom.} \end{array}, \\
 \\
 & \begin{array}{c} \text{Diagram 3: A vertical line from } A \text{ at the top goes to a box labeled } \eta_A, \text{ then to a box labeled } \mu, \text{ and finally to } H. \end{array} \\
 & = \begin{array}{c} \text{Diagram 4: A vertical line from } A \text{ at the top goes to a box labeled } \eta_A, \text{ then to a box labeled } \mu, \text{ and finally to } H. \end{array} \\
 & = \begin{array}{c} \text{Diagram 5: A vertical line from } A \text{ at the top goes to a box labeled } \epsilon, \text{ then to a box labeled } \eta_H, \text{ and finally to } H. \end{array}.
 \end{aligned}$$

P is called a 2-cycle from A to $H \otimes H$ if the following conditions are satisfied:

$$\begin{aligned}
 (2-C) : \quad & \begin{array}{c} \text{Diagram 1: A vertical line from } A \text{ at the top splits into two lines. The left line goes to a box labeled } \phi, \text{ and the right line goes to a box labeled } P. \text{ Below } \phi, \text{ the line goes to a box labeled } P. \text{ Below } P, \text{ the line goes to a box labeled } P. \text{ The right line from } P \text{ goes to a box labeled } P. \text{ All four boxes } \phi, P, P, P \text{ are connected to three vertical lines labeled } H \text{ at the bottom.} \end{array} \\
 & = \begin{array}{c} \text{Diagram 2: A vertical line from } A \text{ at the top splits into two lines. The left line goes to a box labeled } P, \text{ and the right line goes to a box labeled } P. \text{ Below } P, \text{ the line goes to a box labeled } P. \text{ Below } P, \text{ the line goes to a box labeled } P. \text{ The right line from } P \text{ goes to a box labeled } P. \text{ All four boxes } P, P, P, P \text{ are connected to three vertical lines labeled } H \text{ at the bottom.} \end{array},
 \end{aligned}$$

$$\begin{array}{c} A \\ | \\ \textcircled{P} \\ | \quad | \\ \textcircled{\epsilon} \quad | \\ H \end{array} = \begin{array}{c} A \\ | \\ \textcircled{P} \\ | \quad | \\ | \quad \textcircled{\epsilon} \\ H \end{array} = \begin{array}{c} A \\ | \\ \textcircled{\epsilon} \\ | \\ \textcircled{\eta_H} \\ | \\ H \end{array} .$$

Q is called a 2-cycle from H to $A \otimes A$ if the following conditions are satisfied:

$$\begin{array}{c} (2-C) : \end{array} \begin{array}{c} H \\ | \\ \textcircled{\alpha} \quad | \\ | \quad | \\ \textcircled{\alpha} \quad | \\ | \quad | \\ A \quad A \quad A \end{array} = \begin{array}{c} H \\ | \\ \textcircled{Q} \quad | \quad \textcircled{\psi} \\ | \quad | \quad | \\ \textcircled{Q} \quad | \quad | \\ | \quad | \quad | \\ A \quad A \quad A \end{array} ,$$

$$\begin{array}{c} H \\ | \\ \textcircled{Q} \\ | \quad | \\ \textcircled{\epsilon} \quad | \\ A \end{array} = \begin{array}{c} H \\ | \\ \textcircled{Q} \\ | \quad | \\ | \quad \textcircled{\epsilon} \\ A \end{array} = \begin{array}{c} H \\ | \\ \textcircled{\epsilon} \\ | \\ \textcircled{\eta_H} \\ | \\ H \end{array} .$$

(A, α) is called a twisted H -module if the following conditions are satisfied:

$$\begin{array}{c} (TM) : \end{array} \begin{array}{c} H \quad H \quad A \\ | \quad | \quad | \\ \textcircled{\alpha} \quad | \quad | \\ | \quad | \quad | \\ \textcircled{\alpha} \quad | \quad | \\ | \quad | \quad | \\ A \end{array} = \begin{array}{c} H \quad H \quad A \\ | \quad | \quad | \\ \textcircled{\sigma} \quad | \quad | \\ | \quad | \quad | \\ \textcircled{\alpha} \quad | \quad | \\ | \quad | \quad | \\ A \end{array} ,$$

$$\begin{array}{c} A \\ | \\ \textcircled{\eta_H} \\ | \\ \textcircled{\alpha} \\ | \\ A \end{array} = \begin{array}{c} A \\ | \\ | \\ | \\ A \end{array} .$$

(H, β) is called a twisted A -module if the following conditions are satisfied:

$$\begin{aligned}
 (TM) : \quad & \begin{array}{c} \text{Diagram 1: } H \text{ line on left, } A \text{ line on top, } A \text{ line on right. } \beta \text{ on } H \text{ line, } \mu \text{ on } A \text{ line.} \\ \text{Diagram 2: } H \text{ line on left, } A \text{ line on top, } A \text{ line on right. } \beta \text{ on } A \text{ line, } \mu \text{ on } H \text{ line.} \end{array} = \\
 & \begin{array}{c} \text{Diagram 3: } H \text{ line on left, } A \text{ line on top, } A \text{ line on right. } \beta \text{ on } H \text{ line, } \eta_A \text{ on } A \text{ line.} \\ \text{Diagram 4: } H \text{ line on left, } A \text{ line on top, } A \text{ line on right. } \beta \text{ on } A \text{ line, } \eta_A \text{ on } H \text{ line.} \end{array} .
 \end{aligned}$$

(A, ϕ) is called a twisted H -comodule if the following conditions are satisfied:

$$\begin{aligned}
 (TCM) : \quad & \begin{array}{c} \text{Diagram 1: } A \text{ line on top, } H \text{ line on left, } H \text{ line on right. } \phi \text{ on } A \text{ line, } P \text{ on } H \text{ line.} \\ \text{Diagram 2: } A \text{ line on top, } H \text{ line on left, } H \text{ line on right. } \phi \text{ on } H \text{ line, } P \text{ on } A \text{ line.} \end{array} = \\
 & \begin{array}{c} \text{Diagram 3: } A \text{ line on top, } H \text{ line on left, } H \text{ line on right. } \phi \text{ on } H \text{ line, } P \text{ on } A \text{ line.} \\ \text{Diagram 4: } A \text{ line on top, } H \text{ line on left, } H \text{ line on right. } \phi \text{ on } A \text{ line, } P \text{ on } H \text{ line.} \end{array} , \\
 & \begin{array}{c} \text{Diagram 5: } A \text{ line on top, } A \text{ line on bottom. } \phi \text{ on } A \text{ line, } \epsilon_H \text{ on } A \text{ line.} \\ \text{Diagram 6: } A \text{ line on top, } A \text{ line on bottom. } \phi \text{ on } A \text{ line.} \end{array} = .
 \end{aligned}$$

(H, ψ) is called a twisted H -comodule if the following conditions are satisfied:

$$\begin{aligned}
 (TCM) : \quad & \begin{array}{c} \text{Diagram 1: } H \text{ line on top, } H \text{ line on left, } A \text{ line on right. } \psi \text{ on } H \text{ line, } Q \text{ on } A \text{ line.} \\ \text{Diagram 2: } H \text{ line on top, } H \text{ line on left, } A \text{ line on right. } \psi \text{ on } A \text{ line, } Q \text{ on } H \text{ line.} \end{array} = \\
 & \begin{array}{c} \text{Diagram 3: } H \text{ line on top, } H \text{ line on left, } A \text{ line on right. } \psi \text{ on } H \text{ line, } Q \text{ on } A \text{ line.} \\ \text{Diagram 4: } H \text{ line on top, } H \text{ line on left, } A \text{ line on right. } \psi \text{ on } A \text{ line, } Q \text{ on } H \text{ line.} \end{array} ,
 \end{aligned}$$

H is said to be cocommutative with respect to (A, α) if the following conditions are satisfied:

A is said to be cocommutative with respect to (H, β) if the following conditions are satisfied:

H is said to be commutative with respect to (A, ϕ) if the following conditions are satisfied:

A is called commutative with respect to (H, ψ) if the following conditions are satisfied:

If (H, α) weakly acts on A such that (A, α) becomes a left H -module, then we say that (H, α) acts on A . Dually, if (H, ϕ) weakly coacts on A such that (A, ϕ) become a left H -comodule, then

we say that (H, ϕ) coacts on A . An action of H on A is called inner if there exists a morphism $u : H \rightarrow A$ with a convolution inverse $\bar{u}$ such that

Let

It is clear that (H, ad) acts on itself. This action ad is called a (left) adjoint action. Dually we can define an (left) adjoint coaction as follows:

Bialgebra A is called an H -module bialgebra if A is an H -module algebra and H -module coalgebra. Bialgebra A is called an H -comodule bialgebra if A is an H -comodule algebra and an H -comodule coalgebra.

We define the relations (BB1)–(BB11) and (BB1')–(BB11') as follows:

(BB1):

The diagram shows an equality between two string diagrams. The left diagram has two vertical strands labeled H and A at the top. The H strand goes down, loops around a node P , and then goes up to a node β . The A strand goes down, loops around a node β , and then goes up to a node P . Both strands then go down to two H labels at the bottom. The right diagram is more complex, involving nodes β , ϕ , and P , and labels H and A . To the right of the equality, there is a separate equation: $\begin{array}{c} H \quad A \\ \diagdown \quad \diagup \\ \beta \\ \downarrow \\ \epsilon \end{array} = \begin{array}{c} H \quad A \\ \downarrow \quad \downarrow \\ \epsilon \quad \epsilon \end{array};$

(BB2):

The diagram shows an equality between two string diagrams. The left diagram has two vertical strands labeled A and A at the top. The A strand goes down, loops around a node μ , and then goes up to a node ϕ . The A strand goes down, loops around a node ϕ , and then goes up to a node β . Both strands then go down to two H labels at the bottom. The right diagram is more complex, involving nodes μ , ϕ , and β , and labels A and H . To the right of the equality, there is a separate equation: $\begin{array}{c} \eta_A \\ \downarrow \\ \phi \\ \diagup \quad \diagdown \\ H \quad A \end{array} = \begin{array}{c} \eta_H \quad \eta_A \\ \downarrow \quad \downarrow \\ H \quad A \end{array};$

(BB3):

The diagram shows an equality between two string diagrams. The left diagram has two vertical strands labeled H and A at the top. The H strand goes down, loops around a node ϕ , and then goes up to a node β . The A strand goes down, loops around a node β , and then goes up to a node ϕ . Both strands then go down to two H labels at the bottom. The right diagram is more complex, involving nodes β and ϕ , and labels H and A .

(BB4) :

= ;

(BB5) :

, $\quad$.

(BB8) :

;

$$(BB9) : \quad \begin{array}{c} H \quad A \\ | \quad | \\ \text{---} Q \text{---} \\ | \quad | \\ A \quad A \end{array} = \begin{array}{c} H \quad A \\ \diagdown \quad \diagup \\ | \quad | \\ \text{---} Q \text{---} \\ | \quad | \\ A \quad A \end{array} ;$$

$$(BB1') : \quad \begin{array}{c} H \quad H \\ | \quad | \\ \text{---} \psi \text{---} \sigma \\ | \quad | \\ H \quad A \end{array} = \begin{array}{c} H \quad H \\ | \quad | \\ \text{---} \sigma \text{---} \psi \\ | \quad | \\ H \quad A \end{array} , \quad \begin{array}{c} \eta_H \\ | \\ \psi \\ | \\ H \quad A \end{array} = \begin{array}{c} \eta_H \quad \eta_A \\ | \quad | \\ H \quad A \end{array} ;$$

$$(BB2') : \quad \begin{array}{c} H \quad A \\ | \quad | \\ \text{---} \alpha \text{---} Q \\ | \quad | \\ A \quad A \end{array} = \begin{array}{c} H \quad A \\ | \quad | \\ \text{---} \psi \text{---} \alpha \\ | \quad | \\ Q \quad A \end{array} , \quad \begin{array}{c} H \quad A \\ | \quad | \\ \alpha \\ | \\ \epsilon \end{array} = \begin{array}{c} H \quad A \\ | \quad | \\ \epsilon \quad \epsilon \end{array} ;$$

$$(BB3') : \quad \begin{array}{c} H \quad A \\ | \quad | \\ \text{---} \alpha \text{---} \psi \\ | \quad | \\ A \quad H \end{array} = \begin{array}{c} H \quad A \\ | \quad | \\ \text{---} \psi \text{---} \alpha \\ | \quad | \\ H \quad A \end{array} ;$$

(BB4') :

(BB5') :

(BB8') :

$$(BB9') : \quad \begin{array}{c} A \quad A \\ \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \\ \mu \\ H \quad A \end{array} = \begin{array}{c} A \quad A \\ \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \\ \mu \\ H \quad A \end{array} ;$$

For convenience, $(BB6')$ and $(B1)$ be the same; $(BB7')$ and $(B6)$ be the same; $(BB10')$ be the same as $(CB4)$ and $\psi\eta_H = \eta_H \otimes \eta_A$; $(BB11')$ be the same as $(B2)$ and $\epsilon\alpha = \epsilon \otimes \epsilon$. Let $(BB6)$ and $(B3)$ are the same; $(BB7)$ and $(B6')$ are the same; $(BB10)$ is the same as $(B4)$ and $\epsilon\beta = \epsilon \otimes \epsilon$; $(BB11)$ is the same as $(CB2)$ and $\phi\eta_A = \eta_H \otimes \eta_A$. Here $(B_1), (B_6)$ and so on are defined in Chapter 3.

$$\begin{aligned} m_{A\alpha,\sigma\#H} &= \begin{array}{c} A \quad H \quad A \quad H \\ \text{---} \quad \text{---} \quad \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \quad \text{---} \quad \text{---} \\ \alpha \quad \sigma \\ A \quad H \end{array} ; \\ m_{A\#\beta,\mu H} &= \begin{array}{c} A \quad H \quad A \quad H \\ \text{---} \quad \text{---} \quad \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \quad \text{---} \quad \text{---} \\ \beta \quad \mu \\ A \quad H \end{array} ; \\ \Delta_{A\phi,P\#H} &= \begin{array}{c} A \quad H \\ \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \\ \phi \quad P \\ A \quad H \end{array} ; \end{aligned}$$

$$\Delta_{A\#^{\psi,Q}H} = \text{Diagram} ;$$

$\eta_{A_{\alpha,\sigma}\#H} = \eta_A \otimes \eta_H = \eta_{A\#_{\beta,\mu}H}$ and $\epsilon_{A^{\phi,P}\#H} = \epsilon_A \otimes \epsilon_H = \epsilon_{A\#^{\psi,Q}H}$. $(A_{\alpha,\sigma}\#H, m_{A_{\alpha,\sigma}\#H}, \eta_{A_{\alpha,\sigma}\#H})$ and $(A\#_{\beta,\mu}H, m_{A\#_{\beta,\mu}H}, \eta_{A\#_{\beta,\mu}H})$ are called crossed products of A and H . $(A^{\phi,P}\#H, \Delta_{A^{\phi,P}\#H}, \epsilon_{A^{\phi,P}\#H})$ and $(A\#^{\psi,Q}H, \Delta_{A\#^{\psi,Q}H}, \epsilon_{A\#^{\psi,Q}H})$ are called crossed coproducts of A and H . If σ and μ or P and Q are trivial, then they are called the smash products or smash coproducts, written as $A_{\alpha}\#H$, $A\#_{\beta}H$, $A^{\phi}\#H$ and $A\#^{\psi}H$. Every smash product is called a semi-direct product in physics. In particular, the smash product $A_{\alpha}\#H$ is often written as $A\#H$. $(A_{\alpha,\sigma}\#^{\psi,Q}H, \Delta_{A\#^{\psi,Q}H}, \epsilon_{A\#^{\psi,Q}H}, m_{A_{\alpha,\sigma}\#H}, \eta_{A_{\alpha,\sigma}\#H})$ and $(A^{\phi,P}\#_{\beta,\mu}H, \Delta_{A^{\phi,P}\#H}, \epsilon_{A^{\phi,P}\#H}, m_{A\#_{\beta,\mu}H}, \eta_{A\#_{\beta,\mu}H})$ are called bicrossproducts. $(A_{\alpha}^{\phi}\#H, \Delta_{A^{\phi}\#H}, \epsilon_{A^{\phi}\#H}, m_{A_{\alpha}\#H}, \eta_{A_{\alpha}\#H})$ and $(A\#_{\beta}^{\psi}H, \Delta_{A\#^{\psi}H}, \epsilon_{A\#^{\psi}H}, m_{A\#_{\beta}H}, \eta_{A\#_{\beta}H})$ are called biproducts.

Note that S. Majid uses different notations.


4.1 Crossed Products

In this section, we give the necessary and sufficient conditions for crossed (co)products to become (co)algebras .


Theorem 4.1.1. *If A is an algebra and H is a bialgebra with $\alpha(\eta_H \otimes id_A) = id_A$ and $\alpha(id_H \otimes \eta_A) = \epsilon_H \eta_A$, then $D = A_{\alpha,\sigma}\#H$ is an algebra with unity element $\eta_D = \eta_A \otimes \eta_H$ iff (H, α) acts weakly on A and (A, α) is a twisted H -module with 2-cocycle σ .*

Proof. It is clear that $\eta_D = \eta_A \otimes \eta_H$ iff $\sigma(id_H \otimes \eta_H) = \sigma(\eta_H \otimes id_H) = \eta_A \epsilon_H$.


If D is an algebra, then the associative law holds, i.e.




By relation (1), we have that



and σ is a 2-cocycle from $H \otimes H$ to A .




and (A, α) is a twisted H -module.




and (H, α) acts weakly on A .


Conversely, if (H, α) acts weakly on A and (A, α) is a twisted H -module with 2-cocycle σ we show that the associative law holds. That is, we show that relation (1) holds.

the left hand of (1) $\stackrel{\text{by } (WA)}{=}$




$\stackrel{\text{by } (WA)}{=}$





by $(2-COC)$



Thus $A_{\alpha,\sigma} \# H$ is an algebra. $\square$

Dually, we have the followings:

Theorem 4.1.1' If H is an algebra and A is a bialgebra with $\beta(\eta_H \otimes id_A) = \epsilon_A \eta_H$ and $\beta(id_H \otimes \eta_A) = id_A$, then $A \#_{\beta, \mu} H$ is an algebra iff (A, β) acts weakly on H and (H, β) is a twisted A -module with 2-cocycle μ .

Theorem 4.1.1'' If A is a coalgebra and H is a bialgebra with $(\epsilon_H \otimes id_A)\phi = id_A$ and $(id_H \otimes \epsilon_A)\phi = \eta_H \epsilon_A$, then $A^{\phi, P} \# H$ is a coalgebra iff (H, ϕ) coacts weakly on A and (A, ϕ) is a twisted H -comodule with 2-cycle P .

Theorem 4.1.1''' If H is a coalgebra and A is a bialgebra with $(\epsilon_H \otimes id_A)\varphi = \eta_A \epsilon_H$ and $(id_H \otimes \epsilon_A) = id_H$, then $A \#^{\psi, Q} H$ is a coalgebra iff (A, ψ) coacts weakly on H and (H, ψ) is a twisted A -comodule with 2-cycle Q .

Corollary 4.1.2. (i) If A is an algebra and H is a bialgebra, then $A_\alpha \# H$ is an algebra iff (A, α) is an H -module algebra;

(ii) If H is an algebra and A is a bialgebra, then $A \#_\beta H$ is an algebra iff (H, β) is an A -module algebra;

(iii) If A is a coalgebra and H is a bialgebra, then $A^\phi \# H$ is a coalgebra iff (A, ϕ) is an H -comodule coalgebra;

(iv) If H is a coalgebra and A is a bialgebra, then $A \#^\psi H$ is a coalgebra iff (H, ψ) is an A -comodule coalgebra.

4.2 Bicrossproducts

In this section, we give the necessary and sufficient conditions for bicrossproducts to become bialgebras.

Lemma 4.2.1. Let A and H be two algebras in $\mathcal{C}$. Let $D = A \otimes H$ is an object of $\mathcal{C}$. Assume that (D, m_D, η_D) and $(D, \Delta_D, \epsilon_D)$ are an algebra and a coalgebra in $\mathcal{C}$ respectively such that

$$\begin{array}{c} A \otimes H \quad \eta_A \otimes H \\ \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \\ D \end{array} = \begin{array}{c} AH \quad H \\ | \quad \text{---} \text{---} \text{---} \\ A \quad H \end{array},$$

or

$$\begin{array}{c} A \otimes \eta_H \quad A \otimes H \\ \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \\ D \end{array} = \begin{array}{c} A \quad AH \\ \text{---} \text{---} \text{---} \\ A \quad H \end{array}.$$

If

$$\begin{array}{c} \eta_A \otimes H \quad \eta_A \otimes H \\ \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \\ D \quad D \end{array} = \begin{array}{c} \eta_A \otimes H \quad \eta_A \otimes H \\ \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \\ D \quad D \end{array},$$

$$\begin{array}{c} \eta_A \otimes H \quad A \otimes \eta_H \\ \text{---} \\ \text{---} \\ \text{---} \\ D \quad D \end{array} = \begin{array}{c} \eta_A \otimes H \quad A \otimes \eta_H \\ \text{---} \\ \text{---} \\ \text{---} \\ D \quad D \end{array},$$

$$\begin{array}{c} A \otimes \eta_H \quad \eta_A \otimes H \\ \text{---} \\ \text{---} \\ \text{---} \\ D \quad D \end{array} = \begin{array}{c} A \otimes \eta_H \quad \eta_A \otimes H \\ \text{---} \\ \text{---} \\ \text{---} \\ D \quad D \end{array},$$

$$\begin{array}{c} A \otimes \eta_H \quad A \otimes \eta_H \\ \text{---} \\ \text{---} \\ \text{---} \\ D \quad D \end{array} = \begin{array}{c} A \otimes \eta_H \quad A \otimes \eta_H \\ \text{---} \\ \text{---} \\ \text{---} \\ D \quad D \end{array},$$

then



$$\begin{array}{c} D \quad D \\ \text{---} \\ \text{---} \\ \text{---} \\ D \quad D \end{array} = \begin{array}{c} D \quad D \\ \text{---} \\ \text{---} \\ \text{---} \\ D \quad D \end{array}.$$



Proof. (i) Assume



$$\begin{array}{c} A \otimes H \quad \eta_A \otimes H \\ \text{---} \\ \text{---} \\ D \end{array} = \begin{array}{c} A \quad H \\ \text{---} \\ A \quad H \end{array}.$$

We first show that



$$\begin{array}{c} A \otimes H \quad \eta_A \otimes H \\ \text{---} \\ \text{---} \\ \text{---} \\ D \quad D \end{array} = \begin{array}{c} A \otimes H \quad \eta_A \otimes H \\ \text{---} \\ \text{---} \\ \text{---} \\ D \quad D \end{array} \dots\dots(1)$$



the left hand of (1) =  by assumption = 

 = 

=  =  = the right hand of (1).

Thus (1) holds. Next we show that

 = (2)

the left hand of (2) =  = 

$$\begin{aligned}
& \stackrel{\text{by (1)}}{=} \begin{array}{c} A \otimes \eta_H \quad A \otimes \eta_H \quad \eta_A \otimes H \\ \text{---} \quad \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \quad \text{---} \\ D \quad D \end{array} = \begin{array}{c} A \otimes \eta_H \quad A \otimes \eta_H \quad \eta_A \otimes H \\ \text{---} \quad \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \quad \text{---} \\ D \quad D \end{array} \\
& = \begin{array}{c} A \otimes \eta_H \quad A \otimes \eta_H \quad \eta_A \otimes H \\ \text{---} \quad \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \quad \text{---} \\ D \quad D \end{array} = \begin{array}{c} A \otimes \eta_H \quad A \otimes \eta_H \quad \eta_A \otimes H \\ \text{---} \quad \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \quad \text{---} \\ D \quad D \end{array} = \text{the right hand of (2)}.
\end{aligned}$$

Thus relation (2) holds.

Finally, we show that

$$\begin{array}{c} D \quad D \\ \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \\ D \quad D \end{array} = \begin{array}{c} D \quad D \\ \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \\ D \quad D \end{array} \dots\dots(3)$$

$$\begin{aligned}
& \text{the left hand of (3)} = \begin{array}{c} A \otimes \eta_H \quad \eta_A \otimes H \quad A \otimes \eta_H \quad \eta_A \otimes H \\ \text{---} \quad \text{---} \quad \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \quad \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \quad \text{---} \quad \text{---} \\ D \quad D \end{array} \\
& \stackrel{\text{by (1)}}{=} \begin{array}{c} A \otimes \eta_H \quad \eta_A \otimes H \quad A \otimes \eta_H \quad \eta_A \otimes H \\ \text{---} \quad \text{---} \quad \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \quad \text{---} \quad \text{---} \\ \text{---} \quad \text{---} \quad \text{---} \quad \text{---} \\ D \quad D \end{array}
\end{aligned}$$

$$\begin{array}{c}
\begin{array}{c}
A \otimes \eta_H \quad \eta_A \otimes H \quad A \otimes \eta_H \quad \eta_A \otimes H \\
\downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
\text{Diagram 1} \\
\downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
D \quad D
\end{array} \\
\stackrel{by (2)}{=} \\
\begin{array}{c}
A \otimes \eta_H \quad \eta_A \otimes H \quad A \otimes \eta_H \quad \eta_A \otimes H \\
\downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
\text{Diagram 2} \\
\downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
D \quad D
\end{array} \\
= \\
\begin{array}{c}
A \otimes \eta_H \quad \eta_A \otimes H \quad A \otimes \eta_H \quad \eta_A \otimes H \\
\downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
\text{Diagram 3} \\
\downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
D \quad D
\end{array} \\
= \\
\begin{array}{c}
A \otimes \eta_H \quad \eta_A \otimes H \quad A \otimes \eta_H \quad \eta_A \otimes H \\
\downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
\text{Diagram 4} \\
\downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
D \quad D
\end{array}
\end{array}$$

$$\begin{array}{c}
A \otimes H \quad A \otimes \eta_H \quad \eta_A \otimes H \\
\downarrow \quad \downarrow \quad \downarrow \\
\text{Diagram: A box with two inputs from the top and two outputs to the bottom, connected by a diagonal line.} \\
\downarrow \quad \downarrow \\
D \quad D
\end{array}
= \text{the right hand of (3)}.$$

This relation (3) holds.

(ii) We can show that relation (3) holds similarly

$$\begin{array}{c}
A \otimes \eta_H \quad A \otimes H \\
\downarrow \quad \downarrow \\
D
\end{array}
= \begin{array}{c}
A \quad AH \\
\downarrow \quad \downarrow \\
A \quad H
\end{array}. \quad \square$$

Theorem 4.2.2. *Let A and H be bialgebras with $(\epsilon_H \otimes id_A)\phi = id_A$ and $(id_H \otimes \epsilon_A)\phi = \eta_H \epsilon_A$ and $\beta(\eta_H \otimes id_A) = \epsilon_A \eta_H$ and $\beta(id_H \otimes \eta_A) = id_A$ and $\epsilon_H \otimes \epsilon_A = \epsilon_A \beta$ and $\phi \eta_A = \eta_H \otimes \eta_A$. Then $D = A^{\phi, P} \#_{\beta, \mu} H$ is a bialgebra iff D is an algebra and a coalgebra, and relations (BB1)–(BB5) hold.*

Proof. (See the proof of [79, Theorem 2.9]) Let $D = A^{\phi, P} \#_{\beta, \mu} H$ be a bialgebra. We have that

$$\begin{array}{c}
A \otimes H \quad A \otimes H \\
\downarrow \quad \downarrow \\
D \quad D
\end{array}
= \begin{array}{c}
A \otimes H \quad A \otimes H \\
\downarrow \quad \downarrow \\
\text{Diagram: A box with two inputs from the top and two outputs to the bottom, connected by a diagonal line.} \\
\downarrow \quad \downarrow \\
D \quad D
\end{array} \quad \dots \quad (1)$$

By (1), we have that

$$\begin{array}{c}
A \otimes \eta_H \quad A \otimes \eta_H \\
\downarrow \quad \downarrow \\
\epsilon \otimes \epsilon \quad \epsilon \otimes \epsilon
\end{array}
= \begin{array}{c}
A \otimes \eta_H \quad A \otimes \eta_H \\
\downarrow \quad \downarrow \\
\text{Diagram: A box with two inputs from the top and two outputs to the bottom, connected by a diagonal line.} \\
\downarrow \quad \downarrow \\
\epsilon \otimes \epsilon \quad \epsilon \otimes \epsilon
\end{array}, \quad \begin{array}{c}
A \quad A \\
\downarrow \quad \downarrow \\
\mu \\
\downarrow \\
\epsilon
\end{array}
= \begin{array}{c}
A \quad A \\
\downarrow \quad \downarrow \\
\epsilon \quad \epsilon
\end{array}.$$

$$\begin{array}{c}
\eta_A \otimes \eta_H \eta_A \otimes \eta_H \\
\downarrow \quad \downarrow \\
\epsilon \otimes id_H \quad \epsilon \otimes id_H
\end{array}
= \begin{array}{c}
\eta_A \otimes \eta_H \quad \eta_A \otimes \eta_H \\
\downarrow \quad \downarrow \\
\text{Diagram: A box with two inputs from the top and two outputs to the bottom, connected by a diagonal line.} \\
\downarrow \quad \downarrow \\
\epsilon \otimes id_H \quad \epsilon \otimes id_H
\end{array} \quad \text{and} \quad \begin{array}{c}
\eta_A \\
\downarrow \\
P \\
\downarrow \quad \downarrow \\
H \quad H
\end{array}
= \begin{array}{c}
\eta_H \quad \eta_H \\
\downarrow \quad \downarrow \\
H \quad H
\end{array}.$$


Similarly, we can obtain (BB1)–(BB5) by relation (1).


Conversely, if D is an algebra and coalgebra, and relations (BB1)–(BB5) hold, we show that the conditions in Lemma 4.2.1 hold. We first show that


$$\begin{array}{c} A \otimes \eta_H \quad A \otimes \eta_H \\ \text{---} \text{---} \\ | \\ \text{---} \text{---} \\ | \quad | \\ D \quad D \end{array} = \begin{array}{c} A \otimes \eta_H \quad A \otimes \eta_H \\ \text{---} \text{---} \\ \text{---} \text{---} \\ | \quad | \\ D \quad D \end{array} .$$


the left hand


by (BB4), (WA), (WCA)











$$by \stackrel{(WCA)}{=} \text{the right hand.}$$

Similarly, we have that

$$\begin{array}{c}
 \eta_A \otimes H \quad \eta_A \otimes H \\
 \text{---} \text{---} \\
 | \\
 D \quad D
 \end{array}
 =
 \begin{array}{c}
 \eta_A \otimes H \quad \eta_A \otimes H \\
 | \quad | \\
 \text{---} \text{---} \\
 | \quad | \\
 D \quad D
 \end{array},$$

$$\begin{array}{c}
 \eta_A \otimes H \quad A \otimes \eta_H \\
 \text{---} \text{---} \\
 | \\
 D \quad D
 \end{array}
 =
 \begin{array}{c}
 \eta_A \otimes H \quad A \otimes \eta_H \\
 | \quad | \\
 \text{---} \text{---} \\
 | \quad | \\
 D \quad D
 \end{array},$$

$$\begin{array}{c}
A \otimes \eta_H \quad \eta_A \otimes H \\
\text{---} \text{---} \text{---} \text{---} \\
| \\
\text{---} \text{---} \text{---} \text{---} \\
D \quad D
\end{array}
=
\begin{array}{c}
A \otimes \eta_H \quad \eta_A \otimes H \\
\text{---} \text{---} \text{---} \text{---} \\
\text{---} \text{---} \text{---} \text{---} \\
\text{---} \text{---} \text{---} \text{---} \\
D \quad D
\end{array}
.$$

Obviously,

$$\begin{array}{c} A \otimes \eta_H \quad A \otimes H \\ \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \\ D \end{array} = \begin{array}{c} A \quad AH \\ \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \\ A \quad H \end{array}.$$

By Lemma 4.2.1, D is a bialgebra. $\square$

Dually we have Theorem 4.2.2' Let A and H be bialgebras with $\alpha(id_H \otimes \eta_A) = \epsilon_H \epsilon_A$ and $\alpha(\eta_H \otimes id_A) = id_A$ and $(\epsilon_H \otimes id_A)\varphi = \epsilon_H \eta_A$ and $(id_H \otimes \epsilon_A)\varphi = id_H$ and $\epsilon_H \otimes \epsilon_A = \epsilon_A \alpha$ and $\varphi \eta_H = \eta_H \otimes \eta_A$. Then $D = A_{\alpha, \sigma} \#^{\psi, Q} H$ is a bialgebra iff D is an algebra and coalgebra, and relations (BB1')–(BB5') hold.

Corollary 4.2.3. *Let A and H be bialgebras. Then*

(i) $A_\alpha \#^\psi H$ is a bialgebra iff (A, α) is an H -module algebra, (H, ψ) is an A -comodule coalgebra, and $(BB3')$, $(BB10')$, $(BB11')$ hold;

(ii) $A^\phi \#_\beta H$ is a bialgebra iff (A, ϕ) is an H -comodule coalgebra, (H, β) is an A -module algebra, and $(BB3)$, $(BB10)$, $(BB11)$ hold.

Corollary 4.2.4. *Let A and H be bialgebras. Then*

(i) $A_{\alpha,\sigma}\#H$ is a bialgebra iff $A_{\alpha,\sigma}\#H$ is an algebra, σ and α are coalgebra morphisms, H is cocommutative with respect to (A, α) , and $(BB8')$ holds;

(ii) $A \#_{\beta, \mu} H$ is a bialgebra iff $A \#_{\beta, \mu} H$ is an algebra, μ and β are coalgebra morphisms, A is cocommutative with respect to (H, β) , and (BB9') holds;

(iii) $A^{\phi, P} \# H$ is a bialgebra iff $A^{\phi, P} \# H$ is an algebra, P and ϕ are coalgebra morphisms, H is commutative with respect to (A, ϕ) , and (BB8) holds;



(iv) $A \#^{\psi, Q} H$ is a bialgebra iff $A \#^{\psi, Q} H$ is an algebra, Q and ψ are coalgebra morphisms, H is commutative with respect to (A, ψ) , and (BB9) holds;

4.3 Biproducts

In this section, we give the necessary and sufficient conditions for biproducts to become bialgebras.

Lemma 4.3.1. *Let H be a bialgebra, (A, α) an H -module algebra and (A, ϕ) is an H -comodule*

coalgebra. Set $D = A_\sigma^\phi \# H$. Then (i) Relation (1) and relation (2) are equivalent.



(ii) Relation (3) is equivalent to (BB7') and relation (4)




Diagram (4) shows an equality between two expressions. On the left, a multiplication node ϕ (a circle with a dot) has two inputs: H (a vertical line) and A (a vertical line). Its output is a vertical line that goes into a comultiplication node ϕ (a circle with a dot). The comultiplication node has two outputs: A and A . On the right, there are two vertical lines, each with a comultiplication node ϕ in the middle. The left vertical line has input H and output A . The right vertical line has input A and output A . The two comultiplication nodes are connected by a crossing line.

.....(4)

(iii) Relation (2) implies relation (5) (6).

Diagram (5) shows an equality between two expressions. On the left, a multiplication node ϕ has inputs AH and A . Its output is a vertical line that goes into a comultiplication node ϕ . The comultiplication node has two outputs: A and A . On the right, there are two vertical lines, each with a comultiplication node ϕ in the middle. The left vertical line has input A and output A . The right vertical line has input A and output A . The two comultiplication nodes are connected by a crossing line.

.....(5)

Diagram (6) shows an equality between two expressions. On the left, a comultiplication node α has inputs AH and A . Its output is a vertical line that goes into a multiplication node ϕ . The multiplication node has two outputs: A and A . On the right, there are two vertical lines, each with a multiplication node ϕ in the middle. The left vertical line has input A and output A . The right vertical line has input A and output A . The two multiplication nodes are connected by a crossing line.

.....(6)

(iv) Relation (6) is equivalent to (BB6') and relation (7)

Diagram (7) shows an equality between two expressions. On the left, a comultiplication node α has inputs H and A . Its output is a vertical line that goes into a multiplication node ϕ . The multiplication node has two outputs: A and A . On the right, there are two vertical lines, each with a multiplication node ϕ in the middle. The left vertical line has input H and output A . The right vertical line has input A and output A . The two multiplication nodes are connected by a crossing line.

.....(7)




(v) Relation (2) implies (3) when $\epsilon_A(\alpha) = \epsilon_A \otimes \epsilon_H$; (vi) ϵ_D is an algebra morphism iff ϵ_A is an algebra morphism and

Diagram shows an equality between two expressions. On the left, a comultiplication node α has inputs H and A . Its output is a vertical line that goes into a multiplication node ϵ . The multiplication node has two outputs: A and A . On the right, there are two vertical lines, each with a multiplication node ϵ in the middle. The left vertical line has input H and output A . The right vertical line has input A and output A . The two multiplication nodes are connected by a crossing line.

(vii)

$$\begin{array}{c} \eta_D \\ \downarrow \\ D \end{array} = \begin{array}{c} \eta_A \quad \eta_H \quad \eta_A \quad \eta_H \\ \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\ A \quad H \quad A \quad H \end{array} \quad \text{iff} \quad \begin{array}{c} \eta_A \\ \downarrow \\ \phi \\ \downarrow \\ H \quad A \end{array} = \begin{array}{c} \eta_H \quad \eta_A \\ \downarrow \quad \downarrow \\ H \quad A \end{array} \quad \text{and} \quad \begin{array}{c} \eta_A \\ \downarrow \\ \downarrow \\ A \quad A \end{array} = \begin{array}{c} \eta_A \quad \eta_A \\ \downarrow \quad \downarrow \\ A \quad A \end{array}.$$




Proof. (i) It is clear that (1) implies (2). Conversely, if relation (2) holds, we have that

the left side of (1) =  =  $\stackrel{\text{by (2)}}{=}$ 

= the right side of (1).

(ii) Assume relation (3) holds. Applying $(id_A \otimes \eta_H \otimes id_A)$ on relation (3), we get relation (4). Applying $(\eta_A \otimes id_H \otimes id_A)$ on relation (3), we get (BB7').

Conversely, assume relation (4) and (BB7') hold. We see that

the left side of (3) $\stackrel{\text{by (4)}}{=}$  =  $\stackrel{(BB7')}{=}$ 

= the right side of (3).

(iii) Applying $id_A \otimes \eta_H \otimes id_A$ on relation (2), we can get relation (5). Applying $id_A \otimes \epsilon_H \otimes id_A$ on relation (2), we can get relation (6).

(iv) Assume relation (6) holds. Applying $\eta_A \otimes id_H \otimes id_A$ on relation (6), we can get relation (7). Applying $id_A \otimes \eta_H \otimes id_A$ on relation (6), we can get relation (BB6'). Conversely, assume (BB6') and relation (7) hold. We see that

the left side of (6) $\stackrel{\text{by (BB6')}}{=} \stackrel{\text{by (7)}}{=} \text{the right side of (6)}.$

(v),(vi) and (vii) are clear.

Theorem 4.3.2. *Let H be a bialgebra, (A, α) an H -module algebra and (A, ϕ) an H -comodule coalgebra. Then the following conditions are equivalent.*

- (i) $A_\alpha^\phi \# H$ is a bialgebra.
- (ii) (A, α) is an H -module coalgebra, (A, ϕ) is an H -comodule algebra, $\Delta_A(\eta_A) = \eta_A \otimes \eta_A$, ϵ_A is an algebra morphism, and (BB6')–(BB7') hold.
- (iii) ϵ_A and ϕ are algebra morphisms, α is a coalgebra morphism, $\Delta_A(\eta_A) = \eta_A \otimes \eta_A$, and (BB6')–(BB7') hold.


Proof. (ii) $\Leftrightarrow$ (iii) is clear. (i) $\Rightarrow$ (ii) follows from Lemma 4.3.1, so it remains to show that (ii) $\Rightarrow$ (i). By Lemma 4.3.1, we only need show that relation (2) holds. We see that


the left side of (2) $\stackrel{\text{by (6)}}{=} \stackrel{\text{by (3)}}{=} \text{the right side of (2)}. \square$

$\square$

Dually we have

Theorem 4.3.4. *Let H be a Hopf algebra, (A, α) an H -module algebra and (A, ϕ) an H -comodule coalgebra with an antipode S_A . Then $D = A_\alpha^\phi \# H$ has an antipode*

$S_D * id_D =$


since (A, ϕ) is an H -comodule coalgebra
 $\quad \quad \quad =$


since (A, α) is an H -module algebra

and

Thus S_D is an antipode of D . $\square$

Lemma 4.3.5. *Let $\mathcal{C}$ be a symmetric braided tensor category and H a bialgebra in $\mathcal{C}$.*

(i) *If (M, α, ϕ) is a left H -module and left H -comodule in $\mathcal{C}$, then (YD) and $(BB7')$ are equivalent.*

(ii) Let (H, R) be a quasitriangular Hopf algebra in $\mathcal{C}$ and (A, α) a Hopf algebra in $({}_H\mathcal{M}(\mathcal{C}), C^R)$. Then (A, ϕ, α) is a Yetter-Drinfeld H -module in ${}^H_H\mathcal{YD}(\mathcal{C})$, where $\phi = (id_H \otimes \alpha)(C_{H,H} \otimes id_A)(R \otimes id_A)$.

(iii) Let (H, r) be a coquasitriangular Hopf algebra in $\mathcal{C}$ and (A, ϕ) a Hopf algebra in $({}_H\mathcal{M}(\mathcal{C}), C^r)$. Then (A, ϕ, α) is a Yetter-Drinfeld H -module in ${}^H_H\mathcal{YD}(\mathcal{C})$, where $\alpha = (r \otimes id_A)(C_{H,H} \otimes id_A)(id_H \otimes \phi)$.

Consequently, by Lemma 4.3.5, Theorem 4.3.2 and Theorem 4.3.4 we have following three corollaries

Corollary 4.3.6. *Let H be a braided Hopf algebra in symmetric braided tensor category $\mathcal{C}$ and (A, α, ϕ) be a bialgebra or Hopf algebra in $({}^H_H\mathcal{YD}(\mathcal{C}), {}^{YD}C)$. Then $A_\alpha^\phi \# H$ is a bialgebra or Hopf algebra in $\mathcal{C}$.*

We can easily get the well-known bosonisation theorem in [83, Theorem 9.4.12].

Corollary 4.3.7. *Let (H, R) be a quasitriangular Hopf algebra in symmetric $\mathcal{C}$ and (A, α) be a Hopf algebra in $({}_H\mathcal{M}(\mathcal{C}), C^R)$. Then $A_\alpha^\phi \# H$ is a Hopf algebra in $\mathcal{C}$, where $\phi = (id_H \otimes \alpha)(C_{H,H} \otimes id_A)(R \otimes id_A)$.*

Corollary 4.3.8. *Let (H, r) be a coquasitriangular Hopf algebra in symmetric $\mathcal{C}$ and (A, ϕ) a Hopf algebra in $({}_H\mathcal{M}(\mathcal{C}), C^r)$. Then $A_\alpha^\phi \# H$ is a Hopf algebra in $\mathcal{C}$, where $\alpha = (r \otimes id_A)(C_{H,H} \otimes id_A)(id_H \otimes \phi)$.*

In particular, these corollaries hold for ordinary (co)quasitriangular Hopf algebras.

We now get the Lagrange's theorem for braided Hopf algebras.

Theorem 4.3.9. *Let H be a finite-dimensional ordinary Hopf algebra, If (A, α) is a finite-dimensional Hopf algebra in ${}^H_H\mathcal{YD}$ and B is a subHopf algebra of A in ${}^H_H\mathcal{YD}$, then*

$$\dim B \mid \dim A.$$

That is, the dimension of B divides the dimension of A .

Proof. By Corollary 4.3.6 and [87, Corollary 3.2.1],

$$\dim (B_\alpha^\phi \# H) \mid \dim (A_\alpha^\phi \# H).$$

Consequently,

$$\dim B \mid \dim A \quad . \quad \square$$

Definition 4.3.10. *A coalgebra A is called super cocommutative or quantum cocommutative if $\Delta = C_{A,A}\Delta$; an algebra A is called super commutative or quantum commutative if $m_A = m_A C_{A,A}$.*

Remark : In the category $\mathcal{Vect}(k)$, a coalgebra A is called cocommutative if $\Delta = C_{A,A}\Delta$; an algebra A is called commutative if $m_A = m_A C_{A,A}$. Here braiding C is usual twist map.

Proposition 4.3.11. (i) If H is a quantum commutative, or quantum cocommutative Hopf algebra, then $S^2 = id_H$.

(ii) If H is a Hopf algebra, then the following conditions are equivalent:

$$(1) mC_{H,H}^{-1}(id_H \otimes S)\Delta_H = \eta_H \epsilon_H;$$

$$(2) mC_{H,H}^{-1}(S \otimes id_H)\Delta_H = \eta_H \epsilon_H;$$

$$(3) S^2 = id_H;$$

Proof. (ii) We first show that (2) implies (3). If (2) holds, we see that

$$\begin{aligned} m(id_H \otimes S)(S \otimes S)\Delta_H &= mC_{H,H}^{-1}(S \otimes id_H)\Delta_H S \quad (\text{by Pro. 1.0.13}) \\ &= \eta_H \epsilon_H \quad (\text{by (2)}) \end{aligned}$$

and

$$\begin{aligned} m(S \otimes id_H)(S \otimes S)\Delta_H &= SmC_{H,H}^{-1}(S \otimes id_H)\Delta_H \quad (\text{by Pro. 1.0.13}) \\ &= \eta_H \epsilon_H \quad (\text{by (2)}). \end{aligned}$$

Thus S^2 is the convolution inverse of S , which implies $S^2 = id_H$. Therefore (2) implies (3). Similarly, (1) implies (3).

We next show that (3) implies (1). If (3) holds, we see that

$$\begin{aligned} \eta_H \epsilon &= m(id_H \otimes S)\Delta_H \\ &= m(S \otimes id_H)(S \otimes S)\Delta_H \\ &= m(S \otimes id_H)C_{H,H}^{-1}\Delta_H S \quad (\text{by Pro. 1.0.13}) \end{aligned}$$

and

$$\begin{aligned} \eta_H \epsilon &= m(id_H \otimes S)\Delta_H \\ &= m(S \otimes id_H)(S \otimes S)\Delta_H \\ &= m(S \otimes id_H)C_{H,H}^{-1}\Delta_H S \quad (\text{by Pro. 1.0.13}) \\ &= \eta_H \epsilon S^{-1} \\ &= mC_{H,H}^{-1}(id_H \otimes S)\Delta_H. \end{aligned}$$

Thus (1) holds. Similarly (3) implies (2).

(i) It follows from (ii) $\square$

In this subsection, we construct a braided Hopf algebra in a Yetter-Drinfeld category by means of a graded Hopf algebra.

Lemma 4.3.12. (see [91, p.1530],[4] and [99]) Assume that H and Λ are two ordinary Hopf algebras. If there exist two bialgebra homomorphisms $\pi_0 : \Lambda \rightarrow H$ and $\iota_0 : H \rightarrow \Lambda$ with $\pi_0 \iota_0 = id_H$, then

- (i) $(\Lambda, \delta^+, \alpha^+)$ is a right H -Hopf module with $\delta^+ =: (id \otimes \pi_0)\Delta$ and $\alpha^+ =: m(id \otimes \iota_0)$
- (ii) $A \# H \stackrel{\alpha^+}{\cong} \Lambda$ (as algebras) with $A =: \Lambda^{coH}$.
- (iii) $\Lambda^{coH} = Im \omega$ with $\omega =: id_\Lambda * (\iota_0 \circ S_H \circ \pi_H) : \Lambda \rightarrow \Lambda$.

Proof. (i) It is easy to check directly that $(\Lambda, \delta^+, \alpha^+)$ is a right H -Hopf module.

(ii) By the fundamental theorem of Hopf modules, we have $\Lambda^{coH} \otimes H \cong \Lambda$ as H -Hopf modules with isomorphism α^+ . We can also check that Λ^{coH} is an H -module algebra with adjoint action ad . Moreover, α^+ preserves the algebra operations of $\Lambda^{coH} \# H$.

(iii) Obviously, for any $x \in A$, $\omega(x) = x$, which implies $A \subseteq Im(\omega)$. On the other hand, for any $x \in \Lambda$, see that

$$\begin{aligned}
\delta^+ \circ \omega(x) &= \sum (id \otimes \pi_0)\Delta(x_1(\iota_0 \circ S \circ \phi_0(x_2))) \\
&= \sum x_1 \iota_0 \circ S \circ \phi_0(x_4) \otimes \phi_0(x_2) S \circ \phi_0(x_3) \\
&= \omega(x) \otimes 1.
\end{aligned}$$

Thus $A = Im(\omega)$. $\square$

Theorem 4.3.13. (see [91, p.1530],[4] and [99]) Assume that H and Λ are two ordinary Hopf algebras. If there exist two bialgebra homomorphisms $\pi_0 : \Lambda \rightarrow H$ and $\iota_0 : H \rightarrow \Lambda$ with $\pi_0 \iota_0 = id_H$, then (A, α, ϕ) is a braided Hopf algebra in ${}^H_H \mathcal{YD}$ and $A \# H \stackrel{\alpha^+}{\cong} \Lambda$ (as Hopf algebras), where $\alpha(h \otimes a) = \sum \iota_0(h_1) a \iota_0(S(h_2))$, $\phi(a) = (\pi_0 \otimes id)\Delta_\Lambda(a)$, for any $h \in H, a \in A$.

Proof. Define $\Delta_A = (\omega \otimes id)\Delta$ and $\epsilon_A = \epsilon_\Lambda$. By the proof of [99, Theorem 3], $(A, m, \eta, \Delta_A, \epsilon)$ become a braided Hopf algebra in ${}^H_H \mathcal{YD}$ under module operation α and comodule operation ϕ .

Set $\Phi = \alpha^+$. For any $x \in A, h \in H$, see that

$$\begin{aligned}
\Delta \circ \Phi(x \# h) &= \sum x_1 \iota_0(h_1) \otimes x_2 \iota_0(h_2) \\
&= \sum \alpha^+(x_1, h_1) \otimes \alpha^+(x_2, h_2) \\
&= (\Phi \otimes \Phi)\Delta(x \# h).
\end{aligned}$$

Thus Φ is a coalgebra map. By Lemma 4.3.12 (ii), we have Φ is a Hopf algebra isomorphism. $\square$

Assume that $\Lambda = \oplus_{i=0}^\infty \Lambda_i$ is a graded Hopf algebra with $H = \Lambda_0$. Let $\pi_0 : \Lambda \rightarrow \Lambda_0$ and $\iota_0 : \Lambda_0 \rightarrow \Lambda$ denote the canonical projection and injection. It is clear that π_0 and ι_0 are two bialgebra homomorphisms. Consequently, Theorem above holds. Λ^{coH} is called the diagram of Λ , written $diag(\Lambda)$.

Chapter 5

The Factorization Theorem of Bialgebras in Braided Tensor Categories

It is well-known that the factorization of domain plays an important role in ring theory. S. Majid in [83, Theorem 7.2.3] studied the factorization of Hopf algebra and showed that $H \cong A \bowtie H$ for two subbialgebras A and B when multiplication m_H is bijective.

S. Majid found a method to turn an ordinary Hopf algebra H into a braided Hopf algebra $\underline{H}$. This method is called the transmutation, and $\underline{H}$ is called the braided group analogue of H (see [82] and [83]). Huixiang Chen in [26] showed that the double cross coproduct $A \bowtie^R H$ of two quasitriangular Hopf algebras is a quasitriangular Hopf algebra. One needs to know the relation among the braided group analogues of the double cross coproduct $A \bowtie^R H$, A and H .

In this chapter we generalize Majid's factorization theorem into braided tensor categories. We show that the braided group analogue of double cross (co)product is double cross (co)product of braided group analogues. We give the factorization theorem of Hopf algebras and the relation between Hopf algebras and their factors in braided tensor categories.

5.1 the braided reconstruction theorem

In this section, we introduce the method (transmutation) turning (co)quasi- triangular Hopf algebra into a braided Hopf algebra, which is due to S. Majid [82].

Let $\mathcal{C}$ be a tensor category, $\mathcal{D}$ a braided tensor category and (F, μ_0, μ) a tensor functor from $\mathcal{C}$ to $\mathcal{D}$ with $\mu_0 = id_I$. Let $Nat(G, T)$ denote all the natural transformations from functor G to functor T . Assume that there is an object B of $\mathcal{D}$ and a natural transformation α in

$\text{Nat}(B \otimes F, F)$. Here $(B \otimes F)(X) = B \otimes F(X)$ for any object X in $\mathcal{D}$. Let

$$\begin{aligned}
 \theta_V(g)_X &= \begin{array}{c} V \quad F(X) \\ | \quad / \\ \textcircled{g} \quad \quad \\ | \quad \quad \\ \textcircled{\alpha_X} \\ | \\ F(X) \end{array}, & \theta_V^{(2)}(h)_{X_1 \otimes X_2} &= \begin{array}{c} V \quad F(X_1) \quad F(X_2) \\ | \quad / \quad / \\ \textcircled{h} \quad \quad \quad \\ | \quad \quad \quad \\ \textcircled{\alpha_{X_1}} \quad \textcircled{\alpha_{X_2}} \\ | \quad \quad | \\ F(X_1) \quad F(X_2) \end{array}, \\
 \theta_V^{(3)}(p)_{X_1 \otimes X_2 \otimes X_3} &= \begin{array}{c} V \quad F(X_1) \quad F(X_2) \quad F(X_3) \\ | \quad / \quad / \quad / \\ \textcircled{p} \quad \quad \quad \\ | \quad \quad \quad \\ \textcircled{\alpha_{X_1}} \quad \textcircled{\alpha_{X_2}} \quad \textcircled{\alpha_{X_2}} \\ | \quad \quad | \quad | \\ F(X_1) \quad F(X_2) \quad F(X_3) \end{array}
 \end{aligned}$$

for any object $X, X_1, X_2, X_3 \in \text{ob}\mathcal{C}$, $g \in \text{Hom}_{\mathcal{D}}(V, B)$, $h \in \text{Hom}_{\mathcal{D}}(V, B \otimes B)$, $p \in \text{Hom}_{\mathcal{D}}(V, B \otimes B \otimes B)$.

Let $V \in \text{ob}\mathcal{D}$, $g, g' \in \text{Hom}_{\mathcal{D}}(V, B)$. θ_V is called injective if $\theta_V(g)_X = \theta_V(g')_X$ for any $X \in \text{ob}\mathcal{D}$ implies $g = g'$. Similarly, we can obtain the definitions about $\theta_V^{(2)}$ and $\theta_V^{(3)}$.

Proposition 5.1.1. (see [82, Proposition 3.8] [83, Theorem 9.4.6, Proposition 9.4.7] If θ_V , $\theta_V^{(2)}$ and $\theta_V^{(3)}$ are injective for any object $V = I, B, B \otimes B, B \otimes B \otimes B$, then

- (i) B is a bialgebra living in $\mathcal{D}$, called the braided bialgebra determined by braided reconstruction;
- (ii) Furthermore, if $\mathcal{C}$ is a rigid tensor category (i.e. every object has a left duality), then B is Hopf algebra, called the braided Hopf algebra determined by braided reconstruction.
- (iii) Furthermore, if $\mathcal{D}$ and $\mathcal{C}$ is braided tensor categories, then B is quasitriangular.

Proof. (i) We define the multiplication, unit, comultiplication and counit of B as follows:

$$\begin{aligned}
 \theta_{B \otimes B}(m_B)_X &= \begin{array}{c} B \quad B \quad F(X) \\ | \quad / \quad / \\ \textcircled{\alpha_X} \quad \quad \\ | \quad \quad \\ \textcircled{\alpha_X} \\ | \\ F(X) \end{array}, & \theta_I(\eta_B) &= \begin{array}{c} F(X) \\ | \\ F(X) \end{array},
 \end{aligned}$$

$$\theta_B^{(2)}(\Delta_B)_{X \otimes Y} = \begin{array}{c} \begin{array}{ccc} B & F(X) & F(Y) \\ | & \diagdown & / \\ & \mu & \\ & | & \\ \alpha_{X \otimes Y} & & \end{array} \\ | \\ \mu^{-1} \\ \begin{array}{ccc} & F(X) & F(Y) \end{array} \end{array}, \quad \epsilon_B = \begin{array}{c} \begin{array}{cc} B & F(I) \\ \diagdown & / \\ \alpha_I \end{array} \end{array}$$

for any object $X, Y, Z \in \text{ob}\mathcal{C}$.

We only show that the coassociative law holds, the others can be shown similarly. That is, we need show that

$$\begin{array}{c} B \\ | \\ \text{---} \\ \diagup \quad \diagdown \\ B \quad B \end{array} = \begin{array}{c} B \\ | \\ \text{---} \\ \diagdown \quad \diagup \\ B \quad B \end{array} \quad \text{.....(1)}$$

We see that

$$\theta_B^{(3)}(\text{---})_{X \otimes Y \otimes Z} = \begin{array}{c} \begin{array}{cccc} & B & F(X) & F(Y) & F(Z) \\ & | & \diagdown & / & \\ & \text{---} & & & \\ & \diagup & \diagdown & & \\ \alpha_X & & \alpha_Y & & \alpha_Z \\ | & & | & & | \\ F(X) & & F(Y) & & F(Z) \end{array} \end{array}$$

$$\begin{array}{c}
 \begin{array}{c}
 \begin{array}{c}
 F(X) \quad F(Y) \quad F(Z) \\
 \downarrow \quad \downarrow \quad \downarrow \\
 \text{---} \quad \text{---} \quad \text{---} \\
 \downarrow \quad \downarrow \quad \downarrow \\
 \alpha_{X \otimes Y} \quad \mu^{-1} \quad \alpha_Z \\
 \downarrow \quad \downarrow \quad \downarrow \\
 F(X) \quad F(Y) \quad F(Z)
 \end{array} \\
 = \\
 \begin{array}{c}
 B \quad F(X) \quad F(Y) \quad F(Z) \\
 \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
 \mu \quad \mu \\
 \downarrow \quad \downarrow \\
 \alpha_{(X \otimes Y) \otimes Z} \\
 \downarrow \\
 \mu^{-1} \\
 \downarrow \quad \downarrow \quad \downarrow \\
 \mu^{-1} \quad F(X) \quad F(Y) \quad F(Z)
 \end{array} \\
 = \\
 \begin{array}{c}
 B \quad F(X) \quad F(Y) \quad F(Z) \\
 \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
 \mu \quad \mu \\
 \downarrow \quad \downarrow \\
 \alpha_{(X \otimes Y) \otimes Z} \\
 \downarrow \\
 \mu^{-1} \\
 \downarrow \quad \downarrow \quad \downarrow \\
 F(X) \quad F(Y) \quad F(Z)
 \end{array}
 \end{array}
 \end{array}$$

$$\begin{array}{c}
 \text{since } (F, \mu) \text{ is tensor functor} \\
 = \\
 \begin{array}{c}
 B \quad F(X) \quad F(Y) \quad F(Z) \\
 \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
 \mu \quad \mu \\
 \downarrow \quad \downarrow \\
 \alpha_{(X \otimes Y) \otimes Z} \\
 \downarrow \\
 \mu^{-1} \\
 \downarrow \quad \downarrow \quad \downarrow \\
 F(X) \quad F(Y) \quad F(Z)
 \end{array}
 \end{array}$$

$$\begin{array}{c}
 B \\
 \downarrow \\
 \theta_B^{(3)} \left(\begin{array}{c} \text{---} \\ \downarrow \quad \downarrow \quad \downarrow \\ B \quad B \quad B \end{array} \right)_{X \otimes Y \otimes Z}
 \end{array}$$

Thus relation (1) holds since $\theta_B^{(3)}$ is injective.

(ii) We define

$$\theta_B(S_B)_X = \begin{array}{c}
 \begin{array}{c}
 B \quad F(X) \\
 \downarrow \quad \downarrow \\
 b_{F(X)} \\
 \downarrow \quad \downarrow \\
 \alpha_{X^*} \\
 \downarrow \quad \downarrow \\
 d_{F(X)} \\
 \downarrow \quad \downarrow \\
 F(X) \quad F(X)
 \end{array}
 \end{array}$$

for any $X \in \text{ob} \mathcal{C}$. We can show that S_B is the antipode of B . In fact

$$\begin{aligned}
\theta_B(S_B * I)_X &= \text{Diagram 1} = \text{Diagram 2} = \text{Diagram 3} \\
&= \text{Diagram 4} = \text{Diagram 5} = \text{Diagram 6}
\end{aligned}$$

The diagrams are string diagrams representing morphisms in a monoidal category. They involve objects B and $F(X)$, and nodes S (multiplication), ϵ (counit), and α_X (associator). The sequence of diagrams shows the simplification of the expression $\theta_B(S_B * I)_X$ through various string diagram identities, ultimately resulting in the product of the counit ϵ on B and the identity on $F(X)$.

Similarly, we get that

$$\theta_B(I * S_B)_X = \text{Diagram}$$

The diagram shows the counit ϵ on B and the identity on $F(X)$.

Obviously,

$$\theta_B(I * S_B)_X = \text{Diagram}$$

The diagram shows the counit ϵ on B , followed by the unit η on $F(X)$, and then the associator α_X .

We showed that S_B is the antipode of B .

(iii) We define

$$\theta_I^{(2)}(R)_{X \otimes Y} = \begin{array}{c} F(X) \quad F(Y) \\ \diagdown \quad \diagup \\ \mu \\ \hline F(C_{X,Y}) \\ \hline \mu^{-1} \\ \diagup \quad \diagdown \\ F(X) \quad F(Y) \end{array} \quad \text{and} \quad \theta_B^{(2)}(\bar{\Delta})_{X \otimes Y} = \begin{array}{c} B \quad F(X) \quad F(Y) \\ \diagdown \quad \diagup \quad \diagdown \\ \alpha \quad \alpha \\ \diagup \quad \diagdown \\ F(X) \quad F(Y) \end{array} .$$

We easily check that $(B, \bar{\Delta}, \epsilon)$ is a coalgebra by modifying the proof of part (i).

We see that

$$\begin{aligned} \theta_I^{(3)}(R)_{X \otimes Y \otimes Z} &= \begin{array}{c} F(X) \quad F(Y) \quad F(Y) \\ \diagdown \quad \diagup \quad \diagdown \\ R \quad R \\ \diagup \quad \diagdown \quad \diagup \\ \alpha \quad \alpha \quad \alpha \\ \diagup \quad \diagdown \quad \diagup \\ F(X) \quad F(Y) \quad F(Z) \end{array} \\ &= \begin{array}{c} F(X) \quad F(Y) \quad F(Y) \\ \diagdown \quad \diagup \quad \diagdown \\ R \quad R \\ \diagup \quad \diagdown \quad \diagup \\ \alpha \quad \alpha \quad \alpha \\ \diagup \quad \diagdown \quad \diagup \\ F(X) \quad F(Y) \quad F(Z) \end{array} = \begin{array}{c} F(X) \quad F(Y) \quad F(Z) \\ \diagdown \quad \diagup \quad \diagdown \\ \mu \\ \hline F(C_{X,Y}) \\ \hline \mu^{-1} \\ \diagup \quad \diagdown \\ R \quad \alpha \\ \diagup \quad \diagdown \quad \diagup \\ F(X) \quad F(Y) \quad F(Z) \end{array} \end{aligned}$$

$$\begin{array}{c}
\begin{array}{c}
F(X) \quad F(Y) \quad F(Z) \\
\diagdown \quad \diagup \quad | \\
\mu \\
\text{---} F(C_{X,Y}) \text{---} \\
\downarrow \mu^{-1} \\
\begin{array}{c}
\text{---} R \text{---} \\
\diagup \quad \diagdown \\
\alpha \quad \alpha \\
\diagdown \quad \diagup \quad | \\
F(X) \quad F(Y) \quad F(Z)
\end{array}
\end{array} \\
= \\
\begin{array}{c}
F(X) \quad F(Y) \quad F(Z) \\
\diagdown \quad \diagup \quad | \\
\mu \\
\text{---} F(C_{X,Y}) \text{---} \\
\downarrow \mu^{-1} \\
\begin{array}{c}
\mu^{-1} \\
\diagdown \quad \diagup \\
\mu \\
\text{---} F(C_{X,Y}) \text{---} \\
\downarrow \mu^{-1} \\
\begin{array}{c}
\diagdown \quad \diagup \quad | \\
F(X) \quad F(Y) \quad F(Z)
\end{array}
\end{array}
\end{array} \\
= \\
\begin{array}{c}
F(X) \quad F(Y) \quad F(Z) \\
\diagdown \quad \diagup \quad | \\
\mu \\
\text{---} F(C_{X,Y}) \text{---} \\
\downarrow \mu^{-1} \\
\begin{array}{c}
\mu^{-1} \\
\diagdown \quad \diagup \\
\mu \\
\text{---} F(C_{X,Z}) \text{---} \\
\downarrow \mu^{-1} \\
\begin{array}{c}
\diagdown \quad \diagup \quad | \\
F(X) \quad F(Y) \quad F(Z)
\end{array}
\end{array}
\end{array}
\end{array}$$

and

$$\begin{array}{c}
\theta_I^{(3)} \left(\begin{array}{c} \text{---} R \text{---} \\ | \\ \cup \end{array} \right)_{X \otimes Y \otimes Z} = \\
\begin{array}{c}
F(X) \quad F(Y) \quad F(Z) \\
\diagdown \quad \diagup \quad | \\
\begin{array}{c}
\text{---} R \text{---} \\
\diagup \quad \diagdown \\
\alpha \quad \alpha \quad \alpha \\
\diagdown \quad \diagup \quad | \\
F(X) \quad F(Y) \quad F(Z)
\end{array}
\end{array} \\
= \\
\begin{array}{c}
F(X) \quad F(Y) \quad F(Z) \\
\diagdown \quad \diagup \quad | \\
\begin{array}{c}
\text{---} R \text{---} \\
\diagup \quad \diagdown \\
\alpha \quad \mu \\
\diagdown \quad \diagup \\
\alpha \\
\downarrow \mu^{-1} \\
\begin{array}{c}
\diagdown \quad \diagup \quad | \\
F(X) \quad F(Y) \quad F(Z)
\end{array}
\end{array}
\end{array}
\end{array}$$

$$\begin{array}{c}
F(X) \quad F(Y) \quad F(Z) \\
\diagdown \quad \diagup \quad \diagup \\
\quad \mu \\
\quad \diagdown \quad \diagup \\
\quad \mu \\
\quad \downarrow \\
F(C_{X,Y \otimes Z}) \\
\downarrow \\
\mu^{-1} \\
\diagdown \quad \diagup \\
\quad \mu^{-1} \\
\diagdown \quad \diagup \\
F(X) \quad F(Y) \quad F(Z)
\end{array}
\quad \text{by Hexagon Axiom} \quad
\begin{array}{c}
F(X) \quad F(Y) \quad F(Z) \\
\diagdown \quad \diagup \quad \diagup \\
\quad \mu \\
\quad \diagdown \quad \diagup \\
\quad \mu \\
\quad \downarrow \\
F(C_{X,Z \otimes id_Y}) \\
\downarrow \\
F(id_Y \otimes C_{X,Z}) \\
\downarrow \\
\mu^{-1} \\
\diagdown \quad \diagup \\
\quad \mu^{-1} \\
\diagdown \quad \diagup \\
F(X) \quad F(Y) \quad F(Z)
\end{array}$$

$$\begin{array}{c}
F(X) \quad F(Y) \quad F(Z) \\
\diagdown \quad \diagup \quad \diagup \\
\quad \mu \\
\quad \diagdown \quad \diagup \\
\quad \mu \\
\quad \downarrow \\
\mu^{-1} \\
\diagdown \quad \diagup \\
F(C_{X,Y}) \\
\diagdown \quad \diagup \\
\quad \mu \\
\quad \downarrow \\
\mu^{-1} \\
\diagdown \quad \diagup \\
F(C_{X,Z}) \\
\diagdown \quad \diagup \\
\quad \mu \\
\quad \downarrow \\
\mu^{-1} \\
\diagdown \quad \diagup \\
\quad \mu^{-1} \\
\diagdown \quad \diagup \\
F(X) \quad F(Y) \quad F(Z)
\end{array}
=
\begin{array}{c}
F(X) \quad F(Y) \quad F(Z) \\
\diagdown \quad \diagup \quad \diagup \\
\quad \mu \\
\quad \downarrow \\
F(C_{X,Y}) \\
\downarrow \\
\mu^{-1} \\
\diagdown \quad \diagup \\
\quad \mu \\
\quad \downarrow \\
F(C_{X,Z}) \\
\downarrow \\
\mu^{-1} \\
\diagdown \quad \diagup \\
F(X) \quad F(Y) \quad F(Z)
\end{array}$$

Thus

$$\theta_I^{(3)} \left(\begin{array}{c} \text{Diagram with } R \text{ and } X \otimes Y \otimes Z \end{array} \right) = \theta_I^{(3)} \left(\begin{array}{c} \text{Diagram with } R \text{ and } X \otimes Y \otimes Z \end{array} \right) \quad \text{and} \quad \begin{array}{c} \text{Diagram with } R \text{ and } H \end{array} = \begin{array}{c} \text{Diagram with } R \text{ and } H \end{array}.$$

Similarly, we can show that (QT2) and (QT3) hold. Therefore, $(B, R, \bar{\Delta})$ is a quasitriangular bialgebra. $\square$

In the subsection, the diagram

$$\begin{array}{cc} U & V \\ & \times \\ V & U \end{array}$$

always denotes the ordinary twisted map: $x \otimes y \longrightarrow y \otimes x$.

Theorem 5.1.2. *Let H be an ordinary bialgebra and (H_1, R) an ordinary quasitriangular Hopf algebra over field k . Let f be a bialgebra homomorphism from H_1 to H . Then there exists a bialgebra B , written as $B(H_1, f, H)$, living in $({}_{H_1}\mathcal{M}, C^R)$. Here $B(H_1, f, H) = H$ as algebra, its counit is ϵ_H , and its comultiplication and antipode are*

$$\begin{array}{c} \text{Diagram 1: } B \text{ over } H \text{ with } f, fS, \text{ and } ad \end{array} \quad \text{and} \quad \begin{array}{c} \text{Diagram 2: } B \text{ over } B \text{ with } f, f, ad, \text{ and } S \end{array} \quad \text{respectively.}$$

Then B is a braided Hopf algebra if H is a finite-dimensional Hopf algebra. Furthermore, if H is an ordinary quasitriangular bialgebra, then B is a braided quasitriangular bialgebra. In particular, when $H = H_1$ and $f = id_H$, $B(H_1, f, H)$ is a braided group, called the braided group analogue of H and written as $\underline{H}$.

Proof. (i) Set $\mathcal{C} = {}_H\mathcal{M}$ and $\mathcal{D} = ({}_{H_1}\mathcal{M}, C^R)$. Let F be the functor by pull-back along f . That is, for any $(X, \alpha_X) \in {}_H\mathcal{M}$, we obtain an H_1 -module (X, α'_X) with $\alpha'_X = \alpha_X(f \otimes id_X)$, written as $(X, \alpha'_X) = F(X)$. For any morphism $g \in Hom_{\mathcal{C}}(U, V)$, define $F(g) = g$. B is a left B -module by adjoint action. Let ${}_B B$ denote the left regular B -module. It is clear that α is a natural transformation from $B \otimes F$ to F . We first show that θ_V is injective for any $V \in {}_{H_1}\mathcal{M}$. If $\theta_V(g) = \theta_V(h)$, i.e.

$$\begin{array}{c} V \quad F(BB) \\ | \quad / \\ \textcircled{g} \quad \quad \quad \\ | \quad \quad \quad \\ \text{---} \alpha_{BB} \text{---} \\ | \\ F(BB) \end{array} = \begin{array}{c} V \quad F(BB) \\ | \quad / \\ \textcircled{h} \quad \quad \quad \\ | \quad \quad \quad \\ \text{---} \alpha_{BB} \text{---} \\ | \\ F(BB) \end{array}, \text{ which implies that } \begin{array}{c} V \\ | \\ \textcircled{g} \\ | \\ B \end{array} = \begin{array}{c} V \\ | \\ \textcircled{h} \\ | \\ B \end{array}$$


where g and h are H_1 -module homomorphisms from V to B . Thus θ_V is injective. Similarly, we can show that $\theta_V^{(2)}$ and $\theta_V^{(3)}$ are injective.

By Proposition 5.1.1, B is a quasitriangular bialgebra living in $(_{H_1}\mathcal{M}, C^R)$ determined by braided reconstruction.


We first show that the comultiplication of B is the same as stated. Now we see that

$$\theta_B^{(2)}(\Delta_B)_{X \otimes Y} = \begin{array}{c} B \quad F(X) \quad F(Y) \\ | \quad \quad \quad / \quad \backslash \\ \quad \quad \quad \textcircled{\mu} \\ \quad \quad \quad / \quad \backslash \\ \text{---} \alpha_{X \otimes Y} \text{---} \\ | \\ \text{---} \mu^{-1} \text{---} \\ / \quad \backslash \\ F(X) \quad F(Y) \end{array} \dots\dots(1)$$

The left side of (1) =



=




$$\begin{array}{c}
\begin{array}{c}
B \quad F(X) \quad F(Y) \\
\downarrow \quad \downarrow \quad \downarrow \\
\Delta_H \\
\downarrow \quad \downarrow \quad \downarrow \\
\begin{array}{c}
\begin{array}{c}
\downarrow \\
R \\
\downarrow \\
f \\
\downarrow \\
S \\
\downarrow \\
f \\
\downarrow \\
ad
\end{array}
\end{array}
\end{array}
\end{array}
=
\begin{array}{c}
\begin{array}{c}
B \quad F(X) \quad F(Y) \\
\downarrow \quad \downarrow \quad \downarrow \\
\Delta_H \\
\downarrow \quad \downarrow \quad \downarrow \\
\begin{array}{c}
\begin{array}{c}
\downarrow \\
\alpha_X \\
\downarrow \\
F(X)
\end{array}
\end{array}
\end{array}
\end{array}
= \text{the right side of (1)}.
\end{array}$$

Thus relation (1) holds.

Furthermore, if H is a finite-dimensional ordinary Hopf algebra, set $\mathcal{C} = \{M \in {}_H\mathcal{M} \mid M \text{ is finite-dimensional}\}$ and the others are the same as part (i). It is clear that $\mathcal{C}$ is rigid. Thus B has a braided-antipode by Proposition 5.1.1 (ii).

We see that




Thus the antipode of B as stated.

Assume $H_1 = H$. Our functor F is the identity Functor form ${}_H\mathcal{M}$ to ${}_H\mathcal{M}$. It follows from the proof of Proposition 5.1.1 (iii) that the quasitriangular structure of $\underline{H}$ is trivial and $\bar{\Delta} = \Delta$. That is, $\underline{H}$ is a braided group. $\square$

Dually we have

Theorem 5.1.3. *Let H be an ordinary bialgebra and (H_1, r) be an ordinary coquasitriangular Hopf algebra over field k . Let f be a Hopf algebra homomorphism from H to H_1 . Then there exists a braided bialgebra B , written as $B(H, f, H_1)$, living in $({}^B\mathcal{M}, C^r)$. Here $B(H_1, f, H) = H$ as coalgebra, its unit is η_H , and its multiplication and antipode are respectively,*



Then B is a braided Hopf algebra if H is a finite-dimensional Hopf algebra. Furthermore, if H is an ordinary coquasitriangular bialgebra, then B is a braided coquasitriangular bialgebra. In particular, when $H = H_1$ and $f = id_H$, $B(H, f, H_1)$ is a braided group, called the braided group analogue of H and written as $\underline{H}$.

5.2 Braided group analogues of double cross products

In this section we show that the braided group analogue of double cross (co)product is double cross (co)product of braided group analogues.

Theorem 5.2.1. (*Factorisation theorem*) (See [83, Theorem 7.2.3]) Let X , A and H be bialgebras or Hopf algebras. Assume that j_A and j_H are bialgebra or Hopf algebra morphisms from A to X and H to X respectively. If $\xi =: m_X(j_A \otimes j_H)$ is an isomorphism from $A \otimes H$ onto X as objects in $\mathcal{C}$, then there exist morphisms

$$\alpha : H \otimes A \rightarrow A \quad \text{and} \quad \beta : H \otimes A \rightarrow H$$

such that $A_\alpha \bowtie_\beta H$ becomes a bialgebra or Hopf algebra and ξ is a bialgebra or Hopf algebra isomorphism from $A_\alpha \bowtie_\beta H$ onto X .

Proof. Set

$$\zeta =: \begin{array}{c} H \quad A \\ \downarrow \quad \downarrow \\ j_H \quad j_A \\ \downarrow \quad \downarrow \\ \xi \\ \downarrow \\ A \quad H \end{array}, \quad \alpha =: \begin{array}{c} H \quad A \\ \downarrow \quad \downarrow \\ \zeta \\ \downarrow \quad \downarrow \\ A \quad \epsilon \end{array} \quad \text{and} \quad \beta =: \begin{array}{c} H \quad A \\ \downarrow \quad \downarrow \\ \zeta \\ \downarrow \quad \downarrow \\ \epsilon \quad H \end{array}.$$

We see

$$\begin{array}{c} H \quad H \quad A \\ \downarrow \quad \downarrow \quad \downarrow \\ \zeta \quad \zeta \quad \zeta \\ \downarrow \quad \downarrow \quad \downarrow \\ j_A \quad j_H \quad j_H \\ \downarrow \quad \downarrow \quad \downarrow \\ X \end{array} = \begin{array}{c} H \quad H \quad A \\ \downarrow \quad \downarrow \quad \downarrow \\ \zeta \quad \zeta \quad \zeta \\ \downarrow \quad \downarrow \quad \downarrow \\ j_A \quad j_H \quad j_H \\ \downarrow \quad \downarrow \quad \downarrow \\ X \end{array} = \begin{array}{c} H \quad H \quad A \\ \downarrow \quad \downarrow \quad \downarrow \\ \zeta \quad \zeta \quad \zeta \\ \downarrow \quad \downarrow \quad \downarrow \\ j_A \quad j_H \quad j_H \\ \downarrow \quad \downarrow \quad \downarrow \\ X \end{array} = \begin{array}{c} H \quad H \quad A \\ \downarrow \quad \downarrow \quad \downarrow \\ \zeta \quad \zeta \quad \zeta \\ \downarrow \quad \downarrow \quad \downarrow \\ j_H \quad j_A \quad j_H \\ \downarrow \quad \downarrow \quad \downarrow \\ X \end{array} = \begin{array}{c} H \quad H \quad A \\ \downarrow \quad \downarrow \quad \downarrow \\ \zeta \quad \zeta \quad \zeta \\ \downarrow \quad \downarrow \quad \downarrow \\ j_H \quad j_H \quad j_A \\ \downarrow \quad \downarrow \quad \downarrow \\ X \end{array} = \begin{array}{c} H \quad H \quad A \\ \downarrow \quad \downarrow \quad \downarrow \\ \zeta \quad \zeta \quad \zeta \\ \downarrow \quad \downarrow \quad \downarrow \\ j_A \quad j_H \\ \downarrow \quad \downarrow \\ X \end{array}.$$

Thus

$$\begin{array}{c} H \quad H \quad A \\ \downarrow \quad \downarrow \quad \downarrow \\ \zeta \quad \zeta \quad \zeta \\ \downarrow \quad \downarrow \quad \downarrow \\ A \quad H \end{array} = \begin{array}{c} H \quad H \quad A \\ \downarrow \quad \downarrow \quad \downarrow \\ \zeta \quad \zeta \quad \zeta \\ \downarrow \quad \downarrow \quad \downarrow \\ A \quad H \end{array}. \quad \dots\dots (1)$$

Similarly we have

$$\begin{array}{c} H \quad A \quad A \\ \downarrow \quad \downarrow \quad \downarrow \\ \zeta \quad \zeta \quad \zeta \\ \downarrow \quad \downarrow \quad \downarrow \\ A \quad H \end{array} = \begin{array}{c} H \quad A \quad A \\ \downarrow \quad \downarrow \quad \downarrow \\ \zeta \quad \zeta \quad \zeta \\ \downarrow \quad \downarrow \quad \downarrow \\ A \quad H \end{array}. \quad \dots\dots (2)$$

We also have

$$\zeta(\eta \otimes id) = id \otimes \eta \text{ and } \zeta(id \otimes \eta) = \eta \otimes id . \quad \dots\dots (3)$$

It is clear that ζ is a coalgebra morphism from $H \otimes A$ to $A \otimes H$, since j_A , j_H and m_X all are coalgebra homomorphisms. Thus we have

$$\text{and } (\epsilon \otimes \epsilon)\zeta = (\epsilon \otimes \epsilon). \quad \dots\dots (4)$$

We now show that (A, α) is an H -module coalgebra:

$$\text{by (1)}$$

$$\text{and } \alpha(\eta \otimes id_A) = (id_A \otimes \epsilon)\zeta(\eta \otimes id_A) \stackrel{\text{by(3)}}{=} id_A.$$


$$\text{We see that } \epsilon \circ \alpha = (\epsilon \otimes \epsilon)\zeta \stackrel{\text{by(4)}}{=} \epsilon \otimes \epsilon \text{ and}$$

$$\text{by(4)}$$


Thus (A, α) is an H -module coalgebra. Similarly, we can show that (H, β) is an A -module coalgebra.

Now we show that conditions (M1)–(M4) in [12,p37] hold. By (3), we easily know that (M1)

holds. Next we show that (M2) holds.



Thus (M2) holds. Similarly, we can get the proofs of (M3) and (M4). Consequently, $A_\alpha \bowtie_\beta H$ is a bialgebra or Hopf algebra by [12, Corollary 1.8]. It suffices to show that ζ is a bialgebra morphism from $A_\alpha \bowtie_\beta H$ to X . Let $D = A_\alpha \bowtie_\beta H$. Since



we have that ξ is a bialgebra morphism from $A_\alpha \bowtie_\beta H$ to X by [132, Lemma 2.5]. $\square$

Theorem 5.2.2. (Co-factorisation theorem) *Let X , A and H be bialgebras or Hopf algebras. Assume that p_A and p_H are bialgebra or Hopf algebra morphisms from X to A and X to H , respectively. If $\xi = (p_A \otimes p_H)\Delta_X$ is an isomorphism from X onto $A \otimes H$ as objects in $\mathcal{C}$, then there exist morphisms:*

$$\phi : A \rightarrow H \otimes A \quad \text{and} \quad \psi : H \rightarrow H \otimes A$$

such that $A^\phi \bowtie^\psi H$ becomes a bialgebra or Hopf algebra and ξ is a bialgebra or Hopf algebra isomorphism from X to $A^\phi \bowtie^\psi H$.

Proof. Set

We can complete the proof by turning upside down the diagrams in the proof of the preceding theorem. $\square$

In the subsection, we always consider Hopf algebras over field k and the diagram

always denotes the ordinary twisted map: $x \otimes y \longrightarrow y \otimes x$. Our diagrams only denote homomorphisms between vector spaces, so two diagrams can have the additive operation.

We now investigate the relation among braided group analogues of quasitriangular Hopf algebras A and H and their double cross coproduct $D = A \bowtie^R H$. Let us recall transmutation. We denote the braided group analogue of any ordinary quasitriangular Hopf algebras H by $\underline{H}$. R is called a weak R -matrix of $A \otimes H$ if R is invertible under convolution and

Let (A, P) and (H, Q) be ordinary finite-dimensional quasitriangular Hopf algebras over field k . Let R be a weak R -matrix of $A \otimes H$. For any $U, V \in CW(A \otimes H) = \{U \in A \otimes H \mid U \text{ is a weak } R\text{-matrix and in the center of } A \otimes H\}$,


$$R_D =: \sum R' P' U' \otimes Q' (R^{-1})'' V'' \otimes P'' (R^{-1})' V' \otimes R'' Q'' U''$$

is a quasitriangular structure of D and every quasitriangular structure of D is of this form ([26, Theorem 2.9]), where $R = \sum R' \otimes R''$, etc.

Lemma 5.2.3. *Under the above discussion, then*

- (i) $\pi_A : D \rightarrow A$ and $\pi_H : D \rightarrow H$ are bialgebra or Hopf algebra homomorphisms, respectively. Here π_A and π_H are trivial action, that is, $\pi_A(h \otimes a) = \epsilon(h)a$ for any $a \in A, h \in H$.
- (ii) $B(D, \pi_A, A) = \underline{A}$ and $B(D, \pi_H, H) = \underline{H}$.
- (iii) $\pi_{\underline{A}} : \underline{D} \rightarrow \underline{A}$ and $\pi_{\underline{H}} : \underline{D} \rightarrow \underline{H}$ are bialgebra or Hopf algebra homomorphisms, respectively.


(ii) It is enough to show $\Delta_B = \Delta_{\underline{A}}$ since $B = \underline{A}$ as algebras, where $B =: B(D, \pi_A, A)$. See



and $\epsilon \circ \pi_{\underline{H}} = \epsilon$. Thus $\pi_{\underline{H}}$ is a coalgebra homomorphism.

We now investigate the relation among braided group analogues of quasitriangular Hopf algebras A and H and their double cross coproduct $D = A \bowtie^R H$.

Theorem 5.2.4. Under the above discussion, let $\xi = (\pi_{\underline{A}} \otimes \pi_{\underline{H}})\Delta_{\underline{D}}$. Then



and ξ is surjective, where ad denotes the left adjoint action of H .


(ii) Furthermore, if A and H are finite-dimensional, then ξ is a bijective map from $\underline{D}$ onto $\underline{A} \otimes \underline{H}$. That is, in braided tensor category $({}_D\mathcal{M}, C^{R_D})$, there exist morphisms ϕ and ψ such that

$$\underline{D} \cong \underline{A}^\phi \bowtie^\psi \underline{H} \quad (\text{as Hopf algebras})$$

and the isomorphism is $(\pi_{\underline{A}} \otimes \pi_{\underline{H}})\Delta_{\underline{D}}$.

(iii) If H is commutative or $V = R$, then $\xi = id_{\underline{D}}$.

Proof. (i)



(ii) By the proof of (i), ξ is bijective and

$$\bar{\xi} = \begin{array}{c} \begin{array}{cc} A & H \\ \downarrow & \downarrow \\ \text{---} \overline{V} \text{---} & \text{---} R \text{---} \\ \downarrow & \downarrow \\ \text{---} S \text{---} & \downarrow \\ \downarrow & \downarrow \\ A & H \\ \downarrow & \downarrow \\ A & H \end{array} \end{array}.$$

Applying the co-factorization theorem 5.2.2, we complete the proof of (ii).

(iii) follows from (i). $\square$


Remark 5.2.5.

Under the assumption of Theorem 5.2.4, if we set


$$\begin{array}{c} A \\ \downarrow \\ \text{---} \phi \text{---} \\ \downarrow \\ H \end{array} \quad \begin{array}{c} A \\ \downarrow \\ \text{---} R \text{---} \\ \downarrow \\ H \end{array} \quad \text{and} \quad \begin{array}{c} H \\ \downarrow \\ \text{---} \psi' \text{---} \\ \downarrow \\ H \end{array} \quad \begin{array}{c} H \\ \downarrow \\ \text{---} \bar{R} \text{---} \\ \downarrow \\ A \end{array},$$

then $A^{\phi'} \bowtie^{\psi'} H = A \bowtie^R H$ by [26, Lemma 1.3]. We now see the relation among ϕ, ψ, R, ϕ' and ψ' :


$$\phi \quad \text{by proof of Th 5.2.2} \quad \zeta(id \otimes \eta) = \begin{array}{c} A \quad \eta \\ \downarrow \\ \text{---} \bar{\xi} \text{---} \\ \downarrow \\ \Delta_D \\ \downarrow \\ \pi_H \quad \pi_A \\ \downarrow \\ H \quad A \end{array} \quad \text{by Th 5.2.4 (iii)}$$




and ψ by proof of Th 5.2.2 $\zeta(\eta \otimes id) =$



by Th 5.2.4 (iii)





Furthermore, $\psi = \psi'$ and $\phi = \phi'$ when H is commutative or $R = V = U^{-1}$. In this case, $\underline{A}^\phi \bowtie^\psi H = \underline{A}^{\phi'} \bowtie^{\psi'} H = \underline{A} \bowtie^R H$ as Hopf algebras living in braided tensor category $({}_D\mathcal{M}, C^{R_D})$.

Example 5.2.6. (cf. [26, Example 2.11]) Let A be Sweedler's four dimensional Hopf algebra H_4 and $H = k\mathbf{Z}_2$ with $\text{char} k \neq 2$. A is generated by g and x with relations

$$g^2 = 1, \quad x^2 = 0, \quad xg = -gx.$$

H is generated by a with $a^2 = 1$. Let

$$P = \frac{1}{2}(1 \otimes 1 + 1 \otimes g + g \otimes 1 - g \otimes g) + \frac{\mu}{2}(x \otimes x + x \otimes gx + gx \otimes gx - gx \otimes x);$$

$$Q = \frac{1}{2}(1 \otimes 1 + 1 \otimes a + a \otimes 1 - a \otimes a);$$

$$R = \frac{1}{2}(1 \otimes 1 + 1 \otimes a + g \otimes 1 - g \otimes a).$$

It is clear that P and Q are quasitriangular structures of A and H respectively. R is a weak R -matrix of $A \otimes H$ with $R = R^{-1}$. Let $D = A \bowtie^R H$. Thus, for any quasitriangular structure R_D

of D , the braided group analogue $\underline{D}$ of D is a double cross coproduct of braided group analogues $\underline{A}$ and $\underline{H}$. That is, in braided tensor category $({}_D\mathcal{M}, C^{R_D})$,

$$\underline{A} \bowtie^R \underline{H} = \underline{A}^\phi \bowtie^\psi \underline{H} \quad (\text{as Hopf algebras}).$$

Here ψ is trivial coaction and $\phi(z) = \sum R^{(1)} z \bar{R}^{(1)} \otimes R^{(2)} \bar{R}^{(2)}$ for any $z \in A$ with $R = \bar{R} = \sum R^{(1)} \otimes R^{(2)}$.

5.3 The factorization of ordinary bialgebras

Throughout this section, we work in the braided tensor category of vector spaces over field k with ordinary twist braiding. In this section, we give the relation between bialgebra or Hopf algebra H and its factors.

Lemma 5.3.1. (cf. [87], [104]). *If H is a finite dimensional Hopf algebra with $\text{char } k = 0$, then the following conditions are equivalent.*

- (i) H is semisimple.
- (ii) H is cosemisimple.
- (iii) $S_H^2 = \text{id}_H$.
- (iv) $\text{tr}(S_H^2) \neq 0$.

Proof. (i) and (ii) are equivalent by [87, Theorem 2.5.2].

(i) and (iii) are equivalent by [104, Proposition 2 (c)].

(iii) and (iv) are equivalent, since (iii) implies (iv) and (iv) implies (iii) by [87, Theorem 2.5.2]. $\square$

Lemma 5.3.2. (cf. [87], [104]) *If H is a finite dimensional Hopf algebra $\text{char } k > (\dim H)^2$, then the following conditions are equivalent.*

- (i) H is semisimple and cosemisimple.
- (ii) $S_H^2 = \text{id}_H$.
- (iii) $\text{tr}(S_H^2) \neq 0$.

Proof. (i) and (iii) are equivalent by [104, Proposition 2 (c)].

(i) implies (ii) by [87, Theorem 2.5.3].

Obviously, (ii) implies (iii). $\square$

When H is a finite-dimensional Hopf algebra, let Λ_H^l and Λ_H^r denote a non-zero left integral and non-zero right integral of H respectively.

Lemma 5.3.3. *Let A , H and $D = A \bowtie H$ all be finite dimensional Hopf algebra. Then*

- (i) *there are $u \in H, v \in A$ such that*

$$\Lambda_D^l = \Lambda_A^l \otimes u, \quad \Lambda_D^r = v \otimes \Lambda_H^r;$$

- (ii) *If $A \bowtie H$ is semisimple, then A and H are semisimple;*

(iii) *If $A \bowtie H$ is unimodular, then $\Lambda_D = \Lambda_A^l \otimes \Lambda_H^r$, and $A \bowtie H$ is semisimple iff A and H are semisimple.*

Proof . (i) Let $a^{(1)}, a^{(2)}, \dots, a^{(n)}$ and $h^{(1)}, h^{(2)}, \dots, h^{(m)}$ be the basis of A and H respectively. Assuming

$$\Lambda_D^l = \sum k_{ij}(a^{(i)} \otimes h^{(j)})$$

where $k_{ij} \in k$, we have that

$$a\Lambda_D^l = \epsilon(a)\Lambda_D^l$$

and

$$\sum \epsilon(a)k_{ij}(a^{(i)} \otimes h^{(j)}) = \sum k_{ij}(aa^{(i)} \otimes h^{(j)}),$$

for any $a \in A$.

Let $x_j = \sum_i k_{ij}a^{(i)}$. Considering $\{h^{(j)}\}$ is a base of H , we get x_j is a left integral of A and there exists $k_j \in k$ such that $x_j = k_j\Lambda_A^l$ for $j = 1, 2, \dots, m$.

Thus

$$\Lambda_D^l = \sum_j k_j(\Lambda_A \otimes h^{(j)}) = \Lambda_A^l \otimes u,$$

where $u = \sum_j k_j h^{(j)}$.

Similarly, we have that $\Lambda_D^r = v \otimes \Lambda_H^r$.

(ii) and (iii) follow from part (i) . $\square$

Proposition 5.3.4. *If A, H and $D = A \bowtie H$ are finite dimensional Hopf algebras, then*

- (i) *$A \bowtie H$ is semisimple and cosemisimple iff A and H are semisimple and cosemisimple;*
- (ii) *If A and H are involutory and character char k of k does not divides $\dim D$, then $A \bowtie H$, A and H are semisimple and cosemisimple.*
- (iii) *$A \bowtie H$ is cosemisimple iff A and H are cosemisimple;*
- (iv) *If $\text{char } k = 0$, then D is semisimple iff D is cosemisimple iff A and H are semisimple iff A and H are cosemisimple iff $S_D^2 = \text{id}$ iff $S_A^2 = \text{id}$ and $S_H^2 = \text{id}$.*

Proof . (i) It follows from [104, Proposition 2].

(ii) It follows from [67, Theorem 4.3] and part (i);

(iii) If $A \bowtie H$ is cosemisimple, then $(A \bowtie H)^*$ is semisimple. Thus A and H are cosemisimple.

Conversely, if A and H are cosemisimple, then $(A \bowtie H)^*$ is semisimple. Thus $A \bowtie H$ is cosemisimple.

(iv) It follows from Lemma 5.3.1. $\square$

Chapter 6

Braided Lie Algebras

The theory of Lie superalgebras has been developed systematically, which includes the representation theory and classifications of simple Lie superalgebras and their varieties [65] [16]. In many physical applications or in pure mathematical interest, one has to consider not only $\mathbf{Z}_2$ - or $\mathbf{Z}$ -grading but also G -grading of Lie algebras, where G is an abelian group equipped with a skew symmetric bilinear form given by a 2-cocycle. Lie algebras in symmetric and more general categories were discussed in [51] and [50]. A sophisticated multilinear version of the Lie bracket was considered in [66] [98]. Various generalized Lie algebras have already appeared under different names, e.g. Lie color algebras, ϵ Lie algebras [110], quantum and braided Lie algebras, generalized Lie algebras [14] and H -Lie algebras [15].

In [81], Majid introduced braided Lie algebras from geometrical point of view, which have attracted attention in mathematics and mathematical physics (see e.g. [83] and references therein).

In this chapter, braided m-Lie algebras induced by multiplication are introduced, which generalize Lie algebras, Lie color algebras and quantum Lie algebras. The necessary and sufficient conditions for the braided m-Lie algebras to be strict Jacobi braided Lie algebras are given. Two classes of braided m-Lie algebras are given, which are generalized matrix braided m-Lie algebras and braided m-Lie subalgebras of $End_F M$, where M is a Yetter-Drinfeld module over B with $\dim B < \infty$. In particular, generalized classical braided m-Lie algebras $sl_{q,f}(GM_G(A), F)$ and $osp_{q,t}(GM_G(A), M, F)$ of generalized matrix algebra $GM_G(A)$ are constructed and their connection with special generalized matrix Lie superalgebra $sl_{s,f}(GM_{\mathbf{Z}_2}(A^s), F)$ and orthosymplectic generalized matrix Lie super algebra $osp_{s,t}(GM_{\mathbf{Z}_2}(A^s), M^s, F)$ are established. The relationship between representations of braided m-Lie algebras and their associated algebras are established.

Throughout, F is a field, G is an additive group and r is a bicharacter of G ; $|x|$ denotes the degree of x and $(\mathcal{C}, C)$ is a braided tensor category with braiding C . We write $W \otimes f$ for $id_W \otimes f$ and $f \otimes W$ for $f \otimes id_W$. Algebras discussed here may not have unity element.

6.1 Braided m-Lie Algebras

In this section we introduce braided m-Lie algebras and (strict) Jacobi braided Lie algebras. We give the necessary and sufficient conditions for the braided m-Lie algebras to be strict Jacobi braided Lie algebras.

Definition 6.1.1. *Let $(L, [\cdot])$ be an object in the braided tensor category $(\mathcal{C}, C)$ with morphism $[\cdot] : L \otimes L \rightarrow L$. If there exists an algebra (A, m) in $(\mathcal{C}, C)$ and monomorphism $\phi : L \rightarrow A$ such that $\phi[\cdot] = m(\phi \otimes \phi) - m(\phi \otimes \phi)C_{L,L}$, then $(L, [\cdot])$ is called a braided m-Lie algebra in $(\mathcal{C}, C)$ induced by multiplication of A through ϕ . Algebra (A, m) is called an algebra associated to $(L, [\cdot])$.*

A Lie algebra is a braided m-Lie algebra in the category of ordinary vector spaces, a Lie color algebra is a braided m-Lie algebra in symmetric braided tensor category $(\mathcal{M}^{FG}, C^r)$ since the canonical map $\sigma : L \rightarrow U(L)$ is injective (see [110, Proposition 4.1]), a quantum Lie algebra is a braided m-Lie algebra in the Yetter-Drinfeld category $({}^B_B\mathcal{YD}, C)$ by [49, Definition 2.1 and Lemma 2.2]), and a “good” braided Lie algebra is a braided m-Lie algebra in the Yetter-Drinfeld category $({}^B_B\mathcal{YD}, C)$ by [49, Definition 3.6 and Lemma 3.7]). For a cotriangular Hopf algebra (H, r) , the (H, r) -Lie algebra defined in [15, 4.1] is a braided m-Lie algebra in the braided tensor category $({}^H\mathcal{M}, C^r)$. Therefore, the braided m-Lie algebras generalize most known generalized Lie algebras.

For an algebra (A, m) in $(\mathcal{C}, C)$, obviously $L = A$ is a braided m-Lie algebra under operation $[\cdot] = m - mC_{L,L}$, which is induced by A through id_A . This braided m-Lie algebra is written as A^- .

If V is an object in $\mathcal{C}$ and $C_{V,V} = C_{V,V}^{-1}$, then we say that the braiding is symmetric on V .

Example 6.1.2. *If H is a braided Hopf algebra in the Yetter-Drinfeld module category $({}^B_B\mathcal{YD}, C)$ with $B = FG$ and $C(x \otimes y) = r(|y|, |x|)y \otimes x$ for any homogeneous elements $x, y \in H$, then $P(H) = \{x \in H \mid x \text{ is a primitive element}\}$ is a braided m-Lie algebra iff the braiding C is symmetric on $P(H)$.*

Indeed, it is easy to check that $P(H)$ is the Yetter-Drinfeld module. By simple computation we have $\Delta([x, y]) = [x, y] \otimes 1 + 1 \otimes [x, y] + (1 - r(|x|, |y|)r(|y|, |x|))x \otimes y$ for any homogeneous elements $x, y \in P(H)$. Thus $[x, y] \in P(H)$ iff $r(|x|, |y|)r(|y|, |x|) = 1$, as asserted.

Theorem 6.1.3. *Let $(L, [\cdot])$ be a braided m-Lie algebra in $(\mathcal{C}, C)$.*

(i) $(L, [\cdot])$ satisfies the braided anti-symmetry (or quantum anti-symmetry):

$$(BAS): [\cdot] = -[\cdot]C_{L,L}$$

if and only if $mC_{L,L} = mC_{L,L}^{-1}$.

(ii) If the braided anti-symmetry holds, then braided m-Lie algebra $(L, [\cdot], m)$ satisfies the (left) braided primitive Jacobi identity:

$$(BJI): [\cdot](L \otimes [\cdot]) + [\cdot](L \otimes [\cdot])(L \otimes C_{L,L}^{-1})(C_{L,L} \otimes L) + [\cdot](L \otimes [\cdot])(C_{L,L}^{-1} \otimes L)(L \otimes C_{L,L}) = 0,$$

and the right braided primitive Jacobi identity:

$$(BJI'): [\cdot]([\cdot] \otimes L) + [\cdot]([\cdot] \otimes L)(L \otimes C_{L,L}^{-1})(C_{L,L} \otimes L) + [\cdot]([\cdot] \otimes L)(C_{L,L}^{-1} \otimes L)(L \otimes C_{L,L}) = 0.$$

Proof . (i) Assume that $(L, [\cdot])$ satisfies the braided anti-symmetry. Since $[\cdot] = -[\cdot]C$ we have $m - mC = mCC - mC$ and $m = mCC$, which implies $mC = mC^{-1}$. The necessity is clear.
(ii)

$$\begin{aligned}
\text{l.h.s. of (BJI)} &= m(L \otimes m) - m(L \otimes m)(L \otimes C) \\
&\quad - mC(L \otimes m) + mC(L \otimes m)(L \otimes C) \\
&\quad + m(L \otimes m)(L \otimes C^{-1})(C \otimes L) \\
&\quad + m(L \otimes m)(L \otimes C)(L \otimes C^{-1})(C \otimes L) \\
&\quad - mC(L \otimes m)(L \otimes C^{-1})(C \otimes L) \\
&\quad + mC(L \otimes m)(L \otimes C)(L \otimes C^{-1})(C \otimes L) \\
&\quad + m(L \otimes m)(C^{-1} \otimes L)(L \otimes C) \\
&\quad - m(L \otimes m)(L \otimes C)(C^{-1} \otimes L)(L \otimes C) \\
&\quad - mC(L \otimes m)(C^{-1} \otimes L)(L \otimes C) \\
&\quad + mC(L \otimes m)(L \otimes C)(C^{-1} \otimes L)(L \otimes C).
\end{aligned}$$

We first check that $-(6\text{th term}) = 12\text{th term}$ and $4\text{th term} = -(10\text{th term})$. Indeed,

$$\begin{aligned}
12\text{th term} &= mC^{-1}(L \otimes m)(L \otimes C^{-1})(C^{-1} \otimes L)(L \otimes C) \\
&= m(m \otimes L)(L \otimes C^{-1})(C^{-1} \otimes L)(L \otimes C^{-1})(C^{-1} \otimes L)(L \otimes C) \\
&= m(L \otimes m)(L \otimes C)(C^{-1} \otimes L)(L \otimes C^{-1})(C^{-1} \otimes L)(L \otimes C) \\
&= m(L \otimes m)(C^{-1} \otimes L)(L \otimes C^{-1})(C \otimes L)(C^{-1} \otimes L)(L \otimes C) \\
&= -(6\text{th term}).
\end{aligned}$$

$$\begin{aligned}
4\text{th term} &= m(m \otimes L)(C^{-1} \otimes L)(L \otimes C^{-1})(C^{-1} \otimes L) \\
&= m(L \otimes m)(L \otimes C^{-1})(C^{-1} \otimes L)(L \otimes C) \\
&= -(10\text{th term}).
\end{aligned}$$

We can similarly show that $1\text{st term} = -(11\text{th term})$, $-(2\text{nd term}) = 8\text{th term}$, $-(3\text{rd term}) = 5\text{th term}$, $-(7\text{th term}) = 9\text{th term}$. Consequently, (BJI) holds. We can similarly show that $(BJI)'$ holds. $\square$

Readers can prove the above with the help of braiding diagrams.

Definition 6.1.4. Let $[\cdot]$ be a morphism from $L \otimes L$ to L in $\mathcal{C}$. If (BJI) holds, then $(L, [\cdot])$ is called a (left) Jacobi braided Lie algebra. If both of (BAS) and (BJI) hold then $(L, [\cdot])$ is called a (left) strict Jacobi braided Lie algebra.

Dually, we can define right Jacobi braided Lie algebras and right strict Jacobi braided Lie algebras. Left (strict) Jacobi braided Lie algebras are called (strict) Jacobi braided Lie algebras in general.

By Theorem 6.1.3 we have

Corollary 6.1.5. *If $(L, [\])$ is a braided m -Lie algebra, then the following conditions are equivalent:*

- (i) $(L, [\])$ is a left (right) strict Jacobi braided Lie algebra.
- (ii) $mC_{L,L} = mC_{L,L}^{-1}$.
- (iii) $[\]C_{L,L} = [\]C_{L,L}^{-1}$.

Furthermore, if L is a space graded by G with bicharacter r , then the braided primitive Jacobi identity (BJI) becomes:

$$r(|c|, |a|)[a, [b, c]] + r(|b|, |a|)[b, [c, a]] + r(|c|, |b|)[c, [a, b]] = 0 \quad (*)$$

for any homogeneous elements $a, b, c \in L$. That is, L is a Jacobi braided Lie algebra if and only if $(*)$ holds. For convenience, we let $J(a, b, c)$ denote the left hand side of $(*)$.

We now recall the (left) braided Lie algebra defined by Majid [81, Definition 4.1]. A (left) braided Lie algebra in $\mathcal{C}$ is a coalgebra (L, Δ, ϵ) in $\mathcal{C}$, equipped with a morphism $[\] : L \otimes L \rightarrow L$ satisfying the axioms:

- (L1) $([\])(L \otimes [\]) = [\]([\] \otimes [\])(L \otimes C \otimes L)(\Delta \otimes L \otimes L)$
- (L2) $C([\] \otimes L)(L \otimes C)(\Delta \otimes L) = (L \otimes [\])(\Delta \otimes L)$
- (L3) $[\]$ is a coalgebra morphism in $(\mathcal{C}, C)$.

Axiom (L1) is called the left braided Jacobi identity.

There exist braided m -Lie algebras which are neither braided Lie algebras nor Jacobi braided Lie algebras as is seen from the following example.

Example 6.1.6. (see [83]) Let $L = F\{x\}/\langle x^n \rangle$ be an algebra in $({}^F\mathbf{Z}_n\mathcal{M}, C^r)$ with a primitive n th root q of 1 and $r(k, m) = q^{km}$ for any $k, m \in \mathbf{Z}_n$, where n is a natural number.

(i) If $n > 3$, then $(L, [\])$ is a braided m -Lie algebra but is neither a Jacobi braided Lie algebra nor a braided Lie algebra of Majid.

(ii) If $n = 3$, then $(L, [\])$ is a braided m -Lie algebra and a Jacobi braided Lie algebra but is neither a strict Jacobi braided Lie algebra nor a braided Lie algebra of Majid.

Proof. (i) Since $J(x, x, x) = 3qx^3(1 - q - q^2 + q^3) \neq 0$ we have that $(L, [\])$ is not a Jacobi braided Lie algebra. If $(L, [\], \Delta, \epsilon)$ is a braided Lie algebra, since $[\]$ is a coalgebra homomorphism, we have that

$$\begin{aligned} [1, 1] &= 0, \text{ implying } \epsilon(1) = 0 \\ [x, x] &= x^2(1 - q), \text{ implying } \epsilon(x^2)(1 - q) = \epsilon(x)^2 \\ [x, x^2] &= x^3(1 - q^2), \text{ implying } \epsilon(x^3)(1 - q^2) = \epsilon(x)\epsilon(x^2) \\ &\dots \\ [x, x^{n-1}] &= x^n(1 - q^{n-1}), \text{ implying } 0 = \epsilon(x^n)(1 - q^{n-1}) = \epsilon(x)\epsilon(x^{n-1}). \end{aligned}$$

Thus $\epsilon(x^m) = 0$ for $m = 0, 1, 2, \dots, n-1$, which contradicts the fact that (L, Δ, ϵ) is a coalgebra.

(ii) Since $J(x^k, x^l, x^m) = 0$ for any $k, l, m \in \mathbf{Z}_3$, we have that $(L, [\])$ is a Jacobi braided Lie algebra. It follows from $[x, x] = x^2(1 - q) \neq -q[x, x]$ that $(L, [\])$ is not a strict Jacobi braided Lie algebra. $\square$

Note that L in the above example may never be a braided Lie algebra in the vector space category ${}_F\mathcal{M}$ with the ordinary flip τ (i.e. $\tau(x \otimes y) = y \otimes x$) as braiding, although an extension of L may become a braided Lie algebra in ${}_F\mathcal{M}$. Furthermore, for the algebra $L = F\{x\}$ in $({}^F\mathbf{Z}\mathcal{M}, C^r)$ with $q^2 \neq 1$ and $r(k, m) = q^{km}$ for any $k, m \in \mathbf{Z}$, the above conclusion holds.

Definition 6.1.7. Let $(L, [\])$ be a braided m -Lie algebra in $(\mathcal{C}, C)$. If M is an object and there exists a morphism $\alpha : L \otimes M \rightarrow M$ such that $\alpha([\] \otimes M) = \alpha(L \otimes \alpha) - \alpha(L \otimes \alpha)(C \otimes M)$, then (M, α) is called an L -module.

6.2 Generalized Matrix Braided m-Lie Algebras

As examples of the braided m-Lie algebras, we introduce the concepts of generalized matrix algebras (see [138]) and generalized matrix braided m-Lie algebras. We construct generalized classical braided m-Lie algebras $sl_{q,f}(GM_G(A), F)$ and $osp_{q,t}(GM_G(A), M, F)$ of generalized matrix algebra $GM_G(A)$. We show how generalized matrix Lie color algebras are related to Lie superalgebras for any abelian group G . That is, we establish the relationship between generalized matrix Lie color algebras and Lie superalgebras.

Let I be a set. For any $i, j, l, k \in I$, we choose a vector space A_{ij} over field F and an F -linear map μ_{ijl} from $A_{ij} \otimes A_{jl}$ into A_{il} (written $\mu_{ijl}(x, y) = xy$) such that $x(yz) = (xy)z$ for any $x \in A_{ij}$, $y \in A_{jl}$, $z \in A_{lk}$. Let A be the external direct sum of $\{A_{ij} \mid i, j \in I\}$. We define the multiplication in A as

$$xy = \left\{ \sum_k x_{ik} y_{kj} \right\}$$

for any $x = \{x_{ij}\}, y = \{y_{ij}\} \in A$. It is easy to check that A is an algebra (possibly without unit). We call A a generalized matrix algebra, or a gm algebra in short, written as $A = \sum \{A_{ij} \mid i, j \in I\}$ or $GM_I(A)$. Every element in A is called a generalized matrix. We can easily define gm ideals and gm subalgebras. It is easy to define upper triangular generalized matrices, strictly upper triangular generalized matrices and diagonal generalized matrices under some total order $\prec$ of I .

Proposition 6.2.1. Let $A = \sum \{A_{ij} \mid i, j \in I\}$ be a gm algebra and G an abelian group with $G = I$. Then A is an algebra graded by G with $A_g = \sum_{i=j+g} A_{ij}$ for any $g \in G$. In this case, the gradation is called a generalized matrix gradation, or gm gradation in short.

Proof. For any $g, h \in G$, see that

$$\begin{aligned} A_g A_h &= \left(\sum_{i=j+g} A_{ij} \right) \left(\sum_{s=t+h} A_{st} \right) \\ &\subseteq \sum_{i=t+h+g} A_{i,t+h} A_{t+h,t} \\ &\subseteq A_{g+h}. \end{aligned}$$

Thus $A = \sum \{A_{ij} \mid i, j \in I\} = \sum_{g \in G} A_g$ is a G -graded algebra. $\square$

If $\mathcal{C}$ is a small preadditive category and $A_{ij} = \text{Hom}_{\mathcal{C}}(j, i)$ is a vector space over F for any $i, j \in I = \text{objects in } \mathcal{C}$, then we may easily show that $\sum \{A_{ij} \mid i, j \in I\}$ is a generalized matrix algebra. Furthermore, if $V = \oplus_{g \in G} V_g$ is a graded vector space over field F with $A_{gh} = \text{Hom}_F(V_h, V_g)$ for any $g, h \in G$, then we call braided m-Lie algebra $A = \sum \{A_{ij} \mid i, j \in G\}$ the general linear braided m-Lie algebra, written as $gl(\{V_g\}, F)$. If $\dim V_g = n_g < \infty$ for any $g \in G$, then $gl(\{V_g\}, F)$ is written as $gl(\{n_g\}, F)$. Its braided m-Lie subalgebras are called linear braided m-Lie algebras. In fact, $gl(\{n_g\}, F) = \{f \in \text{End}_F V \mid \ker f \text{ is finite codimensional}\}$. When G is finite, we may view $gl(\{n_g\}, F)$ as a block matrix algebra over F . When $G = 0$, we denote $gl(\{n_g\}, F)$ by $gl(n, F)$, which is the ordinary ungraded general linear Lie algebra.

Assume that D is a directed graph (D is possibly an infinite directed graph and also possibly not a simple graph). Let I denote the vertex set of D , x_{ij} an arrow from i to j and $x = (x_{i_1 i_2}, x_{i_2 i_3}, \dots, x_{i_{n-1} i_n})$ a path from i_1 to i_n via arrows $x_{i_1 i_2}, x_{i_2 i_3}, \dots, x_{i_{n-1} i_n}$. For two paths $x = (x_{i_1 i_2}, x_{i_2 i_3}, \dots, x_{i_{n-1} i_n})$ and $y = (y_{j_1 j_2}, y_{j_2 j_3}, \dots, y_{j_{m-1} j_m})$ of D with $i_n = j_1$, we define the multiplication of x and y as

$$xy = (x_{i_1 i_2}, x_{i_2 i_3}, \dots, x_{i_{n-1} i_n}, y_{j_1 j_2}, y_{j_2 j_3}, \dots, y_{j_{m-1} j_m}).$$

Let A_{ij} denote the vector space over field F with basis being all paths from i to j , where $i, j \in I$. Notice that we view every vertex i of D as a path from i to i , written e_{ii} and $e_{ii}x_{ij} = x_{ij}e_{jj} = x_{ij}$. We can naturally define a linear map from $A_{ij} \otimes A_{jk}$ to A_{ik} as $x \otimes y = xy$ for any two pathes $x \in A_{ij}, y \in A_{jk}$. We may easily show that $\sum \{A_{ij} \mid i, j \in I\}$ is a generalized matrix algebra, which is called a path algebra, written as $A(D)$ (see, [3, Chapter 3]).

Example 6.2.2. *There are finite-dimensional braided m-Lie algebras in braided tensor category $({}^{\mathbf{FZ}_3}\mathcal{M}, C^r)$ with a primitive 3th root q of 1 and $r(k, m) = q^{km}$ for any $k, m \in \mathbf{Z}_3$. Indeed, for any natural number n , we can construct a generalized matrix braided m-Lie algebra $A = \sum \{A_{ij} \mid i, j \in \mathbf{Z}_3\}$ such that $\dim A = n$.*

(i) *Let $A_{ij} = 0$ when $i \neq j$ but $A_{11} = F$. Thus $\dim A = 1$.*

(ii) *Let $A_{ij} = 0$ when $i \neq j$ but $A_{11} = A_{22} = F$. Thus $\dim A = 2$.*

(iii) *Let $A_{ij} = 0$ when $i \neq j$ but $A_{11} = A_{22} = A_{33} = F$. Thus $\dim A = 3$.*

(iv) *Let D be a directed graph with vertex set $\mathbf{Z}_3$ and only one arrow from 1 to 2. Set $A = A(D)$. It is clear $\dim(A_{ij}) = 0$ when $i \neq j$ but $\dim(A_{11}) = \dim(A_{22}) = \dim(A_{33}) = \dim(A_{12}) = 1$. Thus $\dim A = 4$.*

(v) *Let D be a directed graph with vertex set $\mathbf{Z}_3$ and only two arrows: one from 1 to 2 and other one from 1 to 3. Set $A = A(D)$. It is clear $\dim(A_{ij}) = 0$ but $\dim(A_{11}) = \dim(A_{22}) = \dim(A_{33}) = \dim(A_{12}) = \dim(A_{13}) = 1$. Thus $\dim A = 5$.*

(vi) *Let D be a directed graph with vertex set $\mathbf{Z}_3$ and only $n + 2$ arrows: one from 1 to 2, one from 2 to 3 and the others from 1 to 3. Set $A = A(D)$. It is clear $\dim(A_{ij}) = 0$ but $\dim(A_{11}) = \dim(A_{22}) = \dim(A_{33}) = \dim(A_{12}) = \dim(A_{23}) = 1$ and $\dim(A_{13}) = n + 1$. Thus $\dim A = n + 6$ for $n = 0, 1, \dots$.*

Let A be a braided m-Lie algebra, G be an abelian group with a bicharacter r and W be a vector space over F .

Definition 6.2.3. If f is an F -linear map from $GM_G(A)$ to W and satisfies the following

$$(i) f(a) = \sum_{g \in G} r(g, g) f(a_{gg})$$

(ii) $f(a_{ij}b_{ji}) = f(b_{ji}a_{ij})$ for any $a, b \in A$ and $i, j \in G$, then f is called a generalized quantum trace function from gm algebra $GM_G(A)$ to W , written $tr_{q,f}$.

Set

$$sl_{q,f}(GM_G(A), F) = \{a \in GM_G(A) \mid tr_{q,f}(a) = 0\}.$$

By computation,

$$tr_{q,f}[a, b] = r(u, u)^{-1} r(g, g) \sum_{g \in G} (1 - r(u, u)^2 r(u, g) r(g, u)) tr_{q,f}(b_{g, g+u} a_{g+u, g})$$

for any homogeneous elements $a, b \in A, u \in G$. Thus we get

Lemma 6.2.4. $sl_{q,f}(GM_G(A), F)$ is a braided m -Lie algebra with $[A, A] \subseteq sl_{q,f}(GM_G(A), F)$ iff

$$\sum_{g \in G} (1 - r(u, u)^2 r(u, g) r(g, u)) tr_{q,f}(b_{g, g+u} a_{g+u, g}) = 0$$

for any homogeneous elements $a, b \in A, u \in G$ with $|a| = u$ and $|b| = -u$. If, in addition, $tr_{q,f}(A_{ij}A_{ji}) \neq 0$ for any $i, j \in G$, then $sl_{q,f}(GM_G(A), F)$ is a braided m -Lie algebra with $[A, A] \subseteq sl_{q,f}(GM_G(A), F)$ iff r is a skew symmetric bicharacter.

If $f(a_{gg}) = a_{gg}$ for any $g \in G$, then from definition 6.2.3 f is a generalized quantum trace from $GM_G(A)$ to $GM_G(A)$. If $GM_G(A) = M_n(F)$ is the full matrix algebra over F with $G = \mathbf{Z}_n$, $r(i, j) = 1$ and $f(a_{gg}) = a_{gg}$ for any $i, j, g \in \mathbf{Z}_n$, then f is the ordinary trace function. If $GM_G(A) = gl(\{n_g\}, F)$ and $f(a_{gg}) = tr(a_{gg})$ (i.e. $f(a_{gg})$ is the ordinary trace of the matrix a_{gg}) for any $g \in G$, then f is the quantum trace of the graded matrix algebra $gl(\{n_g\}, F)$. In this case, $sl_{q,f}(gl(\{n_g\}, F))$ is simply written as $sl_q(\{n_g\}, F)$.

Let G be an abelian group with a bicharacter r . If t is an F -linear map from $GM_G(A)$ to $GM_G(A)$ such that $t(A_{ij}) \subseteq A_{ji}$, $(t(a))_{ij} = t(a_{ji})$ and $t(ab) = t(b)t(a)$ for any $i, j \in G, a, b \in GM_G(A)$, then t is called a generalized transpose on $GM_G(A)$. Given $0 \neq M \in GM_G(A)$, for any $u \in G$, let $osp_{q,t}(GM_G(A), M, F)_u = \{a \in GM_G(A)_u \mid t(a_{u+g, g}) M_{u+g, h} = -r(g, u) M_{g, u+h} a_{u+h, h} \text{ for any } g, h \in G\}$ and $osp_{q,t}(GM_G(A), M, F) = \oplus_{u \in G} osp(GM_G(A), M, F)_u$.

Lemma 6.2.5. $osp_{q,t}(GM_G(A), M, F)$ is a braided m -Lie algebra iff

$$\sum_{g \in G} (1 - r(u, v) r(v, u)) M_{g, h+u+v} a_{h+u+v, v+h} b_{v+h, h} = 0$$

for any homogeneous elements $a, b \in A, u, v \in G$ with $|a| = u$ and $|b| = v$. If, in addition, for any $i, j, k \in G$, $a_{ij} \neq 0$ implies $a_{ij}A_{jk} \neq 0$, then $osp_{q,t}(GM_G(A), M, F)$ is a braided m -Lie algebra iff r is a skew symmetric bicharacter.

Proof. Obviously, $osp_{q,t}(GM_G(A), M, F)_u$ is a subspace. It remains to check that $osp_{q,t}(GM_G(A), M, F)$ is closed under bracket operation. For any $a \in osp_{q,t}(GM_G(A), M, F)_u, b \in osp_{q,t}(GM_G(A), M, F)_v$

and $u, v, g, h \in G$, set $w = u + v$. See that

$$\begin{aligned}
t([a, b]_{w+g, g})M_{w+g, h} &= t((ab - r(v, u)ba)_{w+g, g})M_{w+g, h} \\
&= t(b_{v+g, g})t(a_{w+g, g+v})M_{w+g, h} \\
&\quad - r(v, u)t(a_{u+g, g})t(b_{w+g, g+u})M_{w+g, h} \\
&= r(g + v, u)r(g, v)M_{g, h+w}b_{h+w, u+h}a_{u+h, h} \\
&\quad - r(v, u)r(g + u, v)r(g, u)M_{g, h+w}a_{h+w, v+h}b_{v+h, h}, \\
-r(g, w)M_{g, w+h}[a, b]_{w+h, h} &= -r(g, w)M_{g, w+h}a_{w+h, v+h}b_{v+h, h} \\
&\quad - r(g, w)r(v, u)M_{g, w+h}b_{w+h, u+h}a_{u+h, h}.
\end{aligned}$$

Thus $[a, b] \in \text{osp}_{q, t}(GM_G(A), M, F)_w$ iff $(r(u, v)r(v, u) - 1)M_{g, w+h}a_{w+h, v+h}b_{v+h, h} = 0$ for any $g, h \in G$. $\square$

We now consider $sl_{q, f}(GM_G(A), F)$ and $\text{osp}_{q, t}(GM_G(A), M, F)$ when the bicharacter r is skew symmetric. In this case, they become Lie color algebras in $({}^FG\mathcal{M}, C^r)$, called special gm Lie color algebra and ortho-symplectic gm Lie color algebra, respectively.

It is well-known that a Lie color algebra $({}^FG\mathcal{M}, C^r)$ with finitely generated G is related to a Lie super algebra by [110, Theorem 2]. We now show how the above gm Lie color algebras are related to Lie superalgebras for any abelian group G .

Let G be an abelian group with a skew symmetric bicharacter r . Set $G_0 = \{g \in G \mid r(g, g) = 1\}$ and $G_1 = \{g \in G \mid r(g, g) = -1\}$. We define a new bicharacter: $r_0(g, h) = -1$ for $g, h \in G_1$ and $r_0(g, h) = 1$ otherwise. It is clear that r_0 is a bicharacter too. Obviously, $(r_0)_0 = r_0$ for any skew symmetric bicharacter r on G . For convenience, let L^s denote the Lie superalgebra $L = L_0 \oplus L_1$ for Lie color algebra L in $({}^FG\mathcal{M}, C^r)$, where $L_0 = \oplus_{i \in G_0} L_i$ and $L_1 = \oplus_{i \in G_1} L_i$ with $[x, y] = xy - r_0(|y|, |x|)yx$ for any $x \in L_g, y \in L_h, g, h \in G$ (see [110, Page 718]).

Let $A = \sum\{A_{ij} \mid i, j \in G\} = GM_G(A)$ be a generalized matrix algebra. Set $B_{\bar{i}, \bar{j}} = \sum_{g \in G_{\bar{i}}, h \in G_{\bar{j}}} A_{gh}$ for any $\bar{i}, \bar{j} \in \mathbf{Z}_2$ and $B = \sum\{B_{\bar{i}, \bar{j}} \mid \bar{i}, \bar{j} \in \mathbf{Z}_2\}$. We denote the generalized matrix algebra $GM_{\mathbf{Z}_2}(B)$ by $GM_{\mathbf{Z}_2}(A^s)$. For $a \in GM_G(A)$, if $b_{\bar{i}, \bar{j}} = \sum_{g \in G_{\bar{i}}, h \in G_{\bar{j}}} a_{gh}$ for any $\bar{i}, \bar{j} \in \mathbf{Z}_2$, then we denote the element b by a^s . When $G = \mathbf{Z}_2$ with $r(g, h) = (-1)^{gh}$ for any $g, h \in \mathbf{Z}_2$, we denote $tr_{q, f}$ by $tr_{s, f}$ and $\text{osp}_{q, t}$ by $\text{osp}_{s, t}$. We have

Theorem 6.2.6. (i) $sl_{q, f}(GM_G(A), F)^s = sl_{s, f}(GM_{\mathbf{Z}_2}(A^s), F)$.

(ii) $\text{osp}_{q, t}(GM_G(A), M, F)^s = \text{osp}_{s, t}(GM_{\mathbf{Z}_2}(A^s), M^s, F)$.

Proof. (i) For any $a \in GM_G(A)$, see that

$$\begin{aligned}
tr_{q, f}(a) &= \sum_{g \in G_0} r_0(g, g)tr_{q, f}(a_{gg}) + \sum_{g \in G_1} r_0(g, g)tr_{q, f}(a_{gg}) \\
&= tr_{s, f}(a).
\end{aligned}$$

This completes the proof of (i).

(ii) For any $a \in (\text{osp}_{q, t}(GM_G(A), M, F)^s)_0$ with $a \in \text{osp}_{q, t}(GM_G(A), M, F)_u$ and $u \in G_0$, let

$\bar{i}, \bar{j} \in \mathbf{Z}_2$ and we see that

$$\begin{aligned} t(a_{\bar{i}, \bar{i}})M_{\bar{i}, \bar{j}} &= \sum_{g, h \in G_{\bar{i}}, k \in G_{\bar{j}}} t(a_{gh})M_{gk} \\ &= - \sum_{h \in G_{\bar{i}}, k \in G_{\bar{j}}} r_0(h, u)M_{h, k+u}a_{k+u, k} \\ &= -M_{\bar{i}, \bar{j}}a_{\bar{j}, \bar{j}}. \end{aligned}$$

This shows $a \in osp_{s,t}(GM_{\mathbf{Z}_2}(A^s), M^s, F)$. We can similarly prove the others. $\square$

In fact, the relations (i) and (ii) above define a G -grading of Lie superalgebras $sl_{s,f}(GM_{\mathbf{Z}_2}(A^s), F)$ and $osp_{s,t}(GM_{\mathbf{Z}_2}(A^s), M^s, F)$, respectively.

We may apply the above results to $gl(\{n_g\}, F)$ with the ordinary quantum trace $tr_q(a) = \sum_{g \in G} r(g, g)tr(a_{gg})$ and the ordinary transpose $t(a) = a'$. In this case, $sl_{q,f}$ and $osp_{q,f}$ are denoted by sl_q and osp_q , respectively. We have

Corollary 6.2.7. (i) $(sl_q(gl(\{V_g\}, F))^s = sl_s(V_{\bar{0}}, V_{\bar{1}}, F)$.

(ii) $(osp_q(gl(\{V_g\}, F))^s = osp_s(V_{\bar{0}}, V_{\bar{1}}, M^s, F)$.

6.3 Braided m-Lie Algebras in the Yetter-Drinfeld Category and Their Representations

In this section, we give another class of braided m-Lie algebras. We shall show that representations of an algebra A associated to a braided m-Lie algebra L are also representations of L . Furthermore, we show that if (M, ψ) is a faithful representation of L , then representations of $End_F M$ are also representations of L .

The category is the Yetter-Drinfeld category $({}^B_B\mathcal{YD}, C)$, where B is a finite dimensional Hopf algebra and C is a braiding with $C(x, y) = \sum (x_{(-1)} \cdot y) \otimes x_{(0)}$ for any $x \in M, y \in N$.

We use the Sweedler's notation for coproducts and comodules, i.e.

$$\Delta(a) = \sum_a a_1 \otimes a_2 \quad \text{and} \quad \psi(x) = \sum_x x_{(-1)} \otimes x_{(0)}$$

when $a \in H$ a coalgebra and $x \in M$ a left H -comodule.

Lemma 6.3.1. (i) If (V, α_V, ϕ_V) and (W, α_W, ϕ_W) are two Yetter-Drinfeld modules over B with $\dim B < \infty$, then $Hom_F(V, W)$ is a Yetter-Drinfeld module under the following module operation and comodule operation: $(b \cdot f)(x) = \sum b_1 \cdot f(S(b_2) \cdot x)$ and $\phi(f) = (S^{-1} \otimes \hat{\alpha})(b_B \otimes f)$, where $\hat{\alpha}$ is defined by $(b^* \cdot f)(x) = \langle b^*, x_{(-1)} S(f(x_{(0)}))_{(-1)} \rangle_{ev} (f(x_{(0)}))_{(0)}$ for any $x \in V, f \in Hom_F(V, W), b^* \in B^*$. Here b_B denotes a coevaluation and $\langle, \rangle_{ev}$ an evaluation of B .

(ii) If (M, α_M, ϕ_M) is a Yetter-Drinfeld modules over B , $End_F M$ is an algebra in $({}^B_B\mathcal{YD}, C)$

Proof. (i) It is clear that $\sum f_{(-1)} f_{(0)}(x) = \sum (f(x_{(0)}))_{(-1)} S^{-1}(x_{(-1)}) \otimes (f(x_{(0)}))_{(0)}$ for any $x \in V, f \in Hom_F(V, W), b \in B$. Using this, we can show that $Hom_F(V, W)$ is a B -comodule. Similarly, we can show that $Hom_F(V, W)$ is a B -module. We now show

$$\sum (b \cdot f)_{(-1)} \otimes (b \cdot f)_{(0)} = \sum b_1 f_{(-1)} S(b_3) \otimes b_2 \cdot f_{(0)} \quad (6.1)$$

for any $f \in \text{Hom}_F(V, W)$, $b \in B$. For any $x \in V$, see that

$$\begin{aligned}
\sum (b \cdot f)_{(-1)} \otimes (b \cdot f)_{(0)}(x) &= \sum b_1(f(S(b_4)) \cdot x_{(0)})_{(-1)} S(b_3) S^{-1}(x_{(-1)}) \\
&\quad \otimes b_2 \cdot (f(S(b_4)) \cdot x_{(0)})_{(0)}, \\
b_1 f_{(-1)} S(b_3) \otimes (b_2 \cdot f_{(0)})(x) &= b_1 f_{(-1)} S(b_4) \otimes b_2 \cdot f_{(0)}((S(b_3) \cdot x)) \\
&= \sum b_1(f(S(b_4) \cdot x_{(0)}))_{(-1)} S(b_3) S^{-1}(x_{(-1)}) b_5 S(b_6) \\
&\quad \otimes b_2 \cdot (f(S(b_4) \cdot x_{(0)}))_{(0)} \\
&= \sum b_1(f(S(b_4)) \cdot x_{(0)})_{(-1)} S(b_3) S^{-1}(x_{(-1)}) \\
&\quad \otimes b_2 \cdot (f(S(b_4)) \cdot x_{(0)})_{(0)}.
\end{aligned}$$

Thus (6.1) holds and $\text{Hom}_F(V, W)$ is a Yetter-Drinfeld module.

(ii) Let $E = \text{End}_F M$ and m denote the multiplication of E . Now we show that m is a homomorphism of B -comodules. It is sufficient to show

$$\sum_{fg} (fg)_{(-1)} \otimes (fg)_{(0)} = \sum_{f,g} f_{(-1)} g_{(-1)} \otimes f_{(0)} g_{(0)}. \quad (6.2)$$

for any $f, g \in E$. Indeed, for any $x \in M$, see that

$$\begin{aligned}
\sum_{fg} (fg)_{(-1)} \otimes (fg)_{(0)}(x) &= \sum_x (fg(x_0))_{(-1)} S^{-1}(x_{(-1)}) \otimes (fg(x_0))_{(0)}, \\
\sum_{f,g} f_{(-1)} g_{(-1)} \otimes f_{(0)} g_{(0)}(x) &= \sum_{f,x} f_{(-1)} (g(x_0))_{(-1)} S^{-1}(x_{(-1)}) \otimes f_{(0)} ((g(x_0))_{(0)}) \\
&= \sum_{f,x} (f((g(x_0))_{(0)}))_{(-1)} \\
&\quad S^{-1}((g(x_0))_{(0)})_{(-1)2} (g(x_0))_{(-1)1} S^{-1}(x_{(-1)}) \\
&\quad \otimes (f((g(x_0))_{(0)}))_{(0)} \\
&= (fg(x_0))_{(-1)} S^{-1}(x_{(-1)}) \otimes (fg(x_0))_{(0)}.
\end{aligned}$$

Thus (6.2) holds. Similarly, we can show that m is a homomorphism of B -modules. $\square$

Example 6.3.2. Let (M, α_M, ϕ_M) be a Yetter-Drinfeld modules over B and L a subobject of $E = \text{End}_F M$. If L is closed under operation $[\] = m - mC_{L,L}$, then $(L, [\])$ is a braided m -Lie subalgebra of E^- .

By Lemma 6.3.1, we may define representations of braided m -Lie algebras in the Yetter-Drinfeld category.

Definition 6.3.3. Let $(L, [\])$ be a braided m -Lie algebra in $({}^B_B \mathcal{YD}, C)$. If M is an object in $({}^B_B \mathcal{YD}, C)$ with morphism $\psi : L \rightarrow \text{End}_F M$ such that ψ is a homomorphism of braided m -Lie algebras, i.e. $\psi[\] = m(\psi \otimes \psi) - m(\psi \otimes \psi)C_{L,L}$, then (M, ψ) is called a representation of $(L, [\])$.

Obviously, (M, ψ) is a representation of L iff (M, α) is an L -module (see definition 6.1.7), where the relation between two operations is $\alpha(a, x) = \psi(a)(x)$ for any $a \in L, x \in M$.

Proposition 6.3.4. *Let $(L, [\cdot])$ be an object in $({}^B_B\mathcal{YD}, C)$ with a morphism $[\cdot] : L \otimes L \rightarrow L$. If M is an object in $({}^B_B\mathcal{YD}, C)$ with monomorphism $\psi : L \rightarrow \text{End}_F M$ such that $\psi[\cdot] = m(\psi \otimes \psi) - m(\psi \otimes \psi)C_{L,L}$, then L is a braided m -Lie algebra and (M, ψ) is a faithful representation of $(L, [\cdot])$.*

Proof. By Lemma 6.3.1 $E = \text{End}_F M$ is an algebra in $({}^B_B\mathcal{YD}, C)$. By Definition 6.1.1, $(L, [\cdot])$ is a braided m -Lie algebra. $\square$

Proposition 6.3.5. *Let $(L, [\cdot])$ be a braided m -Lie algebra in $({}^B_B\mathcal{YD}, C)$ induced by multiplication of A through ϕ .*

- (i) *If (M, ψ) is a representation of algebra A , then $(M, \psi\phi)$ is a representation of L .*
- (ii) *If (M, ψ) is a representation of braided m -Lie algebra A^- , then $(M, \psi\phi)$ is a representation of L .*
- (iii) *If (M, ψ) is a faithful representation of L and (N, φ) is a representation of algebra $\text{End}_F M$, then $(N, \varphi\psi)$ is a representation of L .*

Proof. (i) and (ii) follow from Definition 6.3.3.

(iii) By Lemma 6.3.1, $E = \text{End}_F M$ is an algebra in $({}^B_B\mathcal{YD}, C)$. Thus L is a braided m -Lie algebra induced by E through ψ . By Proposition 6.3.5, we complete the proof. $\square$

Let $(L, [\cdot])$ be a braided m -Lie subalgebra of the path algebra $(F(D, \rho))^-$ with relations, then a representation (V, f) of D with $f_\sigma = 0$ for any $\sigma \in \rho$ is also a representation of L (see [3, Proposition II.1.7]).

Part II

Hopf Algebras Living in Symmetric Tensor Categories

Chapter 7

The Quasitriangular Hopf Algebras in Symmetric Braided Tensor Categories

In this chapter, we study the structures of Hopf algebras living in a symmetric braided tensor category $(\mathcal{C}, C)$. We obtain that (H, R) is a quasitriangular bialgebra living in $\mathcal{C}$ iff $({}_H\mathcal{M}, C^R)$ is a braided tensor category. We show that the antipode of (co)quasitriangular Hopf algebra living in $\mathcal{C}$ is invertible. Next we also prove that S^2 is inner when almost cocommutative Hopf algebra (H, R) living in $\mathcal{C}$ has an invertible antipode S . In particular, we structure the Drinfeld (co)double $D(H)$ in symmetric braided tensor category with left duality and we prove that it is (co)quasitriangular.

We give some basic concepts as follows:

Let H and A be two bialgebras in a symmetric braided tensor categories $(\mathcal{C}, C)$, and for any $U, V, W \in \text{ob}\mathcal{C}$, assume that

$$\begin{aligned} R : I &\rightarrow H \otimes H \quad , & r : H \otimes H &\rightarrow I, \\ \tau : H \otimes A &\rightarrow I \quad , & \sigma : H \otimes H &\rightarrow I, \\ P : I &\rightarrow H \otimes A \quad , & Q : I &\rightarrow H \otimes H. \end{aligned}$$

where r and R are invertible under convolution, and they are morphisms in $\mathcal{C}$.

A bialgebra $(H, m, \eta, \Delta, \epsilon)$ with convolution-invertible R in $\text{Hom}_{\mathcal{C}}(I, H \otimes H)$ is called a quasitriangular bialgebra living in braided tensor category $\mathcal{C}$ if the following conditions hold (QT1):

$$\begin{array}{c} \text{Diagram 1: } \text{A cup-shaped node (multiplication) with two vertical lines from below labeled } HH \text{ and one vertical line from below labeled } H. \text{ A circle labeled } R \text{ is on the top line.} \\ \text{Diagram 2: } \text{A cap-shaped node (comultiplication) with two vertical lines going up labeled } H \text{ and } H, \text{ and one vertical line going up labeled } H. \text{ A circle labeled } R \text{ is on the top line.} \end{array} = ;$$

(QT2):

Diagrammatic equation (QT2): The left side shows a vertical line labeled H on the left and a cup labeled HH on the right, with a circle labeled R on the line connecting them. The right side shows three vertical lines labeled H , H , and H from left to right. The first and second lines are connected by a circle labeled R at the top, and the second and third lines are connected by a circle labeled R at the top. The lines cross each other.

(QT3):

Diagrammatic equation (QT3): The left side shows two vertical lines labeled H at the bottom. The left line has a circle labeled R on its left side, and the right line has a circle labeled R on its right side. The lines cross each other. The right side shows two vertical lines labeled H at the bottom. The left line has a circle labeled R on its right side, and the right line has a circle labeled R on its left side. The lines cross each other.

A bialgebra $(H, m, \eta, \Delta, \epsilon)$ with convolution-invertible r in $\text{Hom}_{\mathcal{C}}(H \otimes H, I)$ is called a co-quasitriangular bialgebra living in braided tensor category $\mathcal{C}$ if the following conditions hold

(CQT1):

Diagrammatic equation (CQT1): The left side shows two vertical lines labeled HH on the left and a cup labeled H on the right, with a circle labeled r on the line connecting them. The right side shows two vertical lines labeled H on the left and a cup labeled HH on the right, with a circle labeled r on the line connecting them.

(CQT2):

Diagrammatic equation (CQT2): The left side shows a vertical line labeled H on the left and a cup labeled HH on the right, with a circle labeled r on the line connecting them. The right side shows three vertical lines labeled H , H , and H from left to right. The first and second lines are connected by a circle labeled r at the bottom, and the second and third lines are connected by a circle labeled r at the bottom. The lines cross each other.

(CQT3):

Diagrammatic equation (CQT3): The left side shows two vertical lines labeled H at the top. The left line has a circle labeled r on its left side, and the right line has a circle labeled r on its right side. The lines cross each other. The right side shows two vertical lines labeled H at the top. The left line has a circle labeled r on its right side, and the right line has a circle labeled r on its left side. The lines cross each other.

τ is called a skew pairing on $H \otimes A$ if the following conditions are satisfied:

(SP1)

$$\begin{array}{c} H \quad A \quad A \\ | \quad | \quad | \\ \text{---} \cup \text{---} \\ | \\ \tau \end{array} = \begin{array}{c} H \quad A \quad A \\ | \quad | \quad | \\ \text{---} \cap \text{---} \\ | \quad | \\ \tau \quad \tau \end{array} ;$$

(SP2):

$$\begin{array}{c} H \quad H \quad A \\ | \quad | \quad | \\ \text{---} \cup \text{---} \\ | \\ \tau \end{array} = \begin{array}{c} H \quad H \quad A \\ | \quad | \quad | \\ \text{---} \cap \text{---} \\ | \quad | \\ \tau \quad \tau \end{array} ;$$

(SP3)

$$\begin{array}{c} H \\ | \\ \eta \\ | \\ \tau \\ | \\ \epsilon \end{array} = \begin{array}{c} H \\ | \\ \epsilon \end{array} ;$$

(SP4)

$$\begin{array}{c} A \\ | \\ \eta_H \\ | \\ \tau \\ | \\ \epsilon \end{array} = \begin{array}{c} A \\ | \\ \epsilon \end{array} .$$

P is called a skew copairing of $H \otimes A$ if the following conditions are satisfied:

(CSP1):

$$\begin{array}{c} P \\ | \quad | \\ \text{---} \cup \text{---} \\ | \quad | \end{array} = \begin{array}{c} P \quad P \\ | \quad | \\ \text{---} \cap \text{---} \\ | \quad | \end{array} ;$$

(CSP2):

$$\begin{array}{c} P \\ | \quad | \\ \text{---} \cup \text{---} \\ | \quad | \\ H \quad H \quad A \end{array} = \begin{array}{c} P \quad P \\ | \quad | \\ \text{---} \cap \text{---} \\ | \quad | \\ H \quad H \quad A \end{array} ;$$

(CSP3):

$$\begin{array}{c} P \\ | \quad | \\ \text{---} \cup \text{---} \\ | \quad | \\ H \quad \epsilon \end{array} = \begin{array}{c} \eta_H \\ | \end{array} ;$$

(CSP 4):

$$\begin{array}{c} P \\ | \quad | \\ \text{---} \cup \text{---} \\ | \quad | \\ \epsilon \quad H \end{array} = \begin{array}{c} \eta_A \\ | \end{array} .$$

τ is called a pairing on $H \otimes A$ if the following conditions are satisfied:

(P1):

(P2):

(P3) and (SP3) are the same, (P4) and (SP4) are the same.

P is called a copairing of $H \otimes A$ if the following conditions are satisfied: (CP1):

(CP2):

(CP3) and (CSP4) are the same. (CP4) and (CSP4) are the same.

r is called almost commutative if r satisfies

R is called almost cocommutative if R satisfies

r is called an universal co- R -matrix on $H \otimes H$ if r is invertible and satisfies (P1), (P2) and (AC);

R is called an universal R -matrix of $H \otimes H$ if R is invertible and satisfies (CP1), (CP2) and (ACO);

σ is called a 2-cocycle on $H \otimes H$ if the following conditions are satisfied:

$$2-COC : \begin{array}{c} H \quad H \quad H \\ \text{Diagram 1} \end{array} = \begin{array}{c} H \quad H \quad H \\ \text{Diagram 2} \end{array} .$$

The diagram shows the 2-cocycle condition. On the left, three vertical lines labeled H at the top. The first two lines cross, with a box labeled σ on the first line after the crossing. The third line has a box labeled σ at the bottom. On the right, the same three lines, but the first line has a box labeled σ at the bottom, and the second and third lines cross, with a box labeled σ on the second line after the crossing.

σ is called an anti-2-cocycle on $H \otimes H$ if the following conditions are satisfied:

$$A2-COC : \begin{array}{c} H \quad H \quad H \\ \text{Diagram 1} \end{array} = \begin{array}{c} H \quad H \quad H \\ \text{Diagram 2} \end{array} .$$

The diagram shows the anti-2-cocycle condition. On the left, three vertical lines labeled H at the top. The first two lines cross, with a box labeled σ on the second line after the crossing. The third line has a box labeled σ at the bottom. On the right, the same three lines, but the first line has a box labeled σ at the bottom, and the second and third lines cross, with a box labeled σ on the second line after the crossing.

Q is called a 2-cycle of $H \otimes H$ if the following conditions are satisfied:

$$(2-C) : \begin{array}{c} \text{Diagram 1} \\ H \quad H \quad H \end{array} = \begin{array}{c} \text{Diagram 2} \\ H \quad H \quad H \end{array} .$$

The diagram shows the 2-cycle condition. On the left, three vertical lines labeled H at the bottom. The first two lines cross, with a box labeled Q on the first line after the crossing. The third line has a box labeled Q at the top. On the right, the same three lines, but the first line has a box labeled Q at the top, and the second and third lines cross, with a box labeled Q on the second line after the crossing.

Q is called an anti-2-cycle of $H \otimes H$ if the following conditions are satisfied:

$$A2-C : \begin{array}{c} \text{Diagram 1} \\ H \quad H \quad H \end{array} = \begin{array}{c} \text{Diagram 2} \\ H \quad H \quad H \end{array} .$$

The diagram shows the anti-2-cycle condition. On the left, three vertical lines labeled H at the bottom. The first two lines cross, with a box labeled Q on the second line after the crossing. The third line has a box labeled Q at the top. On the right, the same three lines, but the first line has a box labeled Q at the top, and the second and third lines cross, with a box labeled Q on the second line after the crossing.

Let τ^{-1} and P^{-1} denote the inverse of τ and P under the convolution respectively. It is easy to check the following:

- (i) τ is a skew pairing iff τ^{-1} is a pairing; P is a skew copairing iff P^{-1} is a copairing;
- (ii) σ is a 2-cocycle iff σ^{-1} is an anti-2-cocycle; Q is a 2-cycle iff Q^{-1} is an anti-2-cycle;
- (iii) $\tau^{-1} = \tau(S \otimes id_A)(= \tau(id_H \otimes S^{-1}))$ if H is a Hopf algebra (or A is a Hopf algebra with invertible antipode) and τ is a pairing;
- (iv) $\tau^{-1} = \tau(id_H \otimes S)(= \tau(S^{-1} \otimes id_A))$ if A is a Hopf algebra (or H is a Hopf algebra with invertible antipode) and τ is a skew pairing;

(v) $P^{-1} = (S \otimes id_A)P (= (id_H \otimes S^{-1})P)$ if H is a Hopf algebra (or A is a Hopf algebra with invertible antipode) and P is a copairing;

(vi) $P^{-1} = (id_H \otimes S)P (= S^{-1} \otimes id_A)P$ if A is a Hopf algebra (or H is a Hopf algebra with invertible antipode) and P is a skew copairing.

In this chapter, we assume that the braided tensor category $(\mathcal{C}, C)$ is symmetric and all the algebras, coalgebras, bialgebras and Hopf algebras are in the category $(\mathcal{C}, C)$, unless otherwise stated, (see [87, section 10.5] or [82, Definition 2.2]). We omit all of the proofs of the dual conclusions since they can be obtained by turning the diagrams in the proofs of the conclusions upside down. Obviously, the category $\mathcal{Vect}(k)$ of ordinary vector spaces is a symmetric braided tensor category, in which the braiding is the ordinary transposition. We can easily get the corresponding conclusions in any symmetric braided categories for most of the conclusions in $\mathcal{Vect}(k)$ by turning the ordinary transposition into a braiding. In this case, we shall omit the proofs of these conclusions.

7.1 Yang-Baxter equation

In this section, we give the relations among quantum Yang-Baxter equation, Yang-Baxter equation, (co)quasitriangular Hopf algebra and the braided tensor category.

Proposition 7.1.1. (i) If (H, R) is a quasitriangular bialgebra, then R satisfies the quantum Yang- Baxter equation:

$$(QYBE) : \quad \begin{array}{c} \text{Diagram 1: Three vertical strands labeled } H \text{ at the bottom. Each strand has a box labeled } R \text{ at the top. The strands are connected by three crossings: the left and middle strands cross, the middle and right strands cross, and the left and right strands cross.} \end{array} = \begin{array}{c} \text{Diagram 2: Three vertical strands labeled } H \text{ at the bottom. Each strand has a box labeled } R \text{ at the top. The strands are connected by three crossings: the left and middle strands cross, the middle and right strands cross, and the left and right strands cross.} \end{array} .$$

(ii) If (H, r) is a coquasitriangular bialgebra, then r satisfies the quantum co-Yang- Baxter equation:

$$(CQYBE) : \quad \begin{array}{c} \text{Diagram 1: Three vertical strands labeled } H \text{ at the top. Each strand has a box labeled } r \text{ at the bottom. The strands are connected by three crossings: the left and middle strands cross, the middle and right strands cross, and the left and right strands cross.} \end{array} = \begin{array}{c} \text{Diagram 2: Three vertical strands labeled } H \text{ at the top. Each strand has a box labeled } r \text{ at the bottom. The strands are connected by three crossings: the left and middle strands cross, the middle and right strands cross, and the left and right strands cross.} \end{array} .$$

Proof. (i) By turning the proofs of [64, Theorem VIII 2.4] into braid diagrams, we obtain the proofs.

(ii) It is the dual case of part (i). $\square$

Proposition 7.1.2. (i) If (H, m, η) is an algebra with two algebraic morphisms $\Delta : H \longrightarrow H \otimes H$ and $\epsilon : H \longrightarrow I$, then H is a bialgebra iff ${}_H\mathcal{M}$ is a tensor category;

(ii) If (H, Δ, ϵ) is a coalgebra with two coalgebraic morphisms $m : H \otimes H \longrightarrow H$ and $\eta : I \longrightarrow H$, then H is a bialgebra iff ${}^H\mathcal{M}$ is a tensor category.

Proof. (i) By turning the proofs of [64, Proposition XI.3.1] into braid diagrams, we obtain the proofs. We also can see Corollary 1.0.15.


(ii) It is a dual case of part (i). $\square$

Theorem 7.1.3. (i) (H, R) is a quasitriangular bialgebra iff $({}_H\mathcal{M}, C^R)$ is a braided tensor category.

(ii) (H, r) is a coquasitriangular bialgebra iff $({}^H\mathcal{M}, C^r)$ is a braided tensor category.

Proof. (i) The necessity can be shown by means of turning the proofs of [64, Proposition VIII. 3.1, Proposition XIII. 1.4] into braid diagrams.

The sufficiency. For left regular H -module V , using $(id_V \otimes C_{V,V}^R)(C_{V,V}^R \otimes id_V) = C_{V \otimes V, V}^R$, we have that



Applying $\eta_H \otimes \eta_H \otimes \eta_H$ on the above, we obtain (CP2). Similarly, we can get (CP1) and (ACO).

(ii) It is the dual case of part (i). $\square$

7.2 The antipode of quasitriangular Hopf algebra

In this section, we show that the antipode of (co)quasitriangular Hopf algebra is invertible, and S^2 is inner or coininner when Hopf algebra (H, R) with invertible antipode S is almost cocommutative or (H, r) with invertible antipode S is almost commutative.

Theorem 7.2.1. Let (H, R) be an almost cocommutative Hopf algebra with invertible antipode S and define

$$u = mC_{H,H}(id_H \otimes S)R.$$

Then

- (i) $u^{-1} = mC_{H,H}(id_H \otimes S^{-1})R^{-1}$;
- (ii) $S^2 = m(m \otimes id_H)(u \otimes id_H \otimes u^{-1})$.

Proof. By turning the proofs of [87, Proposition 10.1.4] into braid diagrams, we obtain the proof. $\square$

Theorem 7.2.2. *Let (H, r) be an almost commutative Hopf algebra with invertible antipode S and define*

$$\mu = rC_{H,H}(id_H \otimes S)\Delta_H \quad .$$

Then

- (i) $\mu^{-1} = r^{-1}C_{H,H}(id \otimes S^{-1})\Delta_H$;
- (ii) $S^2 = (\mu \otimes id_H \otimes \mu^{-1})(\Delta \otimes id_H)\Delta_H$.

Proof. It is a dual case of Theorem 7.2.1. $\square$

Theorem 7.2.3. *Let (H, r) be a coquasitriangular Hopf algebra and define*

$$\lambda = r(id_H \otimes S)\Delta_H \quad .$$

Then

- (i) $\lambda^{-1} = r^{-1}(S \otimes id_H)\Delta_H$;
- (ii) $S^2 = (\lambda^{-1} \otimes id_H \otimes \lambda)(\Delta \otimes id_H)\Delta_H$;
- (iii) $S^{-2} = (\lambda \otimes id_H \otimes \lambda^{-1})(\Delta \otimes id_H)\Delta_H$.

Proof. By turning the proofs of [39, Theorem 1.3] into braid diagrams, we obtain the proofs. $\square$

Theorem 7.2.4. *Let (H, R) be a quasitriangular Hopf algebra and define*

$$v = m(id_H \otimes S)R \quad .$$

Then

- (i) $v^{-1} = m(S \otimes id_H)R^{-1}$;
- (ii) $S^2 = m(m \otimes id_H)(v^{-1} \otimes id_H \otimes v)$;
- (iii) $S^{-2} = m(m \otimes id_H)(v \otimes id_H \otimes v^{-1})$.

Proof. It is a dual case of Theorem 7.2.3. $\square$

7.3 Drinfeld double

In this section, we first construct H^σ by a 2-cocycle σ , as in [39, Theorem 1.6]. Next, we show that Drinfeld (co)double is a (co)quasitriangular Hopf algebra.

Proposition 7.3.1. *Let H be a bialgebra with a 2-cocycle σ on $H \otimes H$. Define H^σ as follows: $H^\sigma = H$ as coalgebra and the multiplication*

$$m_{H^\sigma} = \text{[Braid diagram for } m_{H^\sigma} \text{]} .$$

(i) H^σ is a bialgebra;

(ii) If H is a Hopf algebra, then H^σ is also a Hopf algebra with antipode

$$S_{H^\sigma} = \text{[Braid diagram for } S_{H^\sigma} \text{]} .$$

(iii) If H is a commutative algebra, then $(H^\sigma, \hat{\sigma})$ is a symmetric coquasitriangular bialgebra, where


$$\hat{\sigma} = \text{[Braid diagram for } \hat{\sigma} \text{]} .$$



(iv) If H is a commutative Hopf algebra, then S_{H^σ} is invertible.


Proof. (i), (ii) and (iv) can be obtained by turning the proofs of [39, Theorem 1.6] into braid diagrams. We now show part (iii).


the left hand of (CQT2) =


=














by commutativity $\underline{=}$

since H is a bialgebra $\underline{=}$

$\underline{=}$ the right hand of (CQT2).

Similarly, we can check that (CQT1) and (CQT3) hold. Obviously, $\hat{\sigma}$ is invertible. Thus $(H^\sigma, \hat{\sigma})$ is coquasitriangular. $\square$

Proposition 7.3.2. *Let H be a bialgebra with a 2-cycle Q of $H \otimes H$. Define H^Q as follows: $H^Q = H$ as algebra and the comultiplication*

$$\Delta_{H^Q} = \begin{array}{c} \begin{array}{c} H^Q \\ \downarrow \end{array} \\ \begin{array}{c} \text{Diagram: A braid with three strands. The top strand has a circle labeled } Q \text{ and the bottom strand has a circle labeled } \bar{Q}. \text{ The strands cross twice.} \end{array} \end{array} .$$

Then

(i) H^Q is a bialgebra;

(ii) Moreover, when H is a Hopf algebra, H^Q is also a Hopf algebra with antipode

$$S_{H^Q} = \begin{array}{c} \begin{array}{c} H^Q \\ \downarrow \end{array} \\ \begin{array}{c} \text{Diagram: A braid with three strands. The top strand has a circle labeled } Q \text{ and the bottom strand has a circle labeled } \bar{Q}. \text{ The strands cross twice.} \end{array} \end{array} .$$

(iii) When H is a cocommutative coalgebra, $(H^Q, \hat{Q})$ is a symmetric quasitriangular bialgebra, where

$$\hat{Q} = \begin{array}{c} \begin{array}{c} \text{Diagram: A braid with two strands. The top strand has a circle labeled } Q \text{ and the bottom strand has a circle labeled } \bar{Q}. \text{ The strands cross twice.} \end{array} \\ \begin{array}{c} H \quad H \end{array} \end{array} .$$

(iv) When H is a cocommutative Hopf algebra, S_{H^Q} is invertible.

Proof. It is a dual case of Proposition 7.3.1. $\square$

Proposition 7.3.3. If H and A are bialgebras and τ is a skew pairing on $(H \otimes A)$, then

(i) $[\tau]$ is a 2-cocycle on $(A \otimes H) \otimes (A \otimes H)$, where

$$[\tau] = \begin{array}{c} \begin{array}{c} A \\ \downarrow \end{array} \quad \begin{array}{c} H \quad A \\ \downarrow \end{array} \quad \begin{array}{c} H \\ \downarrow \end{array} \\ \begin{array}{c} \text{Diagram: A braid with three strands. The top strand has a circle labeled } A \text{ and the bottom strand has a circle labeled } \epsilon. \text{ The strands cross twice.} \end{array} \end{array} .$$

(ii) τ is invertible under convolution iff $[\tau]$ is invertible. Furthermore, $[\tau]^{-1} = [\tau^{-1}]$;

(iii) When τ is invertible, $(A \otimes H)^{[\tau]}$ is a bialgebra. In this case,

$$m_{A \bowtie_{\tau} H} =$$

and we write $A \bowtie_{\tau} H$ for $(A \otimes H)^{[\tau]}$.

Proof. The proofs of part (i) and (ii) are easily gotten by turning proofs of [39, Proposition 1.5] into braid diagrams.

(iii) It follows from Proposition 7.3.1. $\square$

Remark 7.3.4.

Let τ be an invertible skew pairing on $H \otimes A$ and define:

$$\begin{array}{c} H \quad A \\ | \quad | \\ \text{---} \alpha \text{---} \\ | \\ A \end{array} = \begin{array}{c} H \quad A \\ | \quad | \\ \text{---} \tau \text{---} \\ | \quad | \\ \text{---} \bar{\tau} \text{---} \\ | \quad | \\ H \quad A \end{array}$$

and

$$\begin{array}{c} H \quad A \\ | \quad | \\ \text{---} \beta \text{---} \\ | \\ H \end{array} = \begin{array}{c} H \quad A \\ | \quad | \\ \text{---} \tau \text{---} \\ | \quad | \\ \text{---} \bar{\tau} \text{---} \\ | \quad | \\ H \quad A \end{array}.$$

It is a straightforward verification to check that $A_{\alpha} \bowtie_{\beta} H = A \bowtie_{\tau} H$.

In fact, we can also show that $A \bowtie_{\tau} H$ is a bialgebra. (see Theorem 3.1.4).

Proposition 7.3.5. Let τ be an invertible skew pairing on $H \otimes A$ and define:

$$\begin{array}{c} H \quad A \\ | \quad | \\ \text{---} \alpha \text{---} \\ | \\ A \end{array} = \begin{array}{c} H \quad A \\ | \quad | \\ \text{---} \tau \text{---} \\ | \quad | \\ \text{---} \bar{\tau} \text{---} \\ | \quad | \\ H \quad A \end{array}$$


and

Then $A_\alpha \bowtie_\beta H$ is a bialgebra.


Proof.

Similarly,

Hhus (M1) holds. We next show that (M2) holds.



and



Thus (M2) holds. Similarly, we can show that (M3) holds. Now we show that (M4) holds.

the left hand of $(M4)$ =

=

=

and

the right hand of $[M4]$ =

=

$$\begin{array}{c}
\begin{array}{cc}
H & A \\
\downarrow & \downarrow \\
\begin{array}{c} \text{---} \tau \text{---} \end{array} & \begin{array}{c} \text{---} \bar{\tau} \text{---} \end{array}
\end{array}
= \text{the left hand of (M4)}.
\end{array}$$

Thus (M4) holds. We finally show that A is an H -module coalgebra. We see that

$$\begin{array}{c}
\begin{array}{c}
\begin{array}{ccc}
H & A & A \\
\downarrow & \downarrow & \downarrow \\
\alpha & \alpha & \alpha
\end{array}
= \begin{array}{c}
\begin{array}{ccc}
H & H & A \\
\downarrow & \downarrow & \downarrow \\
\tau & \tau & \bar{\tau}
\end{array}
\end{array}
\\
\\
\begin{array}{c}
\begin{array}{ccc}
H & H & A \\
\downarrow & \downarrow & \downarrow \\
\tau & \tau & \bar{\tau}
\end{array}
= \begin{array}{c}
\begin{array}{ccc}
H & H & A \\
\downarrow & \downarrow & \downarrow \\
\tau & \tau & \bar{\tau}
\end{array}
\end{array}
\end{array}
\\
\\
\begin{array}{c}
\begin{array}{ccc}
H & H & A \\
\downarrow & \downarrow & \downarrow \\
\tau & \tau & \bar{\tau}
\end{array}
= \begin{array}{c}
\begin{array}{ccc}
H & H & A \\
\downarrow & \downarrow & \downarrow \\
\tau & \tau & \bar{\tau}
\end{array}
\end{array}
\end{array}$$

and

Therefore, A an H -module. We also see that

and

Thus A is an H -module coalgebra. Similarly, we can show that H is an A -module coalgebra.

□

$$[P] = \begin{array}{c} \textcircled{\eta_A} \\ | \\ \textcircled{P} \\ | \\ \textcircled{\eta_H} \end{array}.$$

- (ii) P is invertible under convolution iff $[P]$ is invertible. Furthermore $[P]^{-1} = [P^{-1}]$;
 (iii) Moreover, when P is invertible, $(A \otimes H)^{[P]}$ is a bialgebra. in this case,

$$\Delta_{A \bowtie^P H} =$$

and we write $A \bowtie^P H$ for $(A \otimes H)^{[P]}$.

Proof. It is a dual case of Proposition 7.3.3. $\square$

Lemma 7.3.7. Let H and A be bialgebras. Let τ be an invertible skew pairing on $(H \otimes A)$ and P be an invertible copairing of $(H \otimes A)$. Set $D = A \bowtie_{\tau} H$. Then $(D, [P])$ is almost cocommutative iff


(AC1)

(AC2)


Proof. It can be obtained by straightforward verification $\square$

Lemma 7.3.8. Let H and A be bialgebras. Let τ be an invertible skew pairing on $(H \otimes A)$ and P be an invertible copairing of $(H \otimes A)$. Set $D = A \bowtie_{\tau} H$. Then $(D, [P])$ is almost cocommutative iff

(ACO1)




(ACO2)



Proof. It is the dual case of Lemma 7.3.7. $\square$


Proposition 7.3.9. *Let A be a Hopf algebra and H be a Hopf algebra with invertible antipode. Let τ be a pairing on $(H \otimes A)$ and P be a skew copairing of $(H \otimes A)$. Set $D = A \bowtie^P H$. If*




then $(D, [\tau])$ is almost commutative.

Proof.



the right hand of (AC1) $\stackrel{by(P1)}{=}$




by assumption $\stackrel{=}{=}$



$\stackrel{by(P1)}{=}$



by assumption $\stackrel{=}{=}$



$$= \text{[Diagram 1]} = \text{[Diagram 2]} = \text{the left hand of (AC1).}$$

Thus (AC1) holds. (AC2) can similarly be shown.

Consequently, $(B, [\tau])$ is almost commutative by Lemma 7.3.7. $\square$

Proposition 7.3.10. . *Let A be a Hopf algebra and H be a Hopf algebra with invertible antipode. Let τ be a skew pairing on $(H \otimes A)$ and P be a copairing of $(H \otimes A)$. Set $D = A \bowtie_{\tau} H$. If*

$$\begin{array}{c} A \\ \text{[Diagram]} \end{array} = A \quad \text{and} \quad \begin{array}{c} H \\ \text{[Diagram]} \end{array} = H,$$

then $(D, [P])$ is almost cocommutative.

Proof. It is a dual case of Proposition 7.3.9. $\square$

Let $\Delta^{cop} = C_{H,H}\Delta$ and $m^{op} = mC_{H,H}$. We denote $(H, \Delta^{cop}, \epsilon)$ by H^{cop} and (H, m^{op}, η) by H^{op} .

Theorem 7.3.11. *Let $(\mathcal{C}, C)$ be a symmetric braided tensor category with left dual and H be a Hopf algebra with antipode S . Let ev_H and $coev_H$ denote the evaluation and coevaluation of H respectively. Set $A = (H^*)^{op}$, $\tau = ev_H C_{H,H}$ and $P = coev_H$.*

Then

- (i) τ is a skew pairing on $H \otimes A$;
- (ii) P is a copairing of $H \otimes A$;
- (iii) Moreover, when S is invertible, $(D(H), [P])$ is a quasitriangular Hopf algebra, where $D(H) = A \bowtie_{\tau} H$, called Drinfeld double of H .

Proof. Using [82, Proposition 2.4], we can get the proofs of part (i) and part (ii).

(iii) It follows from Proposition 7.3.1, Proposition 7.3.3 and Proposition 7.3.10. $\square$

Theorem 7.3.12. *Let $(\mathcal{C}, C)$ be a symmetric braided tensor category with left duality and H be a Hopf algebra with antipode S . Let ev_H and $coev_H$ denote the evaluation and coevaluation of H respectively. Set $A = (H^*)^{cop}$, $\tau = ev_H C_{H,H}$ and $P = coev_H$.*

Then

- (i) τ is a pairing on $H \otimes A$;
- (ii) P is a skew copairing of $H \otimes A$;
- (iii) Moreover, when S is invertible, $(CD(H), [\tau])$ is a coquasitriangular Hopf algebra, where $CD(H) = A \bowtie^P H$, called Drinfeld codouble of H .

Proof. Using [82, Proposition 2.4], we can get the proofs of part (i) and part (ii).

(iii) It follows from Proposition 7.3.2, Proposition 7.3.6 and Proposition 7.3.9. $\square$

Let $(\mathcal{C}, C)$ be a symmetric braided tensor category with left duality. Let A and H be bialgebras. We easily check that τ^* is a (skew copairing) copairing of $A^* \otimes H^*$ if τ is a skew pairing (pairing) of $H \otimes A$, and P^* is a skew pairing (pairing) on $A^* \otimes H^*$ if P is a copairing (skew pairing) of $H \otimes A$. Therefore we can easily check the following.

Proposition 7.3.13. *Let $(\mathcal{C}, C)$ be a symmetric braided tensor category with left duality.*

- (i) *If (H, r) is a coquasitriangular bialgebra (Hopf algebra), then $(H^*, r^* C_{H^*, H^*})$ is a quasitriangular bialgebra (Hopf algebra);*
- (ii) *If (H, R) is a quasitriangular bialgebra (Hopf algebra), then $(H^*, C_{H^*, H^*} R^*)$ is a coquasitriangular bialgebra (Hopf algebra).*

Now, we see the relation between Drinfeld double in this chapter and the one defined by Drinfeld in $\mathcal{Vect}_f(k)$ of ordinary finite-dimensional vector spaces over field k with ordinary twist map. Obviously, $\mathcal{Vect}_f(k)$ is a symmetric braided tensor category with a left duality (see [64, Example 1, P347]). For any $H \in \mathcal{Vect}_f(k)$, it has been known that $Hom_k(H, k)$ has two kinds of bialgebra structures. On one hand, it has a bialgebra structure under convolution. In this case, we write $H^\hat{*}$ for $Hom_k(H, k)$. On the other hand, $Hom_k(H, k)$ is a duality of H in braided tensor category $\mathcal{Vect}_f(k)$. Therefore it is a bialgebra. In this case, we write H^* for $Hom_k(H, k)$. The two bialgebras have the following relation:

$$H^* = (H^{op \ cop})^\hat{*} \quad \text{as bialgebra.}$$

Therefore, the definition of Drinfeld double in Theorem 7.3.11 is the same

as what Drinfeld did since $(H^*)^{op} = (H^\hat{*})^{cop}$.

Considering Theorem 7.3.12 and [87, Proposition 10.3.14], we obtain $(D(H^\hat{*}))^\hat{*} = CD(H)$ if we identify $H^{\hat{*}\hat{*}}$ as H .

Remark 7.3.14.

(It is possible that $\mathcal{C}$ is not symmetric) If U and V have left duality U^* and V^* , respectively, then $U^* \otimes V^*$ and $V^* \otimes U^*$ both are the left dualities of $U \otimes V$. Their evaluations and coevaluations are

$$d_{U \otimes V} = (d_U \otimes d_V)(id_{U^*} \otimes C_{V^*, U} \otimes id_V), \quad b_{U \otimes V} = (id_U \otimes (C_{V, U^*})^{-1} \otimes id_{V^*})(b_U \otimes b_V);$$

$$d_{U \otimes V} = d_V(id_{V^*} \otimes d_U \otimes id_V), \quad b_{U \otimes V} = (id_U \otimes b_V \otimes id_{U^*})b_U,$$

respectively. $H^{\hat{*}}$ is the left duality of braided bialgebra H under the first; H^* is the left duality of braided bialgebra H under the second.

If both V^* and $V^{*'}$ are left dualities of V , then $V^* \cong V^{*'}$ in $\mathcal{C}$. In fact, assume $d_V : V^* \otimes V \rightarrow I$ and $b_V : I \rightarrow V \otimes V^*$ are an evaluation and a coevaluation of V , respectively; $d'_V : V^{*'} \otimes V \rightarrow I$ and $b'_V : I \rightarrow V \otimes V^{*'}$ are another evaluation and another coevaluation of V , respectively. Let $\phi = (d_V \otimes id_{V^{*'}})(id_{V^*} \otimes b'_V) : V^* \rightarrow V^{*'}$ and $\psi = (d'_V \otimes id_{V^*})(id_{V^{*'}} \otimes b_V) : V^{*' \rightarrow V^*}$. It is clear that $\phi\psi = id_{V^{*'}}$ and $\psi\phi = id_{V^*}$. Thus $V^* \cong V^{*'}$.

Remark 7.3.15.

If H has a left duality H^* in braided tensor category $\mathcal{C}$ (It is possible that $\mathcal{C}$ is not symmetric), then the following conditions are equivalent:

- (i) H is of a symmetric evaluation, i.e.

$$(id_U \otimes d_H)(C_{H^*,U} \otimes id_H) = (d_H \otimes id_U)(id_{H^*} \otimes C_{U,H})$$

for $U = H, H^*$;


- (ii) $C_{U,V} = C_{V,U}^{-1}$ for $U, V = H, H^*$;
- (iii) $C_{H,H} = C_{H,H}^{-1}$;
- (iv) $C_{H^*,H^*} = C_{H^*,H^*}^{-1}$.

Remark 7.3.16.



Theorem 7.3.11 can be stated as follows:

Let H be a finite Hopf algebra (i.e. H has a left duality H^*) with $C_{H,H} = C_{H,H}^{-1}$ (It is possible that $\mathcal{C}$ is not symmetric). Set $A = (H^*)^{op}$ and $\tau = d_H C_{H,A}$. Then $(D(H), [b])$ is a quasi-triangular Hopf algebra in $\mathcal{C}$ with $[b] = \eta_A \otimes b \otimes \eta_H$.


Proof.



and



Thus



We complete the proof. $\square$

7.4 (Co) quasitriangular Hopf algebras of double cross (co)products

In this section, we study the relation among (co)quasitriangular structures of the double cross coproduct $A \bowtie^R H$ (double cross product $A \bowtie_r H$) and A and H . We omit all of the proof in this section since we can obtain them by turning the proof in [26] into diagrams.

Assume that

$$r : A \otimes H \rightarrow I \quad \text{and} \quad R : I \rightarrow A \otimes H,$$

are morphisms in $\mathcal{C}$.

R is called a weak R -matrix of $A \otimes H$ if R is an invertible copairing of $A \otimes H$. r is called a weak r -comatrix on $A \otimes H$ if r is an invertible pairing on $A \otimes H$. Obviously, R is a weak R -matrix of $A \otimes H$ iff $C_{A,H}R$ is an invertible skew copairing of $H \otimes A$. $rC_{A,H}$ is a weak r -comatrix on $H \otimes A$ iff r is an invertible copairing on $A \otimes H$.

Therefore, if r is a weak r -comatrix on $A \otimes H$ and let $\tau = rC_{A,H}$, then $A \bowtie_\tau H$ is a bialgebra by Theorem 7.3.3. For convenience, we also denote $A \bowtie_\tau H$ by $A \bowtie_r H$.

Lemma 7.4.1. *Let R be a weak R -matrix of $A \otimes H$, and define*

$$\phi = \begin{array}{c} \begin{array}{cc} A & \\ \textcircled{R} & \textcircled{\bar{R}} \\ \text{---} & \text{---} \\ \text{---} & \text{---} \\ \text{---} & \text{---} \\ H & A \end{array} \end{array} \quad \text{and} \quad \psi = \begin{array}{c} \begin{array}{cc} H & \\ \textcircled{R} & \textcircled{\bar{R}} \\ \text{---} & \text{---} \\ \text{---} & \text{---} \\ \text{---} & \text{---} \\ H & A \end{array} \end{array} .$$

Then $A^\phi \bowtie^\psi H$ is a bialgebra. In this case, we denote $A^\phi \bowtie^\psi H$ by $A \bowtie^R H$. Furthermore, $A \bowtie^R H$ is a Hopf algebra if A and H are Hopf algebras.

Considering the universal property of double bicrossproducts in Lemma 3.2.2, we have

Lemma 7.4.2. *Let $D = A \bowtie^R H$ be a bialgebra. Assume that $p_A : B \rightarrow A$ and $p_H : B \rightarrow H$ are bialgebra morphisms. Then $p = (p_A \otimes p_H)\Delta_B$ is a bialgebra morphism from $B \rightarrow D$ iff (UT):*

$$\begin{array}{c} \begin{array}{cc} B & \\ \textcircled{R} & \textcircled{\bar{R}} \\ \text{---} & \text{---} \\ \text{---} & \text{---} \\ \text{---} & \text{---} \\ A & H \end{array} \end{array} = \begin{array}{c} \begin{array}{cc} B & \\ \textcircled{p_H} & \textcircled{p_A} \\ \text{---} & \text{---} \\ \text{---} & \text{---} \\ \text{---} & \text{---} \\ A & H \end{array} \end{array} .$$

Corollary 7.4.3. *Let R be a weak R -matrix of $H \otimes H$. Then (H, R) is a quasitriangular bialgebra iff the comultiplication Δ of H is a bialgebra morphism from H to $H \otimes H$.*

Theorem 7.4.4. *$D = A \bowtie^R H$ has the following universal property with respect to (π_A, π_H) . Assume that $p_A : B \rightarrow A$ and $p_H : B \rightarrow H$ are bialgebra morphisms. If p_A and p_H satisfy (UT) defined in Lemma 7.4.2, then there exists a unique bialgebra morphism p from B to D such that $\pi_A p = p_A$ and $\pi_H p = p_H$.*

Theorem 7.4.5. *If $A^\phi \bowtie^\psi H$ admits a quasitriangular structure, then A and H admit quasitriangular structures, and there exists a weak R -matrix R of $A \otimes H$ such that $A^\phi \bowtie^\psi H = A \bowtie^R H$ and*

$$\phi = \begin{array}{c} \begin{array}{cc} & A \\ \textcircled{R} & \textcircled{\bar{R}} \\ \text{---} & \text{---} \\ \text{---} & \text{---} \\ \text{---} & \text{---} \\ H & A \end{array} \end{array} \quad \text{and} \quad \psi = \begin{array}{c} \begin{array}{cc} & H \\ \textcircled{R} & \textcircled{\bar{R}} \\ \text{---} & \text{---} \\ \text{---} & \text{---} \\ \text{---} & \text{---} \\ H & A \end{array} \end{array} .$$

If

$$\begin{array}{ll} U : I \rightarrow A \otimes H, & V : I \rightarrow A \otimes H, \\ P : I \rightarrow A \otimes A, & Q : I \rightarrow H \otimes H, \end{array}$$

are morphisms in $\mathcal{C}$, we define

$$[U, P, Q, V] = \begin{array}{c} \begin{array}{cccc} \textcircled{U} & \textcircled{P} & \textcircled{Q} & \textcircled{V} \\ \text{---} & \text{---} & \text{---} & \text{---} \\ \text{---} & \text{---} & \text{---} & \text{---} \\ \text{---} & \text{---} & \text{---} & \text{---} \\ A & H & A & H \end{array} \end{array} .$$

From now on, unless otherwise stated, R is a fixed weak R -matrix of $A \otimes H$.

Theorem 7.4.6. *If (A, P) and (H, Q) are quasitriangular bialgebras, then $(A \bowtie^R H, [R, P, Q, R^{-1}])$ is a quasitriangular bialgebra.*

Corollary 7.4.7. *$A^\phi \bowtie^\psi H$ admits a quasitriangular structure iff A and H admit a quasitriangular structure, and there exists a weak R -matrix R of $A \otimes H$ such that $A^\phi \bowtie^\psi H = A \bowtie^R H$.*

Corollary 7.4.8. *Let R be a weak R -matrix of $H \otimes H$. Then $(H \bowtie^R H, [R, R, R, R^{-1}])$ is a quasitriangular bialgebra iff (H, R) is a quasitriangular bialgebra.*

Lemma 7.4.9. *Let R be a weak R -matrix of $H \otimes H$. Then the following conditions are equivalent.*

- (i) (H, R) is a triangular bialgebra.
- (ii) $(A \bowtie^R H, [R, R, R, R^{-1}])$ is a triangular bialgebra.
- (iii) $\Delta_H : H \longrightarrow H \bowtie^R H$ is a bialgebra morphism and $[R, R, R, R^{-1}] = (\Delta_H \otimes \Delta_H)(R)$, i.e. Δ_H is a quasitriangular morphism.

for any morphisms $P : I \rightarrow A \otimes H$, $g : I \rightarrow A$, $h : I \rightarrow H$.

Let

$$QT(H) = \{P \mid P \text{ a quasitriangular structure of } H\}.$$

Lemma 7.4.10. Φ is a bijective map from $CW(A, H) \times CW(A, A) \times CW(H, H) \times CW(A, H)$ onto $CW(A \bowtie^R H, A \bowtie^R H)$, where $\Phi(U, P, Q, V) = [U, P, Q, V]$.

Lemma 7.4.11. Let R and $\bar{R}$ be two weak R -matrices of $A \otimes H$. Then $A \bowtie^R H = A \bowtie^{\bar{R}} H$ iff there exists $U \in CW(A, H)$ such that $\bar{R} = RU$.

Theorem 7.4.12. Assume (A, P) and (H, Q) are quasitriangular bialgebras and $U, V \in CW(A, H)$. Then $(A \bowtie^R H, [R, P, Q, R^{-1}][U, \eta, \eta, V])$ is a quasitriangular bialgebra. Conversely, if $(A \bowtie^R H, \mathcal{R})$ is a quasitriangular bialgebra, then there exist P, Q, U, V such that (A, P) and (H, Q) are quasitriangular bialgebras with $\mathcal{R} = [R, P, Q, R^{-1}][U, \eta, \eta, V]$ and $U, V \in CW(A, H)$.

Proposition 7.4.13. There exists a bijective map from $QT(A) \times QT(H) \times CW(A, H) \times CW(A, H)$ onto $QT(A \bowtie^R H)$ by sending (P, Q, U, V) to $[R, P, Q, R^{-1}][U, \eta_H, \eta_A, V]$.

Dually, we can get the followings about double cross coproducts.

Lemma 7.4.1° Let r be a weak r -comatrix on $A \otimes H$, and define

$$\alpha = \text{diagram} \quad \text{and} \quad \beta = \text{diagram}.$$

Then $A_\alpha \bowtie_\beta H$ is a bialgebra. In this case, we denote $A_\alpha \bowtie_\beta H$ by $A \bowtie_r H$. Furthermore, $A_\alpha \bowtie_\beta H$ is a Hopf algebra if A and H is Hopf algebras.

Lemma 7.4.2° Let $D = A \bowtie_r H$ be a bialgebra. Assume that $j_A : A \longrightarrow B$ and $j_H : H \longrightarrow B$ are bialgebra morphisms. Let $j = m_B(j_A \otimes j_H)$ be a bialgebra morphism from D to B iff (CUT):

$$\text{diagram} = \text{diagram}.$$

Corollary 7.4.3° Let r be a weak r -comatrix on $H \otimes H$. Then (H, r) is a coquasitriangular bialgebra iff the multiplication m_H of H is a bialgebra morphism from $H \otimes H$ to H .

Theorem 7.4.4° $D = A \bowtie_r H$ has the following universal property with respect to (i_A, i_H) . Assume that $j_A : A \longrightarrow B$ and $j_H : H \longrightarrow B$ are bialgebra morphisms. If j_A and j_H satisfy (CUT) defined in Lemma 7.4.2, then there exists a unique bialgebra morphism j from D to B such that $j i_A = j_A$ and $j i_H = j_H$.

Theorem 7.4.5° If $A_\alpha \bowtie_\beta H$ admits a coquasitriangular structure, then A and H admit coquasitriangular structures, and there exists a weak r -comatrix r on $A \otimes H$ such that $A_\alpha \bowtie_\beta H = A \bowtie_r H$ and

$$\alpha = \begin{array}{c} \begin{array}{cc} H & A \\ \downarrow & \downarrow \\ \text{Diagram 1} \end{array} \end{array} \quad \text{and} \quad \beta = \begin{array}{c} \begin{array}{cc} H & A \\ \downarrow & \downarrow \\ \text{Diagram 2} \end{array} \end{array} .$$

(Note: The diagrams are string diagrams representing morphisms. Diagram 1 for α has inputs H and A at the top, and outputs A and H at the bottom. It contains a box labeled r on the H line and a box labeled $\bar{r}$ on the A line. Diagram 2 for β has inputs H and A at the top, and outputs H and A at the bottom. It contains a box labeled r on the H line and a box labeled $\bar{r}$ on the A line.)

If

$$\begin{aligned} u : A \otimes H &\rightarrow I, & v : A \otimes H &\rightarrow I, \\ p : A \otimes A &\rightarrow I, & q : H \otimes H &\rightarrow I, \end{aligned}$$

are morphisms, we set

$$[u, p, q, v] = \begin{array}{c} \begin{array}{cccc} A & H & A & H \\ \downarrow & \downarrow & \downarrow & \downarrow \\ \text{Diagram 3} \end{array} \end{array} .$$

(Note: Diagram 3 is a complex string diagram with four inputs A, H, A, H at the top and four outputs at the bottom. It contains boxes labeled p , q , u , and v connected by various crossings and lines.)

Theorem 7.4.6° Assume (A, p) and (H, q) are coquasitriangular bialgebras. Then $(A \bowtie_r H, [r, p, q, r^{-1}])$ is a coquasitriangular bialgebra.


Corollary 7.4.7° $A_\alpha \bowtie_\beta H$ admits a coquasitriangular structure iff A and H admit a coquasitriangular structure, and there exists a weak r -comatrix r of $A \otimes H$ such that $A_\alpha \bowtie_\beta H = A \bowtie_r H$.

Corollary 7.4.8° Let r be a weak r -comatrix on $H \otimes H$. Then $(H \bowtie_r H, [r, r, r, r^{-1}])$ is a coquasitriangular bialgebra iff (H, r) is a coquasitriangular bialgebra.

Lemma 7.4.9° Let r be a weak r -comatrix on $H \otimes H$. Then the following are equivalent.

- (i) (H, r) is a cotriangular bialgebra.
- (ii) $(A \bowtie_r H, [r, r, r, r^{-1}])$ is a cotriangular bialgebra.
- (iii) $m_H : H \bowtie_r H \longrightarrow H$ is a bialgebra morphism and $[r, r, r, r^{-1}](m_H \otimes m_H) = r$, i.e. m_H is a coquasitriangular morphism.

Let $CCW(A \otimes H)$ denote the set of all weak r -comatrix r on $A \otimes H$ satisfying the following conditions:



for any morphisms $f : A \otimes H \rightarrow I$, $g : A \rightarrow I$, $h : H \rightarrow I$. Let

$$CQT(H) = \{p \mid p \text{ is a coquasitriangular structure of } H\}.$$

From now on, unless otherwise stated, r is a fixed weak r -comatrix on $A \otimes H$.

Lemma 7.4.10° ϕ is a bijective map from $CCW(A, H) \times CCW(A, A) \times CCW(H, H) \times CCW(A, H)$ to $CCW(A \bowtie_r H, A \bowtie_r H)$, where $\phi(u, p, q, v) = [u, p, q, v]$.

Lemma 7.4.11° Let r and $\bar{r}$ be two weak r -comatrices on $A \otimes H$. Then $A \bowtie_r H = A \bowtie_{\bar{r}} H$ iff there exists $u \in CCW(A, H)$ such that $\bar{r} = r * u$.

Theorem 7.4.12° Assume (A, p) and (H, q) are coquasitriangular bialgebras and $u, v \in CCW(A, H)$. Then $(A \bowtie_r H, [r, p, q, r^{-1}] * [u, \epsilon, \epsilon, v])$ is a coquasitriangular bialgebra. Conversely, if $(A \bowtie_r H, \bar{r})$ is a coquasitriangular bialgebra, then there exist p, q, u, v such that (A, p) and (H, q) are coquasitriangular bialgebras with $\bar{r} = [r, p, q, r^{-1}] * [u, \epsilon, \epsilon, v]$ and $u, v \in CCW(A, H)$.

Proposition 7.4.13° There exists a bijective map from $CQT(A) \times CQT(H) \times CCW(A, H) \times CCW(A, H)$ to $CQT(A \bowtie_r H)$ by sending (s, t, u, v) to $[r, s, t, r^{-1}] * [u, \epsilon, \epsilon, v]$.

Chapter 8

The Duality Theorem for Braided Hopf Algebras

In this chapter, we give the duality theorem for Hopf algebras living in a symmetric braided tensor category. Using the result, we show that

$$(R\#H)\#H^{\hat{*}} \cong M_n(R) \quad \text{as algebras}$$

when H is a finite-dimensional Hopf algebra living in symmetric braided category determined by (co)triangular structures. In particular, the duality theorem holds for any finite-dimensional super-Hopf algebra H

Supersymmetry has attracted a great deal of interest from physicists and mathematicians (see [30], [88]). It finds numerous applications in particle physics and statistical mechanics. Of central importance in supersymmetric theories is the Z_2 -graded structure which permeates them. Superalgebras and super-Hopf algebras are most naturally very important.

It is well-known that in the work of C^* -algebras and von Neumann algebras for abelian groups the crossed product R crossed by G crossed by $\hat{G}$ is isomorphic to R tensor the compact operators. This result does have implications for symmetries in quantum theory. One may see for example Pedersen's book on C^* -algebras [96]. This offers some clues as to why duality theorem might be useful in mathematical physics. Its generalization to Hopf-von Neumann algebras was known again , see for example the book of Stratila on modular theory [113] or the papers on Hopf-von Neumann algebras (or the book on Kac algebras) by Enock and Schwartz. Blattner and Montgomery strip off the functional analysis and duplicate the result at the level of Hopf algebras [20]. But if one looks more deeply into the literature one finds indeed its origins in mathematical physics. However, since most algebras in physics are not finite-dimensional one does need to say some words about what one thinks the operator-algebra version should be. What plays the role of compact operators in the super case ?

In this chapter, we show that the subalgebras $H\bar{\otimes}H^{\hat{*}}$ of $End_k H$ plays the role of compact operators in the super case. We obtain the duality theory in symmetric tensor category. Using this theory, we study super-Hopf algebras. If super(quantum)group H acts on a superalgebra

R , then the duality of the super(quantum)group acts on the crossed product. We show that the repeated crossed product is isomorphic to $M_n(R)$ when $\dim H = n$

8.1 The duality theory for Hopf algebras living in symmetric tensor categories

In this section we obtain the duality theory for Hopf algebras living in symmetric tensor category $\mathcal{C}$.

Throughout this section H is a Hopf algebra living in $\mathcal{C}$ with a left duality H^* and (R, α) is a left H -module algebra in $\mathcal{C}$. Let $H^{\hat{*}} = (H^*)^{op \ cop}$ and $R \# H$ be an algebra.

It follows from [76] that

Lemma 8.1.1. *If (R, α) is an H -module algebra, let $\phi = (id \otimes \alpha)(b' \otimes id) : R \rightarrow H^{*op} \otimes R$, where $b' = C_{H, H^{*op}} b$. Then*

$$\phi m = (m \otimes m)(id \otimes C_{R, H^*} \circ p \otimes id)(\phi \otimes \phi).$$

Lemma 8.1.2. *If H is a Hopf algebra living in symmetric tensor category with a left duality H^* , then*

(1) $(H, \rightharpoonup)$ is a left $H^{\hat{\star}}$ -module algebra under the module operation $\rightharpoonup = (id \otimes d)(C_{H^{\hat{\star}}, H} \otimes id)(id \otimes \Delta)$.

(2) $(H^{\hat{*}}, \rightharpoonup)$ is a left H -module algebra under the module operation $\rightharpoonup = (id \otimes d)(id \otimes C_{H, H^{\hat{*}}})(C_{H, H^{\hat{*}}} \otimes id)(id \otimes \Delta)$.

(3) $(H, \leftarrow)$ is a right $H^{\hat{*}}$ -module algebra under the module operation $\leftarrow = (d \otimes id)(C_{H, H^{\hat{*}}} \otimes id)(id \otimes C_{H, H^{\hat{*}}})(\Delta \otimes id)$.

(4) $(H^{\hat{*}}, \leftarrow)$ is a right H -module algebra under the module operation $\leftarrow = (d \otimes id)(id \otimes C_{H^{\hat{*}}, H})(\Delta \otimes id)$.

Lemma 8.1.3. *Let H be a Hopf algebra living in symmetric tensor category with a left duality H^* . If we define multiplication and unity on $H \otimes H^*$ as follows:*

$$\begin{array}{c} H \otimes H^{\hat{*}} \quad H \otimes H^{\hat{*}} \\ \quad \cup \quad \quad \quad \cup \\ \quad \text{---} \quad \quad \quad \text{---} \\ \quad \downarrow \quad \quad \quad \downarrow \\ H \otimes H^{\hat{*}} \end{array} = \begin{array}{c} H \quad H^{\hat{*}} \quad H \quad H^{\hat{*}} \\ \quad \cup \quad \quad \quad \cup \\ \quad \text{---} \quad \quad \quad \text{---} \\ \quad \downarrow \quad \quad \quad \downarrow \\ H \quad \quad \quad H \end{array}$$

and $\eta_{H \otimes H^*} = b_H$, then $H \otimes H^*$ is an algebra, written as $H \bar{\otimes} H^*$.

In fact, if $\mathcal{C}$ is a symmetric Yetter-Drinfeld category, then $H \bar{\otimes} H^{\hat{*}}$ can be viewed as a subalgebra of $End_k(H)$.

Lemma 8.1.4. *λ is an algebraic morphism and ρ is an anti-algebraic morphism, where $\lambda = (m \otimes d \otimes id)(id \otimes C_{H^*, H} \otimes id \otimes id)(id \otimes id \otimes \Delta \otimes id)(id \otimes id \otimes b)$ and $\rho = (d \otimes m \otimes id)(id \otimes id \otimes C_{H, H} \otimes id)(id \otimes C_{H, H} \otimes id \otimes id)(id \otimes id \otimes \Delta \otimes id)(id \otimes id \otimes b)$.*

Proof.

$$\begin{array}{c}
 H\#H^* \quad H\#H^* \\
 \downarrow \\
 \textcircled{\lambda} \\
 \downarrow \\
 H\bar{\otimes}H^*
 \end{array}
 =
 \begin{array}{c}
 H \quad H^* \quad H \quad H^* \\
 \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
 \text{Diagram 1} \\
 \downarrow \quad \downarrow \\
 H \quad H^*
 \end{array}
 =
 \begin{array}{c}
 H \quad H^* \quad H \quad H^* \\
 \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
 \text{Diagram 2} \\
 \downarrow \quad \downarrow \\
 H \quad H^*
 \end{array}$$

$$=
 \begin{array}{c}
 H \quad H^* \quad H \quad H^* \\
 \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
 \text{Diagram 3} \\
 \downarrow \quad \downarrow \\
 H \quad H^*
 \end{array}
 \quad \text{and}$$

$$\begin{array}{c}
 H\#H^* \quad H\#H^* \\
 \downarrow \quad \downarrow \\
 \textcircled{\lambda} \quad \textcircled{\lambda} \\
 \downarrow \\
 H\bar{\otimes}H^*
 \end{array}
 =
 \begin{array}{c}
 H \quad H^* \quad H \quad H^* \\
 \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
 \text{Diagram 4} \\
 \downarrow \quad \downarrow \\
 H \quad H^*
 \end{array}
 =
 \begin{array}{c}
 H \quad H^* \quad H \quad H^* \\
 \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
 \text{Diagram 5} \\
 \downarrow \quad \downarrow \\
 H \quad H^*
 \end{array}$$

$$\begin{array}{c}
H \quad H^{\hat{*}} \quad H \quad H^{\hat{*}} \\
\vdots \\
= \text{[String Diagram]} = \text{[Simplified String Diagram]} = \begin{array}{c} H \# H^{\hat{*}} \quad H^{\hat{*}} \# H \\ \downarrow \text{[Circle with } \lambda \text{]} \\ H^{\hat{\otimes}} H^{\hat{*}} \end{array}
\end{array}$$

Thus λ is an algebraic morphism.

Similarly, we can show that ρ is an anti-algebraic morphism. $\square$

Lemma 8.1.5. λ and ρ satisfy the following:

(1)

$$\begin{array}{c}
H \quad H^{\hat{*}} \quad H^{\hat{*}} \quad H \\
\vdots \\
\begin{array}{c} H \# H^{\hat{*}} \quad H^{\hat{*}} \# H \\ \downarrow \text{[Circle with } \lambda \text{]} \\ H^{\hat{\otimes}} H^{\hat{*}} \end{array} = \text{[Large String Diagram with } S, \lambda, \rho \text{]} \downarrow \\
H^{\hat{\otimes}} H^{\hat{*}}
\end{array}$$

(2)

$$\begin{array}{c}
\begin{array}{c}
H^{\hat{*}} \# H \quad H^{\hat{*}} \# H \\
\downarrow \quad \downarrow \\
\rho \quad \lambda \\
\downarrow \\
H \bar{\otimes} H^{\hat{*}}
\end{array}
=
\begin{array}{c}
\begin{array}{c}
H^{\hat{*}} \quad H \quad H \quad H^{\hat{*}} \\
\downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
\text{Diagram with } S, \rho, \lambda, \text{ and } \bar{\otimes} \text{ nodes} \\
\downarrow \\
H \bar{\otimes} H^{\hat{*}}
\end{array}
\end{array}$$

Proof. We show (1) by following five steps. It is easy to check the following (i) and (ii).

(i)

$$\begin{array}{c}
\begin{array}{c}
H \# \eta_{H^{\hat{*}}} \quad \eta_{H^{\hat{*}}} \# H \\
\downarrow \quad \downarrow \\
\lambda \quad \rho \\
\downarrow \\
H \bar{\otimes} H^{\hat{*}}
\end{array}
=
\begin{array}{c}
\begin{array}{c}
H \# \eta_{H^{\hat{*}}} \quad \eta_{H^{\hat{*}}} \# H \\
\downarrow \quad \downarrow \\
\rho \quad \lambda \\
\downarrow \\
H \bar{\otimes} H^{\hat{*}}
\end{array}
\end{array}$$

(ii)

$$\begin{array}{c}
\begin{array}{c}
\eta_H \# H^{\hat{*}} \quad H^{\hat{*}} \# \eta_H \\
\downarrow \quad \downarrow \\
\lambda \quad \rho \\
\downarrow \\
H \bar{\otimes} H^{\hat{*}}
\end{array}
=
\begin{array}{c}
\begin{array}{c}
\eta_H \# H^{\hat{*}} \quad H^{\hat{*}} \# \eta_H \\
\downarrow \quad \downarrow \\
\rho \quad \lambda \\
\downarrow \\
H \bar{\otimes} H^{\hat{*}}
\end{array}
\end{array}$$

(iii)

$$\begin{array}{c}
 \begin{array}{c}
 H^{\hat{*}} \# \eta_H \quad H^{\hat{*}} \# \eta_H \\
 \downarrow \quad \downarrow \\
 \textcircled{\rho} \quad \textcircled{\lambda} \\
 \downarrow \\
 H \bar{\otimes} H^{\hat{*}}
 \end{array}
 =
 \begin{array}{c}
 \begin{array}{cccc}
 H^{\hat{*}} & \eta_H & H & \eta_{H^{\hat{*}}}
 \end{array} \\
 \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
 \textcircled{-} \quad \textcircled{\rho} \\
 \downarrow \quad \downarrow \\
 \textcircled{\lambda} \\
 \downarrow \\
 H \bar{\otimes} H^{\hat{*}}
 \end{array}
 .
 \end{array}$$

$$\begin{array}{c}
 \text{In fact, the right side} =
 \begin{array}{c}
 \begin{array}{cccc}
 H^{\hat{*}} & \eta_H & H & \eta_{H^{\hat{*}}}
 \end{array} \\
 \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
 \text{Diagram with crossings and caps} \\
 \downarrow \quad \downarrow \\
 H \quad H^{\hat{*}}
 \end{array}$$

$$=
 \begin{array}{c}
 \begin{array}{cc}
 H^{\hat{*}} & H \\
 \downarrow & \downarrow \\
 \text{Diagram with crossings and caps} \\
 \downarrow \quad \downarrow \\
 H & H^{\hat{*}}
 \end{array}
 = \text{the left side.}
 \end{array}$$

Thus (iii) holds.

(iv)

$$\begin{array}{c}
 H^{\hat{*}} \quad \eta_{H^{\hat{*}}} \quad H^{\hat{*}} \quad H \\
 \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
 \text{Diagram 1: A braiding diagram with four strands. The top two strands are labeled } H^{\hat{*}} \text{ and } \eta_{H^{\hat{*}}}. \text{ The bottom two strands are labeled } H^{\hat{*}} \text{ and } H. \text{ The diagram consists of a series of crossings and a box labeled } S \text{ on the top two strands.} \\
 \\
 H \# \eta_{H^{\hat{*}}} \quad H^{\hat{*}} \# \eta_H \\
 \downarrow \quad \downarrow \\
 \text{Diagram 2: A braiding diagram with four strands. The top two strands are labeled } H \# \eta_{H^{\hat{*}}} \text{ and } H^{\hat{*}} \# \eta_H. \text{ The bottom two strands are labeled } H \bar{\otimes} H^{\hat{*}}. \text{ The diagram consists of a series of crossings and a box labeled } S \text{ on the top two strands.} \\
 \\
 H \bar{\otimes} H^{\hat{*}}
 \end{array}$$

$$\begin{array}{c}
 H^{\hat{*}} \quad \eta_{H^{\hat{*}}} \quad H^{\hat{*}} \quad \eta_H \\
 \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
 \text{Diagram 3: A braiding diagram with four strands. The top two strands are labeled } H^{\hat{*}} \text{ and } \eta_{H^{\hat{*}}}. \text{ The bottom two strands are labeled } H^{\hat{*}} \text{ and } \eta_H. \text{ The diagram consists of a series of crossings and a box labeled } S \text{ on the top two strands.} \\
 \\
 \text{The right side} \quad \text{by (iii)} \quad = \\
 \text{Diagram 4: A braiding diagram with four strands. The top two strands are labeled } H^{\hat{*}} \text{ and } \eta_{H^{\hat{*}}}. \text{ The bottom two strands are labeled } H^{\hat{*}} \text{ and } \eta_H. \text{ The diagram consists of a series of crossings and a box labeled } S \text{ on the top two strands.} \\
 \\
 H \quad H^{\hat{*}}
 \end{array}$$

$$\begin{array}{c}
\begin{array}{c} H \quad H^* \\ \text{[Diagram with a box labeled } S \text{]} \\ H \quad H^* \end{array} \\
= \\
\begin{array}{c} H \quad H^* \\ \text{[Diagram with a box labeled } S \text{]} \\ H \quad H^* \end{array}
\end{array}$$

and

$$\begin{array}{c}
\begin{array}{c} H \quad \eta_{H^*} \quad H^* \quad \eta_H \\ \text{[Diagram with crossings]} \\ H \quad H^* \end{array} \\
\text{the left side} = \\
\begin{array}{c} H \quad H^* \\ \text{[Diagram with a box labeled } S \text{]} \\ H \quad H^* \end{array}
\end{array}$$

Thus (iv) holds.

(v)

$$\begin{array}{c}
\begin{array}{c} \eta_H \# H^* \quad \eta_{H^*} \# H \\ \text{[Diagram with boxes } \lambda \text{ and } \rho \text{]} \\ H \otimes H^* \end{array} \\
= \\
\begin{array}{c} \eta_H \quad H^* \quad \eta_{H^*} \quad H \\ \text{[Diagram with crossings and boxes } \lambda \text{ and } \rho \text{]} \\ H \otimes H^* \end{array}
\end{array}$$

In fact, the right side =

H $H^{\hat{*}}$

and

the left side =

H $H^{\hat{*}}$ H $H^{\hat{*}}$


=


H $H^{\hat{*}}$ H $H^{\hat{*}}$


= the right side.

Thus (v) holds.

Now we show that the relation (1) holds.







$$\begin{array}{c}
\begin{array}{cccc}
H & H^{\hat{*}} & H^{\hat{*}} & H
\end{array} \\
\begin{array}{c}
\text{Diagram with four strands, crossings, and nodes } S, \rho, \lambda. \\
\text{The diagram shows a sequence of crossings and nodes. The top node is } S. \text{ Below it are two crossings. Then two more crossings. Then nodes } \rho \text{ and } \lambda. \text{ The strands then cross again and end at } H \bar{\otimes} H^{\hat{*}}.
\end{array}
\end{array}
= \text{the right side of (1)}.$$

(2) We can similarly show relation (2). $\square$

Lemma 8.1.6. *If the antipode of H is invertible, then λ is invertible with inverse*

$$\lambda^{-1} = \begin{array}{c} \begin{array}{cc} H & H^{\hat{*}} \\ \text{Diagram with two strands, a crossing, and node } \bar{S}. \\ \text{The diagram shows two strands, one from } H \text{ and one from } H^{\hat{*}}. \text{ They cross, then the } H^{\hat{*}} \text{ strand goes through node } \bar{S}. \text{ They cross again and end at } H \text{ and } H^{\hat{*}}. \end{array} \end{array} .$$

Lemma 8.1.7. *$R\#H$ becomes an $H^{\hat{*}}$ -module algebra under the module operation $\rightharpoonup' = (id_{H^{\hat{*}}} \otimes \rightharpoonup)(C_{H^{\hat{*}}, R} \otimes id_H)$.*

Theorem 8.1.8. *Let H be a Hopf algebra living in a symmetric tensor category $\mathcal{C}$. If H has a left duality H^* and the antipode S of H is invertible, then*

$$(R\#H)\#H^{\hat{*}} \cong R \otimes (H \bar{\otimes} H^{\hat{*}}) \quad \text{as algebras}$$

where $H \bar{\otimes} H^{\hat{*}}$ is defined in Lemma 8.1.3.

(i)

Let

and

where $b' = C_{H, H^*} b_H$.

We define

and

We see that

since w is algebraic

$$\begin{array}{c}
\begin{array}{c} R \quad H\#H^{\hat{*}} \\ \text{Diagram 1: A box with } S \text{ and } \alpha \text{ connected by } w. \text{ Inputs } R, H\#H^{\hat{*}} \text{ from bottom.} \end{array} \\
= \\
\begin{array}{c} R \quad H\#H^{\hat{*}} \\ \text{Diagram 2: Similar to Diagram 1, but with } \alpha \text{ and } w \text{ in different positions.} \end{array} \\
= \\
\begin{array}{c} R \quad H\#H^{\hat{*}} \\ \text{Diagram 3: Two separate vertical lines.} \end{array}
\end{array}$$

Similarly, we have $\Phi\Psi = id$. Thus Φ is invertible.

Now we show that Φ is algebraic.

$$\begin{array}{c}
\text{Let} \\
\begin{array}{c} (R\#H)\#H^{\hat{*}} \\ \downarrow \\ \Phi' \\ \downarrow \\ R \otimes (H\bar{\otimes}H^{\hat{*}}) \end{array} = \begin{array}{c} R \quad \eta_H \quad H\#H^{\hat{*}} \\ \text{Diagram: A complex network of boxes } S, \alpha, \lambda, \rho \text{ and lines.} \end{array}
\end{array}$$

It is clear that $\Phi = (id \otimes \lambda^{-1})\Phi'$. Consequently, we only need show that Φ' is algebraic.

$$\begin{array}{c}
\text{Let} \\
\begin{array}{c} R \\ \downarrow \\ \xi \\ \downarrow \\ R \otimes (H\bar{\otimes}H^{\hat{*}}) \end{array} = \begin{array}{c} R \quad \eta_H \\ \text{Diagram: A complex network of boxes } S, \alpha, \lambda, \rho \text{ and lines.} \end{array}
\end{array}$$


We have that

$$\begin{array}{c}
(R\#H)\#H^{\hat{*}} \\ \downarrow \\ \Phi' \\ \downarrow \\ R \otimes (H\bar{\otimes}H^{\hat{*}}) \end{array} = \begin{array}{c} R \quad H\#H^{\hat{*}} \\ \text{Diagram: A box with } \xi \text{ and } \lambda \text{ connected by a line.} \end{array}$$

We claim that

$$\begin{array}{c}
\begin{array}{c}
H \quad H^* \quad R \\
\downarrow \quad \downarrow \quad \downarrow \\
\lambda \quad \xi \\
\downarrow \quad \downarrow \\
R \quad H \bar{\otimes} H^*
\end{array}
=
\begin{array}{c}
H \quad H^* \quad R \\
\downarrow \quad \downarrow \quad \downarrow \\
\downarrow \quad \downarrow \quad \downarrow \\
\xi \quad \lambda \\
\downarrow \quad \downarrow \\
R \quad H \bar{\otimes} H^*
\end{array}
.
\end{array}
\quad \dots (*)$$

$$\begin{array}{c}
\text{the left side} =
\begin{array}{c}
H \quad H^* \quad R \quad \eta_H \\
\downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
\lambda \quad b' \quad \alpha \\
\downarrow \quad \downarrow \quad \downarrow \\
S \quad \downarrow \quad \downarrow \\
\downarrow \quad \downarrow \quad \downarrow \\
\rho \quad \downarrow \quad \downarrow \\
R \quad H \bar{\otimes} H^*
\end{array}
\quad \text{by Lemma 8.1.5(1)}
=
\begin{array}{c}
H \quad H^* \quad R \quad \eta_H \\
\downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
\downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
S \quad \downarrow \quad \downarrow \quad \downarrow \\
\downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\
\rho \quad \downarrow \quad \downarrow \quad \downarrow \\
R \quad H \bar{\otimes} H^*
\end{array}
\end{array}$$



$$= \text{[Diagram 1]} = \text{[Diagram 2]} = \text{[Diagram 3]} .$$

Thus relation (*) holds.

Next, we check that ξ is algebraic. We see that

$$\text{[Diagram 1]} = \text{[Diagram 2]} = \text{[Diagram 3]} = \text{[Diagram 4]} = \text{[Diagram 5]}$$

and obviously

$$\text{[Diagram 1]} = \text{[Diagram 2]} .$$

Thus ξ is algebraic.

Now we show that Φ' is algebraic.

$$\begin{array}{c}
\begin{array}{c} (R\#H)\#H^{\hat{*}} \quad (R\#H)\#H^{\hat{*}} \\ \downarrow \quad \downarrow \\ \Phi' \quad \Phi' \\ \downarrow \\ R \otimes (H \bar{\otimes} H^{\hat{*}}) \end{array} = \begin{array}{c} \begin{array}{c} R \quad H \quad H^{\hat{*}} \quad R \quad H \quad H^{\hat{*}} \\ \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\ \xi \quad \lambda \quad \xi \quad \lambda \\ \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\ R \quad H \bar{\otimes} H^{\hat{*}} \end{array} \end{array} \\
\\
= \begin{array}{c} \begin{array}{c} R \quad H \quad H^{\hat{*}} \quad R \quad H \quad H^{\hat{*}} \\ \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\ \xi \quad \lambda \quad \xi \quad \lambda \\ \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\ R \quad H \bar{\otimes} H^{\hat{*}} \end{array} \quad \text{by } (*) \quad \begin{array}{c} R \quad H \quad H^{\hat{*}} \quad R \quad H \quad H^{\hat{*}} \\ \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\ \xi \quad \alpha \quad \lambda \quad \xi \\ \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\ R \quad H \bar{\otimes} H^{\hat{*}} \end{array} \\
\\
\text{by Lemma 8.1.4} \quad \begin{array}{c} R \quad H \quad H^{\hat{*}} \quad R \quad H \quad H^{\hat{*}} \\ \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\ \xi \quad \alpha \quad \lambda \quad \xi \\ \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\ R \quad H \bar{\otimes} H^{\hat{*}} \end{array} \quad \text{since } \xi \text{ is algebraic} \quad \begin{array}{c} R \quad H \quad H^{\hat{*}} \quad R \quad H \quad H^{\hat{*}} \\ \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \quad \downarrow \\ \alpha \quad \lambda \quad \xi \\ \downarrow \quad \downarrow \quad \downarrow \\ R \quad H \bar{\otimes} H^{\hat{*}} \end{array} \\
\\
= \begin{array}{c} (R\#H)\#H^{\hat{*}} \quad (R\#H)\#H^{\hat{*}} \\ \downarrow \quad \downarrow \\ \Phi' \\ \downarrow \\ R \otimes (H \bar{\otimes} H^{\hat{*}}) \end{array} .
\end{array}$$

It is clear that

$$\begin{array}{ccc}
 \eta_{(R\#H)\#H^*} & & \eta_{R\otimes(H\bar{\otimes}H^*)} \\
 | & & | \\
 \textcircled{\Phi'} & = & \\
 | & & | \\
 R\otimes(H\bar{\otimes}H^*) & & R\otimes(H\bar{\otimes}H^*)
 \end{array}$$

Thus Φ' is algebraic. $\square$

8.2 The duality theory for Yetter-Drinfeld Hopf algebras

In this section, we give the duality theory for Yetter-Drinfeld Hopf algebra H .

A braided Hopf algebra in Yetter-Drinfeld categories is called a Yetter-Drinfeld Hopf algebra.

In this section, all Yetter-Drinfeld categories are symmetric.

If H is a finite-dimensional Hopf algebra living in a symmetric tensor category $\mathcal{C}$ determined by (co)triangular structure, then $H\bar{\otimes}H^* = \text{End}_k(H)$ since $\dim(H\bar{\otimes}H^*) = \dim(\text{End}_k H)$ and $H\bar{\otimes}H^*$ is a subalgebra of $\text{End}_k H$. We can easily check that

$$R\otimes \text{End}_k H \cong M_n(R) \quad \text{as algebras}$$

where $M_n(R) = \{x \mid x \text{ is a } n \times n\text{-matrix over } k\}$

Considering Theorem 8.1.8, we have :

Theorem 8.2.1. (*Duality theorem*) *Let H be a finite-dimensional Yetter-Drinfeld Hopf algebra Then*

$$(R\#H)\#H^* \cong M_n(R) \quad (\text{as algebras}).$$

Some interesting braided group crossed products are the Weyl algebras, which have been studied, for example in [77]. Now we give an example in super case.

Example 8.2.2. (see [83, P502]) *Let R denote the Gassmann plane $C^{0|0}$. That is, $R = \mathbf{C}(\theta)$ with odd coordinate θ and relation $\theta^2 = 0$. The comultiplication and antipode are*

$$\Delta(\theta) = \theta \otimes 1 + 1 \otimes \theta \quad \text{and} \quad S(\theta) = -\theta.$$

R is a super-Hopf algebra with $\dim R = 2$. Let $H = R$ and α be the adjoint action. Thus (R, α) is a left H -module algebra in super case. Since H is finite-dimensional we have that

$$(R\#H)\#H^* \cong M_n(R) \quad (\text{as superalgebras}).$$

by Theorem 8.2.1. $\square$

In fact, we can obtain duality theory for crossed coproducts by only turning the above diagrams upside down.

Part III

Hopf Algebras Living in the Category of Vector spaces

Chapter 9

The Relation Between Decomposition of Comodules and Coalgebras

The decomposition of coalgebras and comodules is an important subject in study of Hopf algebras. T. Shudo and H. Miyamoto [112] showed that C can be decomposed into a direct sum of its indecomposable subcoalgebras of C . Y.H. Xu [121] showed that the decomposition was unique. He also showed that M can uniquely be decomposed into a direct sum of the weak-closed indecomposable subcomodules of M (we call the decomposition the weak-closed indecomposable decomposition) in [124]. In this chapter, we give the relation between the two decomposition. We show that if M is a full, W -relational hereditary C -comodule, then the following conclusions hold:

- (1) M is indecomposable iff C is indecomposable;
- (2) M is relative-irreducible iff C is irreducible;
- (3) M can be decomposed into a direct sum of its weak-closed relative-irreducible subcomodules iff C can be decomposed into a direct sum of its irreducible subcoalgebras.

We also obtain the relation between coradical of C -comodule M and radical of algebra $C(M)^*$

Let k be a field, M be a C -comodule, N be a subcomodule of M , E be a subcoalgebras of C and P be an ideal of C^* . As in [124], we define:

$$E^{\perp C^*} = \{f \in C^* \mid f(E) = 0\}.$$

$$P^{\perp C} = \{c \in C \mid P(c) = 0\}.$$

$$N^{\perp C^*} = \{f \in C^* \mid f \cdot N = 0\}.$$

$$P^{\perp M} = \{x \in M \mid P \cdot x = 0\}.$$

Let $\langle N \rangle$ denote $N^{\perp C^* \perp M}$. $\langle N \rangle$ is called the closure of N . If $\langle N \rangle = N$, then N is called closed. If $N = C^*x$, then $\langle N \rangle$ is denoted by $\langle x \rangle$. If $\langle x \rangle \subseteq N$ for any $x \in N$, then N is called weak-closed. It is clear that any closed subcomodule is weak-closed. If $\rho(N) \subseteq N \otimes E$, then N is called an E -subcomodule of M . Let

$$M_E = \sum \{N \mid N \text{ is a subcomodule of } M \text{ and } \rho(N) \subseteq N \otimes E\}.$$

We call M_E a component of M over E . If M_E is some component of M and $M_F \subseteq M_E$ always implies $M_F = M_E$ for any non-zero component M_F , then M_E is called a minimal component of M .

Let $\{m_\lambda \mid \lambda \in \Lambda\}$ be the basis of M and $C(M)$ denote the subspace of C spanned by

$W(M) = \{c \in C \mid \text{there exists an } m \in M \text{ with } \rho(m) = \sum m_\lambda \otimes c_\lambda \text{ such that } c_{\lambda_0} = c \text{ for some } \lambda_0 \in \Lambda\}$.

E. Abe in [1, P129] checked that $C(M)$ is a subcoalgebra of C . It is easy to know that if E is subcoalgebra of C and $\rho(M) \subseteq M \otimes E$, then $C(M) \subseteq E$. If $C(M) = C$, then M is called a full C -comodule. If D is a simple subcoalgebra of C and $M_D \neq 0$ or $D = 0$, then D is called faithful to M . If every simple subcoalgebra of C is faithful to M , then M is called a component faithfulness C -comodule.

Let X and Y be subspaces of coalgebra C . Define $X \wedge Y$ to be the kernel of the composite

$$C \xrightarrow{\Delta} C \otimes C \rightarrow C/X \otimes C/Y.$$

$X \wedge Y$ is called a wedge of X and Y .

9.1 The relation between the decomposition of comodules and coalgebras

Lemma 9.1.1. *Let N and L be subcomodules of M . Let D and E be subcoalgebras of C . Then*

- (1) $N^{\perp C^*} = (C(N))^{\perp C^*}$;
- (2) $M_E = M_{C(M_E)}$;
- (3) N is closed iff there exists a subcoalgebra E such that $N = M_E$;
- (4) $M_D \neq 0$ iff $D^{\perp C^* \perp M} \neq 0$;
- (5) If $D \cap C(M) = 0$, then $M_D = 0$;
- (6) If D is a simple subcoalgebra of C , then $C(M_D) = \begin{cases} D & \text{if } M_D \neq 0 \\ 0 & \text{if } M_D = 0 \end{cases}$;
- (7) If $D \cap E = 0$, then $M_D \cap M_E = 0$;
- (8) If D and E are simple subcoalgebras and $M_D \cap M_E = 0$ with $M_D \neq 0$ or $M_E \neq 0$, then $D \cap E = 0$.
- (9) If M_E is the minimal component of M and $0 \neq N \subseteq M_E$, then $C(M_E) = C(N)$.

Proof. (1) Let $\{m_\lambda \mid \lambda \in \Lambda\}$ be a basis of N . For any $f \in N^{\perp C^*}$, if $c \in W(N)$, then there exists $m \in N$ with $\rho(m) = \sum_{\lambda \in \Lambda} m_\lambda \otimes c_\lambda$ such that $c_{\lambda_0} = c$ for some $\lambda_0 \in \Lambda$. Since $f \cdot m = \sum m_\lambda f(c_\lambda) = 0$, $f(c_\lambda) = 0$ for any $\lambda \in \Lambda$. Obviously, $f(c) = f(c_{\lambda_0}) = 0$. Considering that $C(N)$ is the space spanned by $W(N)$, we have $f \in C(N)^{\perp C^*}$. Conversely, if $f \in C(N)^{\perp C^*}$, then $f \cdot m = \sum m_\lambda f(c_\lambda) = 0$ for any $m \in N$, i.e. $f \in N^{\perp C^*}$. This shows that $N^{\perp C^*} = C(N)^{\perp C^*}$.

(2) Since M_E is an E -subcomodule of M , $C(M_E) \subseteq E$ and $M_{C(M_E)} \subseteq M_E$. Since M_E is a $C(M_E)$ -subcomodule, $M_E \subseteq M_{C(M_E)}$.

(3) If N is closed, then $N^{\perp C^* \perp M} = N$. Let $E = C(N)$. Obviously, $N \subseteq M_E$. By Lemma 9.1.1(1), $N^{\perp C^*} \cdot M_E = E^{\perp C^*} \cdot M_E = 0$. Thus $M_E \subseteq N^{\perp C^* \perp M}$ and $M_E \subseteq N$. This shows that

$M_E = N$. Conversely, if $M_E = N$, obviously, $(M_E) \subseteq (M_E)^{\perp C^* \perp M}$. Thus it is sufficient to show that

$$\langle M_E \rangle \subseteq M_E.$$

Let $L = (M_E)^{\perp C^* \perp M} = \langle M_E \rangle$. We see that $C(L)^{\perp C^*} = L^{\perp C^*} = (M_E)^{\perp C^* \perp M \perp C^*} = (M_E)^{\perp C^*} = C(M_E)^{\perp C^*}$. Thus $C(L) = C(M_E)$ and $L \subseteq M_{C(L)} = M_{C(M_E)} = M_E$.

(4) If $D^{\perp C^* \perp M} \neq 0$, let $\{m_\lambda \mid \lambda \in \Lambda\}$ be a basis of $D^{\perp C^* \perp M}$ and $\rho(x) = \sum m_\lambda \otimes d_\lambda$ for any $x \in D^{\perp C^* \perp M}$. Since $D^{\perp C^*} \cdot x = \sum_\lambda m_\lambda D^{\perp C^*}(d_\lambda) = 0$, $d_\lambda \in D^{\perp C^* \perp C} = D$ for any $\lambda \in \Lambda$, which implies that $D^{\perp C^* \perp M}$ is a D -subcomodule. Therefore $0 \neq D^{\perp C^* \perp M} \subseteq M_D$, i.e. $M_D \neq 0$. Conversely, if $M_D \neq 0$, then we have that

$$\begin{aligned} 0 \neq M_D &\subseteq (M_D)^{\perp C^* \perp M} \\ &= C(M_D)^{\perp C^* \perp M} \quad (\text{by part (1)}) \\ &\subseteq D^{\perp C^* \perp M}. \end{aligned}$$

(5) Since $M_D \subseteq M$, $C(M_D) \subseteq C(M)$. Obviously, $C(M_D) \subseteq D$. Thus $C(M_D) \subseteq C(M) \cap D = 0$, which implies that $M_D = 0$.

(6) If $M_D = 0$, then $C(M_D) = 0$. If $M_D \neq 0$, then $0 \neq C(M_D) \subseteq D$. Since D is a simple subcoalgebra, $C(M_D) = D$.

(7) If $x \in M_D \cap M_E$, then $\rho(x) \in (M_D \otimes D) \cap (M_E \otimes E)$, and

$$\rho(x) = \sum_{i=1}^n x_i \otimes d_i = \sum_{j=1}^m y_j \otimes e_j, \quad (9.1)$$

where $x_i, \dots, x_n$ is linearly independent and $d_i \in D$ and $e_j \in E$. Let $f_l \in M^*$ with $f_l(x_i) = \delta_{li}$ for $i, l = 1, 2, \dots, n$. Let $f_l \otimes id$ act on equation (9.1). We have that $d_l = \sum_{j=1}^m f_l(y_j) e_j \in E$, which implies $d_l \in D \cap E = 0$ and $d_l = 0$ for $l = 1, \dots, n$. Therefore $M_D \cap M_E = 0$.

(8) If $D \cap E \neq 0$, then $D = E$. Thus $M_D = M_E$ and $M_D \cap M_E = M_E = M_D = 0$. We get a contradiction. Therefore $D \cap E = 0$.

(9) Obviously, $C(N) \subseteq C(M_E)$. Conversely, since $0 \neq N \subseteq M_{C(N)} \subseteq M_{C(M_E)} = M_E$ and M_E is a minimal component, $M_{C(N)} = M_E$. By the definition of component, $C(M_E) \subseteq C(N)$. Thus $C(N) = C(M_E)$. $\square$

Proposition 9.1.2. *If E is a subcoalgebra of C , then the following conditions are equivalent.*

- (1) M_E is a minimal component of M .
- (2) $C(M_E)$ is a simple subcoalgebra of C .
- (3) M_E is a minimal closed subcomodule of M .

Proof. It is easy to check that $M_E = 0$ iff $C(M_E) = 0$. Thus (1), (2) and (3) are equivalent when $C(M_E) = 0$. We now assume that $M_E \neq 0$.

(1) $\implies$ (2) Since $0 \neq M_E$, there exists a non-zero finite dimensional simple subcomodule N of M such that $N \subseteq M_E$. By [124, Lemma 1.1], N is a simple C^* -submodule of M . Since M_E is a minimal component, $C(N) = C(M_E)$ by Lemma 9.1.1 (9). Let $D = C(N) = C(M_E)$. By

Lemma 9.1.1(1), $(0 : N)_{C^*} = N^{\perp C^*} = C(N)^{\perp C^*} = D^{\perp C^*}$. Thus N is a faithful simple $C^*/D^{\perp C^*}$ -module, and so $C^*/D^{\perp C^*}$ is a simple algebra. It is clear that D is a simple subcoalgebra of C .

(2) $\implies$ (3) If $0 \neq N \subseteq M_E$ and N is a closed subcomodule of M , then by Lemma 9.1.1(3) there exists a subcoalgebra F of C such that $N = M_F$. Since $0 \neq C(N) = C(M_F) \subseteq C(M_E)$ and $C(M_E)$ is simple, $C(M_F) = C(M_E)$. By Lemma 9.1.1(2), $N = M_F = M_{C(M_F)} = M_{C(M_E)} = M_E$, which implies that M_E is a minimal closed subcomodule.

(3) $\implies$ (1), If $0 \neq M_F \subseteq M_E$, then M_F is a closed subcomodule by Lemma 9.1.1(3) and $M_E = M_F$, i.e. M_E is a minimal component.

This completes the proof. $\square$

Let

$\mathcal{C}_0 = \{D \mid D \text{ is a simple subcoalgebra of } C\};$

$\mathcal{C}(M)_0 = \{D \mid D \text{ is a simple subcoalgebra of } C \text{ and } D \subseteq C(M)\};$

$\mathcal{M}_0 = \{N \mid N \text{ is a minimal closed subcomodule of } M\};$

$\mathcal{C}(M)_1 = \{D \mid D \text{ is a faithful simple subcoalgebra of } C \text{ to } M\};$

$\mathcal{C}_0 = \sum \{D \mid D \text{ is a simple subcoalgebra of } C\};$

$\mathcal{M}_0 = \sum \{N \mid N \text{ is a minimal closed subcomodule of } M\};$

$\mathcal{C}_0$ and $\mathcal{M}_0$ are called the coradical of coalgebra C and the coradical of comodule M respectively. If $\mathcal{M}_0 = M$, then M is called cosemisimple. By Lemma 9.1.1(5),

$$\mathcal{C}(M)_1 \subseteq \mathcal{C}(M)_0$$

Theorem 9.1.3. $\psi \begin{cases} \mathcal{C}(M)_1 \longrightarrow \mathcal{M}_0 \\ D \longmapsto M_D \end{cases} \text{ is bijective.}$

Proof. By Proposition 9.1.2, ψ is a map. Let D and $E \in \mathcal{C}(M)_1$ and $\psi(D) = \psi(E)$, i.e. $M_D = M_E$. By Lemma 9.1.1 (6), we have that $D = C(D_D) = C(M_E) = E$. If $N \in \mathcal{M}_0$, then $N = M_{C(N)}$ and $C(N)$ is a simple subcoalgebra by Lemma 9.1.1(3) and Proposition 9.1.2. Thus $\psi(C(N)) = N$, which implies that ψ is surjective. $\square$

In [124] and [121], Xu defined the equivalence relation for coalgebra and for comodule as follows:

Definition 9.1.4. We say that $D \sim E$ for D and $E \in \mathcal{C}_0$ iff for any pair of subclasses $\mathcal{C}_1$ and $\mathcal{C}_2$ of $\mathcal{C}_0$ with $D \in \mathcal{C}_1$ and $E \in \mathcal{C}_2$ such that $\mathcal{C}_1 \cup \mathcal{C}_2 = \mathcal{C}_0$ and $\mathcal{C}_1 \cap \mathcal{C}_2 = \emptyset$, there exist elements $D_1 \in \mathcal{C}_1$ and $E_1 \in \mathcal{C}_2$ such that $D_1 \wedge E_1 \neq E_1 \wedge D_1$. Let $[D]$ denote the equivalence class which contains D .

We say that $N \sim L$ for N and $L \in \mathcal{M}_0$ iff for any pair of subclasses $\mathcal{M}_1$ and $\mathcal{M}_2$ of $\mathcal{M}_0$ with $N \in \mathcal{M}_1$ and $L \in \mathcal{M}_2$ such that $\mathcal{M}_1 \cup \mathcal{M}_2 = \mathcal{M}_0$ and $\mathcal{M}_1 \cap \mathcal{M}_2 = \emptyset$, there exist elements $N_1 \in \mathcal{M}_1$ and $L_1 \in \mathcal{M}_2$ such that $N_1 \wedge L_1 \neq L_1 \wedge N_1$. Let $[N]$ denote the equivalence class which contains N .

Definition 9.1.5. If $D \wedge E = E \wedge D$ for any simple subcoalgebras D and E of C , then C is called π -commutative. If $N \wedge L = L \wedge N$ for any minimal closed subcomodules N and L of M , then M is called π -commutative.

Obviously, every cocommutative coalgebra is π -commutative. By [124, Theorem 3.8 and Theorem 4.18], M is π -commutative iff M can be decomposed into a direct sum of the weak-closed relative-irreducible subcomodules of M iff every equivalence class of M contains only one element. By [121], C is π -commutative iff C can be decomposed into a direct sum of irreducible subcoalgebras of C iff equivalence every class of C contains only one element.

Lemma 9.1.6. *Let D , E and F be subcoalgebras of C . N , L and T be subcomodules of M . Then*

- (1) $M_D \wedge M_F = M_{C(M_D) \wedge C(M_F)} \subseteq M_{D \wedge F}$;
- (2) If D and E are faithful simple subcoalgebras of C to M , then $M_D \wedge M_E = M_{D \wedge E}$;
- (3) $M_{D+E} \supseteq M_D + M_E$;
- (4) If $F = \sum \{ D_\alpha \mid \alpha \in \Omega \}$ and $\{ D_\alpha \mid \alpha \in \Omega \} \subseteq \mathcal{C}_0$, then $M_F = \sum \{ M_{D_\alpha} \mid \alpha \in \Omega \}$. In particular, $M_{C_0} = M_0$.
- (5) $(N + L) \wedge T \supseteq N \wedge T + L \wedge T$;
- (6) $(D + E) \wedge F \supseteq D \wedge F + E \wedge F$.

Proof. (1) We see that

$$\begin{aligned} M_D \wedge M_E &= \rho^{-1}(M \otimes (M_D)^{\perp C^* \perp C} \wedge (M_E)^{\perp C^* \perp C}) \quad (\text{by [124, Proposition 2.2(1)]}) \\ &= \rho^{-1}(M \otimes C(M_D) \wedge C(M_E)) \quad (\text{by Lemma 9.1.1(1)}). \end{aligned}$$

By the definition of component, subcomodule $M_D \wedge M_E \subseteq M_{C(M_D) \wedge C(M_E)}$. It follows from the equation above that $M_{C(M_D) \wedge C(M_E)} \subseteq M_D \wedge M_E$. Thus

$$M_D \wedge M_E = M_{C(M_D) \wedge C(M_E)}$$

and

$$M_{C(M_D) \wedge C(M_E)} \subseteq M_{D \wedge E}.$$

(2) Since D and E are faithful simple subcoalgebras of C to M , $C(M_D) = D$ and $C(M_E) = E$ by Lemma 9.1.1(6). By Lemma 9.1.6(1), $M_D \wedge M_E = M_{D \wedge E}$.

(3) It is trivial.

(4) Obviously $M_F \supseteq \sum \{ M_{D_\alpha} \mid \alpha \in \Omega \}$. Conversely, let $N = M_F$. Obviously N is an F -comodule. For any $x \in N$, let $L = C^*x$. it is clear that L is a finite dimensional comodule over F . By [114, Lemma 14.0.1], L is a completely reducible module over F^* . Thus L can be decomposed into a direct sum of simple F^* -submodules:

$$L = L_1 \oplus L_2 \oplus \cdots \oplus L_n,$$

where L_i is a simple F^* -submodule. By [124, Proposition 1.16], $\langle L_i \rangle = (L_i)^{\perp F^* \perp N}$ is a minimal closed F -subcomodule of N . By Theorem 9.1.3, there exists a simple subcoalgebra D_{α_i} of F such that $\langle L_i \rangle = N_{D_{\alpha_i}}$. Obviously, $N_{D_{\alpha_i}} \subseteq M_{D_{\alpha_i}}$. Thus $L \subseteq \sum_{i=1}^n \langle L_i \rangle = \sum_{i=1}^n N_{D_{\alpha_i}} \subseteq \sum \{ M_{D_\alpha} \mid \alpha \in \Omega \}$. Therefore $M_F = \sum \{ M_{D_\alpha} \mid \alpha \in \Omega \}$. If $C_0 = F = \sum \{ D \mid D \in \mathcal{C}_0 \}$, then $M_{C_0} = M_F = \sum \{ M_D \mid D \in \mathcal{C}_0 \} = M_0$ by Theorem 9.1.3,

(5) and (6) are trivial $\square$

Lemma 9.1.7. *Let N be a subcomodule of M , and let D , E and F be simple subcoalgebras of C . Then*

(1) $D \sim 0$ iff $D = 0$; $M_D \sim 0$ iff $M_D = 0$;

We called the equivalence class which contains zero a zero equivalence class.

(2) If D and E are faithful to M and $M_D \sim M_E$, then $D \sim E$;

(3) $[M_D] \subseteq \{M_E \mid E \in [D]\}$;

(4) If D is faithful to M , then $[M_D] \subseteq \{M_E \mid E \in [D] \text{ and } E \text{ is faithful to } M\}$.

Proof. (1) If $D \sim 0$ and $D \neq 0$, let $\mathcal{C}_1 = \{0\}$ and $\mathcal{C}_2 = \{F \mid F \neq 0, F \in \mathcal{C}_0\}$. Thus $\mathcal{C}_1 \cup \mathcal{C}_2 = \mathcal{C}_0$ and $\mathcal{C}_1 \cap \mathcal{C}_2 = \emptyset$ with $0 \in \mathcal{C}_1$ and $D \in \mathcal{C}_2$. But for any $D_1 \in \mathcal{C}_1$ and $E_1 \in \mathcal{C}_2$, since $D_1 = 0$, $D_1 \wedge E_1 = E_1 \wedge D_1 = E_1$. We get a contradiction. Thus $D = 0$. Conversely, if $D = 0$, obviously $D \sim 0$. Similarly, we can show that $M_D \sim 0$ iff $M_D = 0$.

(2) For any pair of subclasses $\mathcal{C}_1$ and $\mathcal{C}_2$ of $\mathcal{C}_0$, if $\mathcal{C}_1 \cap \mathcal{C}_2 = \emptyset$ and $\mathcal{C}_1 \cup \mathcal{C}_2 = \mathcal{C}_0$ with $D \in \mathcal{C}_1$ and $E \in \mathcal{C}_2$, let $\mathcal{M}_1 = \{M_F \mid F \in \mathcal{C}_1 \text{ and } F \text{ is faithful to } M\}$ and $\mathcal{M}_2 = \{M_F \mid F \in \mathcal{C}_2 \text{ and } F \text{ is faithful to } M\}$. By Theorem 9.1.3, $\mathcal{M}_0 = \mathcal{M}_1 \cup \mathcal{M}_2$ and $\mathcal{M}_1 \cap \mathcal{M}_2 = \emptyset$. Obviously, $M_D \in \mathcal{M}_1$ and $M_E \in \mathcal{M}_2$. Since $M_D \sim M_E$, there exist $M_{D_1} \in \mathcal{M}_1$ and $M_{E_1} \in \mathcal{M}_2$ such that $M_{D_1} \wedge M_{E_1} \neq M_{E_1} \wedge M_{D_1}$, where $D_1 \in \mathcal{C}_1$ and $E_1 \in \mathcal{C}_2$. By Lemma 9.1.6(2), $M_{D_1 \wedge E_1} \neq M_{E_1 \wedge D_1}$. Thus $D_1 \wedge E_1 \neq E_1 \wedge D_1$. Obviously D and $D_1 \in \mathcal{C}_1$. Meantime E and $E_1 \in \mathcal{C}_2$. By Definition 9.1.4, $D \sim E$.

(3) Obviously, $M_F \sim M_D$ for any $M_F \in [M_D]$. If $M_D \neq 0$, then $M_F \neq 0$ by Lemma 9.1.7(1). By Lemma 9.1.7(2), $F \sim D$. Thus $M_F \in \{M_E \mid E \in [D]\}$. If $M_D = 0$, by Lemma 9.1.7(1), $M_F = 0 = M_D \in \{M_E \mid E \in [D]\}$.

(4) If $M_F \in [M_D]$, then $M_D \sim M_F$. If $D = 0$, then $M_D = 0$. By Lemma 9.1.7(1), $M_F = 0$. Thus $M_F = M_D = 0 \in \{M_E \mid E \in [D] \text{ and } E \text{ is faithful to } M\}$. If $D \neq 0$, we have that $M_D \neq 0$ since D is faithful to M . By Lemma 9.1.7(1), $M_F \neq 0$. By Lemma 9.1.7(3), $M_F \in \{M_E \mid E \in [D] \text{ and } E \text{ is faithful to } M\}$. $\square$

Theorem 9.1.8. *Let $\{\mathcal{E}_\alpha \mid \alpha \in \bar{\Omega}\}$ be all of the equivalence classes of C . and $E_\alpha = \sum\{E \mid E \in \mathcal{E}_\alpha\}$. Then*

(1) *For any $\alpha \in \bar{\Omega}$, there exists a set I_α and subclasses $\mathcal{E}(\alpha, i) \subseteq \mathcal{E}_\alpha$ such that $\cup\{\mathcal{E}(\alpha, i) \mid i \in I_\alpha\} = \mathcal{E}_\alpha$ and $\{M_{\mathcal{E}(\alpha, i)} \mid \alpha \in \bar{\Omega}, i \in I_\alpha\}$ is the set of the equivalence classes of M (they are distinct except for zero equivalence class), where $M_{\mathcal{E}(\alpha, i)}$ denotes $\{M_D \mid D \in \mathcal{E}(\alpha, i)\}$.*

(2) *If M is a component faithfulness C -comodule, then $\{M_{\mathcal{E}(\alpha, i)} \mid \alpha \in \bar{\Omega}, i \in I_\alpha\}$ is the set of the distinct equivalence classes of M .*

(3)

$$M = \sum_{\alpha \in \bar{\Omega}} \oplus M_{(E_\alpha)^{(\infty)}} = \sum_{\alpha \in \bar{\Omega}} \sum_{i \in I_\alpha} \oplus (M_{\mathcal{E}(\alpha, i)})^{(\infty)} = \sum_{\alpha \in \bar{\Omega}} \oplus (M_{E_\alpha})^{(\infty)}$$

and for any $\alpha \in \bar{\Omega}$,

$$M_{(E_\alpha)^{(\infty)}} = \sum_{i \in I_\alpha} (M_{\mathcal{E}(\alpha, i)})^{(\infty)} = (M_{E_\alpha})^{(\infty)}$$

where $E(\alpha, i) = \sum\{E \mid E \in \mathcal{E}(\alpha, i)\}$.

Proof. (1) By Theorem 9.1.3 and Lemma 9.1.7(3) we can immediately get part (1).

(2) If M is a component faithfulness C -comodule, then $\mathcal{C}(M)_1 = \mathcal{C}(M)_0 = \mathcal{C}_0$. It follows from Theorem 9.1.3 and part (1) that $\{M_{\mathcal{E}(\alpha,i)} \mid \alpha \in \bar{\Omega}, i \in I_\alpha\}$ consists of all the distinct equivalence classes of M .

(3) We see that

$$\begin{aligned}
M &= M_C = M_{\sum\{(E_\alpha)^{(\infty)} \mid \alpha \in \bar{\Omega}\}} \text{ (by [121])} \\
&\supseteq \sum_{\alpha \in \bar{\Omega}} M_{(E_\alpha)^{(\infty)}} \text{ (by Lemma 9.1.6(3))} \\
&\supseteq \sum_{\alpha \in \bar{\Omega}} \sum_{n=0}^{\infty} M_{\wedge^{n+1} E_\alpha} \text{ (by Lemma 9.1.6(3))} \\
&\supseteq \sum_{\alpha \in \bar{\Omega}} \sum_{n=0}^{\infty} \wedge^{n+1} M_{E_\alpha} \text{ (by Lemma 9.1.6(1))} \\
&= \sum_{\alpha \in \bar{\Omega}} \sum_{n=0}^{\infty} \wedge^{n+1} M_{\sum_{i \in I_\alpha} E(\alpha,i)} \text{ (by Theorem 9.1.8(1))} \\
&\supseteq \sum_{\alpha \in \bar{\Omega}} \sum_{n=0}^{\infty} \wedge^{n+1} \sum_{i \in I_\alpha} M_{E(\alpha,i)} \text{ (by Lemma 9.1.6(3))} \\
&\supseteq \sum_{\alpha \in \bar{\Omega}} \sum_{n=0}^{\infty} \sum_{i \in I_\alpha} \wedge^{n+1} M_{E(\alpha,i)} \text{ (by Lemma 9.1.6(5)(6))} \\
&= \sum_{\alpha \in \bar{\Omega}} \sum_{i \in I_\alpha} \sum_{n=0}^{\infty} \wedge^{n+1} M_{E(\alpha,i)} \\
&= \sum_{\alpha \in \bar{\Omega}} \sum_{i \in I_\alpha} (M_{E(\alpha,i)})^{(\infty)} \\
&= M \text{ (by [124, (4.10) in Theorem 4.15] and Lemma 9.1.6(4) and part (1)).}
\end{aligned}$$

Thus

$$M = \sum_{\alpha \in \bar{\Omega}} M_{(E_\alpha)^{(\infty)}} = \sum_{\alpha \in \bar{\Omega}} \sum_{i \in I_\alpha} (M_{E(\alpha,i)})^{(\infty)} \quad (9.2)$$

and

$$M = \sum_{\alpha \in \bar{\Omega}} \sum_{i \in I_\alpha} \oplus (M_{E(\alpha,i)})^{(\infty)}$$

by [124, Theorem 4.15] and part (1). We see that

$$\begin{aligned}
M_{E(\alpha,i)} \wedge M_{E(\alpha,i)} &\subseteq M_{(E_\alpha)^{(\infty)}} \wedge M_{(E_\alpha)^{(\infty)}} \\
&\subseteq M_{(E_\alpha)^{(\infty)} \wedge (E_\alpha)^{(\infty)}} \text{ (by Lemma 9.1.6(1))} \\
&= M_{(E_\alpha)^{(\infty)}} \text{ (by [54, Proposition 2.1.1])},
\end{aligned}$$

Thus

$$M_{(E_\alpha)^{(\infty)}} \supseteq (M_{E(\alpha,i)})^{(\infty)}$$

for any $i \in I_\alpha$ and for any $\alpha \in \bar{\Omega}$, and

$$M_{(E_\alpha)^{(\infty)}} \supseteq \sum_{i \in I_\alpha} (M_{E(\alpha,i)})^{(\infty)} \quad (9.3)$$

If $M_{(E_\alpha)^{(\infty)}} \cap \sum_{\beta \in \bar{\Omega}, \beta \neq \alpha} M_{(E_\beta)^{(\infty)}} \neq 0$, then there exists a non-zero simple subcomodule $C^*x \subseteq M_{(E_\alpha)^{(\infty)}} \cap \sum_{\beta \in \bar{\Omega}, \beta \neq \alpha} M_{(E_\beta)^{(\infty)}}$. By [124, Proposition 1.16], $\langle x \rangle$ is a minimal closed subcomodule of M . By Lemma 9.1.1(3), $M_{(E_\alpha)^{(\infty)}}$ is a closed subcomodule of M . By [124, Lemma 3.3], there exists $\gamma \in \bar{\Omega}$ with $\gamma \neq \alpha$ such that $C^*x \subseteq M_{(E_\gamma)^{(\infty)}}$. Thus $M_{(E_\gamma)^{(\infty)}} \cap M_{(E_\alpha)^{(\infty)}} \neq 0$. By Lemma 9.1.1 (7), $(E_\gamma)^{(\infty)} \cap (E_\alpha)^{(\infty)} \neq 0$, which contradicts [121]. Thus for any $\alpha \in \bar{\Omega}$, $M_{(E_\alpha)^{(\infty)}} \cap \sum_{\beta \in \bar{\Omega}, \beta \neq \alpha} M_{(E_\beta)^{(\infty)}} = 0$, which implies that

$$M = \sum_{\alpha \in \bar{\Omega}} \oplus M_{(E_\alpha)^{(\infty)}}.$$

It follows from equations (9.2) and (9.3) that

$$M_{(E_\alpha)^{(\infty)}} = \sum_{i \in I_\alpha} (M_{E(\alpha, i)})^{(\infty)} \quad (9.4)$$

We see that

$$\begin{aligned} M_{E_\alpha} \wedge M_{E_\alpha} &\subseteq M_{E_\alpha \wedge E_\alpha} \quad (\text{by Lemma 9.1.6 (1)}) \\ &\subseteq M_{(E_\alpha)^{(\infty)}}. \end{aligned}$$

Thus

$$M_{(E_\alpha)^{(\infty)}} \supseteq (M_{E_\alpha})^{(\infty)} \supseteq \sum_{i \in I_\alpha} (M_{E(\alpha, i)})^{(\infty)}$$

and

$$M_{(E_\alpha)^{(\infty)}} = (M_{E_\alpha})^{(\infty)}$$

by relation (9.4). This completes the proof. $\square$

Definition 9.1.9. *If M can be decomposed into a direct sum of two non-zero weak-closed subcomodules, then M is called decomposable. If N is a subcomodule of M and N contains exactly one non-zero minimal closed submodule, then N is said to be relative-irreducible.*

Corollary 9.1.10. *C is a coalgebra.*

- (1) *If C is π -commutative, then every C -comodule M is also π -commutative;*
- (2) *If C can be decomposed into a direct sum of its irreducible subcoalgebras, then every C -comodule M can also be decomposed into a direct sum of its relative-irreducible subcomodules;*
- (3) *If C is decomposable, then every component faithfulness C -comodule M is decomposable;*
- (4) *If C is irreducible, then every non-zero C -comodule M is relative-irreducible;*
- (5) *C is irreducible iff every component faithfulness C -comodule M is relative-irreducible.*

Proof. (1) For any pair of non-zero closed subcomodules N and L of M , by Theorem 9.1.3, there exist faithful simple subcoalgebras D and E of C to M such that $N = M_D$ and $L = M_E$. By Lemma 9.1.6(2), $N \wedge L = M_D \wedge M_E = M_{D \wedge E} = M_{E \wedge D} = M_E \wedge M_D = L \wedge N$. Thus M is π -commutative.

(2) Since C can be decomposed into a direct sum of its irreducible subcoalgebras, every equivalence class of C contains only one element by [121]. By Theorem 9.1.8(1), every equivalence class of M also contains only one element. Thus it follows from [124, Theorem 4.18] that M can be decomposed into a direct sum of its relative-irreducible subcomodules.

(3) If C is decomposable, then there are at least two non-zero equivalence classes in C . By Theorem 9.1.8(2), there are at least two non-zero equivalence classes in M . Thus M is decomposable.

(4) If C is irreducible, then there is only one non-zero simple subcoalgebra of C and there is at most one non-zero minimal closed subcomodule in M by Theorem 9.1.3. Considering $M \neq 0$, we have that M is relative-irreducible.

(5) If C is irreducible, then every component faithfulness C -comodule M is relative-irreducible by Corollary 9.1.10(4). Conversely, let $M = C$ be the regular C -comodule. If D is a non-zero

simple subcoalgebra of C , then $0 \neq D \subseteq M_D$. Thus M is a component faithfulness C -comodule. Since M is a relative-irreducible C -comodule by assumption, there is only one non-zero minimal closed subcomodule in M and so there is also only one non-zero simple subcoalgebra in C by Theorem 9.1.3. Thus C is irreducible. $\square$

Lemma 9.1.11. *Let N be a C -subcomodule of M and $\emptyset \neq L \subseteq M$. Then*

- (1) $C^* \cdot L = C(M)^* \cdot L$; $N^{\perp C^* \perp M} = N^{\perp C(M)^* \perp M}$;
- (2) N is a (weak-) closed C -subcomodule iff N is a (weak-)closed $C(M)$ -subcomodule;
- (3) N is a minimal closed C -subcomodule iff N is a minimal closed $C(M)$ -subcomodule;
- (4) N is an indecomposable C -subcomodule iff N is an indecomposable $C(M)$ -subcomodule;
- (5) N is a relative-irreducible C -subcomodule iff N is a relative-irreducible $C(M)$ -subcomodule.

Proof. (1) Let $C = C(M) \oplus V$, where V is a subspace of C . If $f \in V^*$, then $f \cdot L = 0$. Thus $C^* \cdot L = (C(M)^* + V^*) \cdot L = C(M)^* \cdot L$. We now show the second equation. Obviously,

$$N^{\perp C^* \perp M} \subseteq N^{\perp C(M)^* \perp M}.$$

Conversely, for any $x \in N^{\perp C(M)^* \perp M}$ and $f \in N^{\perp C^*}$, there exist $f_1 \in C(M)^*$ and $f_2 \in V^*$ such that $f = f_1 + f_2$. Obviously, $f \cdot x = f_1 \cdot x = 0$. Thus $x \in N^{\perp C^* \perp M}$ and $N^{\perp C(M)^* \perp M} \subseteq N^{\perp C^* \perp M}$. Therefore

$$N^{\perp C^* \perp M} = N^{\perp C(M)^* \perp M}.$$

(2) If N is a weak-closed C -subcomodule, then $(C^* \cdot x)^{\perp C^* \perp M} \subseteq N$ for any $x \in N$. We see that

$$\begin{aligned} (C^* \cdot x)^{\perp C^* \perp M} &= (C(M)^* \cdot x)^{\perp C^* \perp M} \quad (\text{by Lemma 9.1.11(1)}) \\ &= (C(M)^* \cdot x)^{\perp C(M)^* \perp M} \quad (\text{by Lemma 9.1.11(1)}) \end{aligned}$$

Thus $(C(M)^* \cdot x)^{\perp C(M)^* \perp M} \subseteq N$ and so N is a weak-closed $C(M)$ -subcomodule. Conversely, if N is a weak-closed $C(M)$ -subcomodule. Similarly, we can show that N is a weak-closed C -subcomodule. We now show the second assertion. If N is a closed C -subcomodule, then $N^{\perp C^* \perp M} = N$ and $N = N^{\perp C^* \perp M} = N^{\perp C(M)^* \perp M}$ by part (1), which implies that N is a closed $C(M)$ -subcomodule. Conversely, if N is a closed $C(M)$ -subcomodule, similarly, we can show that N is a closed C -subcomodule.

Similarly the others can be proved. $\square$

Proposition 9.1.12. *Let every simple subcoalgebra in $C(M)$ be faithful to M .*

- (1) *If M is an indecomposable C -comodule, then $C(M)$ is also an indecomposable subcoalgebra.*
- (2) *M is a relative-irreducible C -comodule iff $C(M)$ is an irreducible subcoalgebra.*

Proof. (1) If M is an indecomposable C -comodule, then M is an indecomposable $C(M)$ -comodule by Lemma 9.1.11 and M is a component faithfulness $C(M)$ -comodule. By Corollary 9.1.10(3), $C(M)$ is indecomposable.

(2) If M is a relative-irreducible C -comodule, then M is a relative-irreducible $C(M)$ -comodule by Lemma 9.1.11(5) and $C(M)$ is irreducible. Conversely, if $C(M)$ is irreducible, then M is a relative-irreducible $C(M)$ -comodule by Corollary 9.1.10(4) and so M is a relative-irreducible C -comodule by Lemma 9.1.11(5). $\square$

Definition 9.1.13. *If $D \sim E$ implies $M_D \sim M_E$ for any simple subcoalgebras D and E in $C(M)$, then M is called a W -relational hereditary C -comodule.*

If M is a W -relational hereditary C -comodule, then $\mathcal{C}(M)_1 = \mathcal{C}(M)_0$. In fact, by Lemma 9.1.1(5), $\mathcal{C}(M)_1 \subseteq \mathcal{C}(M)_0$. If $\mathcal{C}(M)_1 \neq \mathcal{C}(M)_0$, then there exists $0 \neq D \in \mathcal{C}(M)_0$ such that $M_D = 0$. Since $M_D \sim M_0$ and M is W -relational hereditary, we have that $D \sim 0$ and $D = 0$ by Lemma 9.1.7(1). We get a contradiction. Thus $\mathcal{C}(M)_1 = \mathcal{C}(M)_0$.

Obviously, every π -commutative comodule is W -relational hereditary. If C is π -commutative, then M is π -commutative by Corollary 9.1.10(1) and every C -comodule M is W -relational hereditary. Furthermore, M is also a component faithfulness $C(M)$ -comodule.

Proposition 9.1.14. *Let the notation be the same as in Theorem 9.1.8. Then the following conditions are equivalent.*

- (1) M is W -relational hereditary.
- (2) For any $\alpha \in \bar{\Omega}$, there is at most one non-zero equivalence class in $\{M_{\mathcal{E}(\alpha, i)} \mid i \in I_\alpha\}$, and $\mathcal{C}(M)_1 = \mathcal{C}(M)_0$.
- (3) For any $D \in \mathcal{C}(M)_0$,
 $[M_D] = \{ M_F \mid F \in [D] \text{ and } F \subseteq C(M) \}$
- (4) For any D and $E \in \mathcal{C}(M)_0$, $M_D \sim M_E$ iff $D \sim E$.
- (5) For any $\alpha \in \bar{\Omega}$, $M_{(E_\alpha)^{(\infty)}}$ is indecomposable, and $\mathcal{C}(M)_1 = \mathcal{C}(M)_0$.
- (6) For any $\alpha \in \bar{\Omega}$, $(M_{E_\alpha})^{(\infty)}$ is indecomposable, and $\mathcal{C}(M)_1 = \mathcal{C}(M)_0$;
- (7) $M = \sum_{\alpha \in \bar{\Omega}} \oplus (M_{E_\alpha})^{(\infty)}$ is its weak-closed indecomposable decomposition, and $\mathcal{C}(M)_1 = \mathcal{C}(M)_0$;
- (8) $M = \sum_{\alpha \in \bar{\Omega}} \oplus M_{(E_\alpha)^{(\infty)}}$ is its weak-closed indecomposable decomposition, and $\mathcal{C}(M)_1 = \mathcal{C}(M)_0$.

Proof. We prove it along with the following lines: (1) $\implies$ (4) $\implies$ (3) $\implies$ (1) (3) $\iff$ (2) $\iff$ (5) $\iff$ (6) $\iff$ (7) $\iff$ (8).

(1) $\implies$ (4) It follows from the discussion above that

$$\mathcal{C}(M)_1 = \mathcal{C}(M)_0.$$

If D and $E \in \mathcal{C}(M)_0$ and $M_D \sim M_E$, then $D \sim E$ by Lemma 9.1.7 (2). If $D \sim E$ and D and $E \in \mathcal{C}(M)_0$, then $M_D \sim M_E$.

(4) $\implies$ (3) Considering Lemma 9.1.7(4) and $\mathcal{C}(M)_1 = \mathcal{C}(M)_0$, we only need to show that

$$\{M_F \mid F \in [D] \text{ and } F \subseteq C(M)\} \subseteq [M_D].$$

For any $F \in [D]$ with $F \subseteq C(M)$, $F \sim D$ and $M_F \sim M_D$ by part (4), which implies that $M_F \in [M_D]$.

(3) $\implies$ (1) It is trivial.

(5) $\iff$ (6) $\iff$ (7) $\iff$ (8) It follows from Theorem 9.1.8(3).

(5) $\iff$ (2) By Theorem 9.1.8 (3),

$$M_{(E_\alpha)^{(\infty)}} = \sum_{i \in I_\alpha} (M_{E(\alpha, i)})^{(\infty)}.$$

Thus part (2) and part (5) are equivalent.

(2) $\implies$ (3) It follows from Lemma 9.1.7 (4).

(3) $\implies$ (2) If there are two non-zero equivalence classes $M_{\mathcal{E}(\alpha, i_1)} \neq 0$ and $M_{\mathcal{E}(\alpha, i_2)} \neq 0$ in $\{M_{\mathcal{E}(\alpha, i)} \mid i \in I_\alpha\}$, then there exist $D_1 \in \mathcal{E}(\alpha, i_1)$ and $D_2 \in \mathcal{E}(\alpha, i_2)$ such that $M_{D_1} \neq 0$ and $M_{D_2} \neq 0$. Let $D = D_1$. Since M_{D_1} and $M_{D_2} \in \{M_F \mid F \in [D] \text{ and } F \subseteq C(M)\}$ and $M_{D_2} \notin [M_D] = M_{\mathcal{E}(\alpha, i_1)}$, this contradicts part (3). Thus there is at most one non-zero equivalence class in $\{M_{\mathcal{E}(\alpha, i)} \mid i \in I_\alpha\}$. $\square$

Proposition 9.1.15. *If M is a full, W -relational hereditary C -comodule, then*

(1) *M is indecomposable iff C is indecomposable;*

(2) *M is relative-irreducible iff C is irreducible.*

(3) *M can be decomposed into a direct sum of its weak-closed relative-irreducible subcomodules iff C can be decomposed into a direct sum of its irreducible subcoalgebras.*

(4) *M is π -commutative iff C is π -commutative.*

Proof. Since M is a full C -comodule, $C(M) = C$. Since M is W -relational hereditary, $\mathcal{C}(M)_1 = \mathcal{C}(M)_0 = \mathcal{C}_0$. Thus M is a component faithfulness C -comodule.

(1) If M is indecomposable, then C is indecomposable by Proposition 9.1.12(1). Conversely, if C is indecomposable, then there is at most one non-zero equivalence class in C . By Proposition 9.1.14(2), there is at most one non-zero equivalence class in M . Thus M is indecomposable.

(4) If M is π -commutative, then there is only one element in every equivalence class of M . By Proposition 9.1.14(4) and Theorem 9.1.3, there is only one element in every equivalence class of C . Thus C is π -commutative by [121]. Conversely, if C is π -commutative, then M is π -commutative by Corollary 9.1.10(1).

(2) It follows from the above discussion and Proposition 9.1.12.

(3) $\iff$ (4) By [124, Theorem 3.8 and Theorem 4.18] and [121], it is easy to check that (3) and (4) are equivalent. $\square$

Proposition 9.1.16. *If $M_D \wedge M_E = M_E \wedge M_D$ implies $D \wedge E = E \wedge D$ for any simple coalgebras D and E in $C(M)$, then M is W -relational hereditary.*

Proof. Let $D \sim E$. For any pair of subclasses $\mathcal{M}_1$ and $\mathcal{M}_2$ of $\mathcal{M}_0$ with $\mathcal{M}_1 \cap \mathcal{M}_2 = \emptyset$ and $\mathcal{M}_1 \cup \mathcal{M}_2 = \mathcal{M}_0$, if $M_D \in \mathcal{M}_1$ and $M_E \in \mathcal{M}_2$, let $\mathcal{C}_1 = \{F \in \mathcal{C}_0 \mid M_F \in \mathcal{M}_1\}$ and $\mathcal{C}_2 = \{F \in \mathcal{C}_0 \mid M_F \in \mathcal{M}_2\}$. By Theorem 9.1.3, $\mathcal{C}_0 = \mathcal{C}_1 \cup \mathcal{C}_2$ and $\mathcal{C}_1 \cap \mathcal{C}_2 = \emptyset$. Obviously, $D \in \mathcal{C}_1$ and $E \in \mathcal{C}_2$. By Definition 9.1.4, there exist $D_1 \in \mathcal{C}_1$ and $E_1 \in \mathcal{C}_2$ such that $D_1 \wedge E_1 \neq E_1 \wedge D_1$. By the assumption condition, we have that $M_{D_1} \wedge M_{E_1} \neq M_{E_1} \wedge M_{D_1}$. Thus $M_D \sim M_E$, i.e. M is W -relational hereditary. $\square$

Proposition 9.1.17. *Let $M = C$ as a right C -comodule. Let $N = D$ and $L = E$ with N and L as subcomodules of M with D and E as right coideals of C . Let X be an ideal of C^* and F be a subcoalgebra of C . Then:*

- (1) $X^{\perp C} = X^{\perp M}$; $N^{\perp C^* \perp M} = C(N)$; $C(M_F) = F$;
- (2) N is a closed subcomodule of M iff D is a subcoalgebra of C ;
- (3) N is a closed subcomodule iff N is a weak-closed subcomodule of M ;
- (4) N is a minimal closed subcomodule of M iff D is a simple subcoalgebra of C ;
- (5) When N and L are closed subcomodules, $N \wedge_M L = D \wedge_C E$, where $\wedge_M$ and $\wedge_C$ denote wedge in comodule M and in coalgebra C respectively;
- (6) M is a full and W -relational hereditary C -comodule and a component faithfulness C -comodule.
- (7) The weak-closed indecomposable decomposition of M as a C -comodule and the indecomposable decomposition of C as coalgebra are the same.

Proof. (1) If $x \in X^{\perp M}$, then $f \cdot x = 0$ for any $f \in X$. Let $\rho(x) = \sum_{i=1}^n x_i \otimes c_i$ and $x_1, \dots, x_n$ be linearly independent. Thus $f(c_i) = 0$ and $i = 1, \dots, n$. Since $x = \sum_{i=1}^n \epsilon(x_i)c_i$, $f(x) = 0$, which implies $x \in X^{\perp C}$.

Conversely, if $x \in X^{\perp C}$, we have that $\rho(x) \in X^{\perp C} \otimes X^{\perp C}$ since $X^{\perp C}$ is subcoalgebra of C . Thus $f \cdot x = 0$ for any $f \in X$, which implies $x \in X^{\perp M}$. Thus $X^{\perp M} = X^{\perp C}$. By Lemma 9.1.1(1), $N^{\perp C^* \perp M} = C(N)^{\perp C^* \perp M}$. By part (1), $C(N)^{\perp C^* \perp M} = C(N)^{\perp C^* \perp C}$. Thus $N^{\perp C^* \perp M} = C(N)^{\perp C^* \perp C} = C(N)$.

Finally, we show that $C(M_F) = F$. Obviously, $C(M_F) \subseteq F$. If we view F as a C -subcomodule of M , then $F \subseteq C(M_F)$. Thus $F = C(M_F)$.

(2) If N is a closed subcomodule of M , then $N^{\perp C^* \perp M} = N$. By Proposition 9.1.17(1), $N^{\perp C^* \perp M} = C(N) = N$. Thus $\rho(N) = \Delta(D) \subseteq N \otimes N = D \otimes D$, which implies that D is a subcoalgebra of C . Conversely, if D is a subcoalgebra of C , then $\rho(N) \subseteq N \otimes N$. Thus $C(N) \subseteq D = N$. By Proposition 9.1.17(1), $N^{\perp C^* \perp M} = C(N) \subseteq N$. Thus $N^{\perp C^* \perp M} = N$, i.e. N is closed.

(3) If N is a closed subcomodule, then N is weak-closed. Conversely, if N is weak-closed, then $\langle x \rangle \subseteq N$ for $x \in N$. Since $\langle x \rangle$ is a closed subcomodule of M , $\langle x \rangle$ is subcoalgebra of C if let $\langle x \rangle$ with structure of coalgebra C . This shows that $\Delta(x) \in \langle x \rangle \otimes \langle x \rangle \subseteq D \otimes D$. Thus D is a subcoalgebra of C . By Proposition 9.1.17(2), N is a closed subcomodule of M .

(4) It follows from part (2).

(5) We only need to show that

$$(N^{\perp C^*} L^{\perp C^*})^{\perp M} = (D^{\perp C^*} E^{\perp C^*})^{\perp C}.$$

Since N and L are closed subcomodules, $N^{\perp C^*} = C(N)^{\perp C^*} = D^{\perp C^*}$ and $L^{\perp C^*} = C(L)^{\perp C^*} = E^{\perp C^*}$ by Proposition 9.1.17(1) and Lemma 9.1.1(1). Thus we only need to show that

$$(D^{\perp C^*} E^{\perp C^*})^{\perp M} = (D^{\perp C^*} E^{\perp C^*})^{\perp C}.$$

The above formula follows from Proposition 9.1.17(1).

(6) By the proof of Corollary 9.1.10(5), we know that M is a component faithfulness C -comodule. Let $\{m_\lambda \mid \lambda \in \Lambda\}$ be a basis of M . For any $c \in C$, $\Delta(c) = \sum m_\lambda \otimes c_\lambda$, by the

definition of $C(M)$, $c_\lambda \in C(M)$. Since $c = \sum \epsilon(m_\lambda)c_\lambda \in C(M)$, $C \subseteq C(M)$, i.e. $C = C(M)$. Consequently, it follows from part (4)(5) that M is W -relational hereditary.

(7) Since M is W -relational hereditary, $M = \sum_{\alpha \in \bar{\Omega}} \oplus M_{(E_\alpha)^{(\infty)}}$ is a weak-closed indecomposable decomposition of M by Proposition 9.1.14. By part (1) (3), $C(M_{(E_\alpha)^{(\infty)}}) = (E_\alpha)^{(\infty)} = (M_{(E_\alpha)^{(\infty)}})^{\perp C^* \perp M} = M_{(E_\alpha)^{(\infty)}}$. Thus $M = \sum_{\alpha \in \bar{\Omega}} \oplus M_{(E_\alpha)^{(\infty)}} = \sum_{\alpha \in \bar{\Omega}} \oplus (E_\alpha)^{(\infty)}$. By [121], $\sum_{\alpha \in \bar{\Omega}} \oplus (E_\alpha)^{(\infty)} = C$ is a indecomposable decomposition of C . Thus the weak-closed indecomposable decomposition of M as a C -comodule and the indecomposable decomposition of C as coalgebra are the same. This completes the proof. $\square$

By Lemma 9.1.6(4) and Proposition 9.1.17, C is cosemisimple iff every C -comodule M is cosemisimple.

9.2 The coradicals of comodules

Proposition 9.2.1. *Let M be a C -comodule, J denote the Jacobson radical of C^* and $r_j(C(M)^*)$ denote the Jacobson radical of $C(M)^*$.*

(1)

$$M_0 = (r_j(C(M)^*))^{\perp M} = \text{Soc}_{C^*} M \quad \text{and} \quad r_j(C(M)^*) = M_0^{\perp C^*};$$

(2) *If we view C as a right C -comodule, then*

$$C_0 = \text{Soc}_{C^*} C = J^{\perp C} = \sum \{D \mid D \text{ is a minimal right coideal of } C\}.$$

Proof. (1) We first show that

$$M_0^{\perp C^*} \subseteq J$$

when M is a full C -comodule. We only need to show that $\epsilon - f$ is invertible in C^* for any $f \in (M_0)^{\perp C^*}$. Let $I = (M_0)^{\perp C^*}$. For any natural number n , $f^{n+1} \cdot (M_0)^{(n)} = 0$ since $f^{n+1} \in I^{n+1}$, where $(M_0)^{(n)} = \wedge^{n+1} M_0$. Thus

$$f^{n+1}(C((M_0)^{(n)})) = 0. \tag{9.5}$$

by Lemma 9.1.1(1) and

$$M = (M_0)^{(\infty)} = \cup \{(M_0)^{(n)} \mid n = 0, 1, \dots\} \tag{9.6}$$

by [124, Theorem 4.7]. We now show that

$$C = \cup \{C((M_0)^{(n)}) \mid n = 0, 1, \dots\} \tag{9.7}$$

Since $(M_0)^{(n)} \subseteq (M_0)^{(n+1)}$, we have that there exists a basis $\{m_\lambda \mid \lambda \in \Lambda\}$ of M such that for every given natural number n there exists a subset of $\{m_\lambda \mid \lambda \in \Lambda\}$, which is a basis of $(M_0)^{(n)}$. For any $c \in W(M)$, there exists $m \in M$ with $\rho(m) = \sum m_\lambda \otimes c_\lambda$ such that $c_{\lambda_0} = c$ for some $\lambda_0 \in \Lambda$. By equation (9.6), there exists a natural number n such that $m \in (M_0)^{(n)}$, which implies that $c \in C((M_0)^{(n)})$ and $C(M) \subseteq \cup_0^\infty C((M_0)^{(n)})$. Thus

$$C = \cup \{C((M_0)^{(n)}) \mid n = 0, 1, \dots\}.$$

Let

$$g_n = \epsilon + f + \cdots + f^n, n = 1, 2, \cdots .$$

For any $c \in C$, there exists a natural number n such that $c \in C((M_0)^{(n)})$ by relation (9.7). We define

$$g(c) = g_n(c) . \quad (9.8)$$

Considering relation (9.5), we have that g is well-defined. Thus $g \in C^*$.

We next show that g is an inverse of $\epsilon - f$ in C^* . For any $c \in C^*$, there exists a natural number n such that $c \in C((M_0)^{(n)})$ by relation (9.7). Thus $\Delta(c) \in C((M_0)^{(n)}) \otimes C((M_0)^{(n)})$. We see that

$$\begin{aligned} g * (\epsilon - f)(c) &= \sum g(c_1)(\epsilon - f)(c_2) \\ &= \sum g_n(c_1)(\epsilon - f)(c_2) \text{ (by relation (9.8))} \\ &= (g_n * (\epsilon - f))(c) \\ &= (\epsilon - f^{n+1})(c) \\ &= \epsilon(c) \text{ (by relation (9.5))} . \end{aligned}$$

Thus $g * (\epsilon - f) = \epsilon$. Similarly, $(\epsilon - f) * g = \epsilon$. Thus $\epsilon - f$ has an inverse in C^* .

We next show that

$$M_0^{\perp C^*} = J$$

when $C(M) = C$. It follows from Lemma 9.1.1(1) that

$$\begin{aligned} (M_0)^{\perp C^*} &= \cap \{N^{\perp C^*} \mid N \text{ is a minimal closed subcomodule of } M\} \\ &= \cap \{C(N)^{\perp C^*} \mid N \text{ is a minimal closed subcomodule of } M\} \\ &\supseteq J \text{ (by [54, Proposition 2.1.4] and Proposition 9.1.2) } \end{aligned}$$

Thus $M_0^{\perp C^*} = J$

We now show that

$$r_j(C(M)^*) = M_0^{\perp C^*} \quad \text{and} \quad (r_j(C(M)^*))^{\perp M} = M_0$$

for any C -comodule M . If M is a C -comodule, then M is full $C(M)$ -comodule and so

$$(r_j(C(M)^*)) = M_0^{\perp C^*} .$$

By [124, Proposition 4.6], M_0 is closed. Thus $((r_j(C(M)^*))^{\perp M} = M_0$.

Finally, we show that $M_0 = Soc_{C^*} M$ for any C -comodule M . By [124, Proposition 1.16], $Soc_{C^*} M \subseteq M_0$. Conversely, if $x \in M_0$, let $N = C^*x$. By Lemma 9.1.6(4), $M_{C_0} = M_0$. Thus N is a finite dimensional C_0 -comodule. By [114, Theorem 14.0.1], $N = N_1 \oplus \cdots \oplus N_n$ and N_i is a simple C_0^* -submodule. By [124, Lemma 1.1], N_i is a C_0 -subcomodule. Thus N_i is a C -comodule. By [124, Lemma 1.1], N_i is a C^* -submodule. If L is a non-zero C^* -submodule of M

and $L \subseteq N_i$, then L is also a $(C_0)^*$ -submodule. Thus $L = N_i$. This shows that N_i is also a simple C^* -submodule. Thus $N_i \subseteq Soc_{C^*}M$ and $N \subseteq Soc_{C^*}M$. It follows from the above proof that $M_0 = Soc_{C^*}M$.

(2) It follows from Proposition 9.1.17 and part (1).

Chapter 10

The Homological Dimensions of Crossed Products

Throughout this chapter, k is a field, R is an algebra over k , and H is a Hopf algebra over k . We say that $R \#_{\sigma} H$ is the crossed product of R and H if $R \#_{\sigma} H$ becomes an algebra over k by multiplication:

$$(a \# h)(b \# g) = \sum_{h, g} a(h_1 \cdot b) \sigma(h_2, g_1) \# h_3 g_2$$

for any $a, b \in R, h, g \in H$, where σ is a k -linear map from $H \otimes H$ to R and $\Delta(h) = \sum h_1 \otimes h_2$ (see [87, Definition 7.1.1].)

Let $lpd(RM)$, $lid(RM)$ and $lfd(RM)$ denote the left projective dimension, the left injective dimension and the left flat dimension of left R -module M respectively. Let $lgD(R)$ and $wD(R)$ denote the left global dimension and the weak dimension of algebra R respectively.

Crossed products are very important algebraic structures. The relation between homological dimensions of algebra R and crossed product $R \#_{\sigma} H$ is often studied. J.C. McConnell and J.C. Robson in [89, Theorem 7.5.6] obtained that

$$rgD(R) = rgD(R * G)$$

for any finite group G with $|G|^{-1} \in k$, where $R * G$ is the skew group ring. It is clear that every skew group ring $R * G$ is crossed product $R \#_{\sigma} kG$ with trivial σ . Zhong Yi in [126] obtained that the global dimension of crossed product $R * G$ is finite when the global dimension of R is finite and some other conditions hold.

In this chapter, we obtain that the global dimensions of R and the crossed product $R \#_{\sigma} H$ are the same; meantime, their weak dimensions are also the same, when H is a finite-dimensional semisimple and cosemisimple Hopf algebra.

10.1 The homological dimensions of modules over crossed products

In this section, we give a relation between homological dimensions of modules over R and $R \#_{\sigma} H$.

If M is a left (right) $R\#_\sigma H$ -module, then M is also a left (right) R -module since we can view R as a subalgebra of $R\#_\sigma H$.

Lemma 10.1.1. *Let R be a subalgebra of algebra A .*

- (i) *If M is a free A -module and A is a free R -module, then M is a free R -module;*
- (ii) *If P is a projective left $R\#_\sigma H$ -module, then P is a projective left R -module;*
- (iii) *If P is a projective right $R\#_\sigma H$ -module and H is a Hopf algebra with invertible antipode, then P is a projective right R -module;*
- (iv) *If*

$$\mathcal{P}_M : \quad \cdots P_n \xrightarrow{d_n} P_{n-1} \cdots \rightarrow P_0 \xrightarrow{d_0} M \rightarrow 0$$

is a projective resolution of left $R\#_\sigma H$ -module M , then $\mathcal{P}_M$ is a projective resolution of left R -module M ;

- (v) *If*

$$\mathcal{P}_M : \quad \cdots P_n \xrightarrow{d_n} P_{n-1} \cdots \rightarrow P_0 \xrightarrow{d_0} M \rightarrow 0$$

is a projective resolution of right $R\#_\sigma H$ -module M and H is a Hopf algebra with invertible antipode, then $\mathcal{P}_M$ is a projective resolution of right R -module M .

Proof. (i) It is obvious.

(ii) Since P is a projective $R\#_\sigma H$ -module, we have that there exists a free $R\#_\sigma H$ -module F such that P is a summand of F . It is clear that $R\#_\sigma H \cong R \otimes H$ as left R -modules, which implies that $R\#_\sigma H$ is a free R -module. Thus it follows from part (i) that F is a free R -module and P is a summand of F as R -module. Consequently, P is a projective R -module.

(iii) By [87, Corollary 7.2.11], $R\#_\sigma H \cong H \otimes R$ as right R -modules. Thus $R\#_\sigma H$ is a free right R -module. Using the method in the proof of part (i), we have that P is a projective right R -module.

(iv) and (v) can be obtained by part (ii) and (iii). $\square$

Lemma 10.1.2. (i) *Let R be a subalgebra of an algebra A . If M is a flat right (left) A -module and A is a flat right (left) R -module, then M is a flat right (left) R -module;*

- (ii) *If F is a flat left $R\#_\sigma H$ -module, then F is a flat left R -module;*
- (iii) *If F is a flat right $R\#_\sigma H$ -module and H is a Hopf algebra with invertible antipode, then F is a flat right R -module;*
- (iv) *If*

$$\mathcal{F}_M : \quad \cdots F_n \xrightarrow{d_n} F_{n-1} \cdots \rightarrow F_0 \xrightarrow{d_0} M \rightarrow 0$$

is a flat resolution of left $R\#_\sigma H$ -module M , then $\mathcal{F}_M$ is a flat resolution of left R -module M ;

- (v) *If*

$$\mathcal{F}_M : \quad \cdots F_n \xrightarrow{d_n} F_{n-1} \cdots \rightarrow F_0 \xrightarrow{d_0} M \rightarrow 0$$

is a flat resolution of right $R\#_\sigma H$ -module M and H is a Hopf algebra with invertible antipode, then $\mathcal{F}_M$ is a flat resolution of M .

Proof. (i) We only show part (i) in the case which M is a right A -module and A is a right R -module; the other cases can be shown similarly. Let

$$0 \rightarrow X \xrightarrow{f} Y$$

be an exact left $R\#_\sigma H$ -module sequence. By umptions,

$$0 \rightarrow A \otimes_R X \xrightarrow{A \otimes f} A \otimes_R Y$$

and

$$0 \rightarrow M \otimes_A (A \otimes_R X) \xrightarrow{M \otimes (A \otimes f)} M \otimes_A (A \otimes_R Y)$$

are exact sequences. Obviously,

$$M \otimes_A (A \otimes_R X) \cong M \otimes_R X \quad \text{and} \quad M \otimes_A (A \otimes_R Y) \cong M \otimes_R Y$$

as additive groups. Thus

$$0 \rightarrow M \otimes_R X \xrightarrow{M \otimes f} M \otimes_R Y$$

is an exact sequence, which implies M is a flat R -module.

(ii)–(v) are immediate consequence of part (i). $\square$

The following is an immediate consequence of Lemma 10.1.1 and 10.1.2.

Proposition 10.1.3. (i) If M is a left $R\#_\sigma H$ -module, then

$$lpd({}_R M) \leq lpd({}_{R\#_\sigma H} M);$$

(ii) If M is a right $R\#_\sigma H$ -module and H is a Hopf algebra with invertible antipode, then

$$rpd(M_R) \leq rpd(M_{R\#_\sigma H});$$

(iii) If M is a left $R\#_\sigma H$ -module, then

$$lfd({}_R M) \leq lfd({}_{R\#_\sigma H} M);$$

(iv) If M is a right $R\#_\sigma H$ -module and H is a Hopf algebra with invertible antipode, then

$$rfd(M_R) \leq rfd(M_{R\#_\sigma H}).$$

Lemma 10.1.4. Let H be a finite-dimensional semisimple Hopf algebra, and let M and N be left $R\#_\sigma H$ -modules. If f is an R -module homomorphism from M to N , and

$$\bar{f}(m) = \sum \gamma^{-1}(t_1) f(\gamma(t_2)m)$$

for any $m \in M$, then $\bar{f}$ is an $R\#_\sigma H$ -module homomorphism from M to N , where $t \in \int_H^r$ with $\epsilon(t) = 1$, γ is a map from H to $R\#_\sigma H$ sending h to $1\#h$ and the convolution-inverse

$$\gamma^{-1}(h) = \sum \sigma^{-1}(S(h_2), h_3) \# S(h_1)$$

for any $h \in H$.

Proof. (See the proof of [87, Theorem 7.4.2]) For any $a \in R, h \in H, m \in M$, we see that

$$\begin{aligned}
\bar{f}(am) &= \sum \gamma^{-1}(t_1) f(\gamma(t_2)am) \\
&= \sum \gamma^{-1}(t_1) f((t_2 \cdot a)\gamma(t_3)m) \\
&= \sum \gamma^{-1}(t_1) (t_2 \cdot a) f(\gamma(t_3)m) \\
&= \sum a \gamma^{-1}(t_1) f(\gamma(t_2)m) \\
&= a \bar{f}(m)
\end{aligned}$$

and

$$\begin{aligned}
\bar{f}(\gamma(h)m) &= \sum \gamma^{-1}(t_1) f(\gamma(t_2)\gamma(h)m) \\
&= \sum \gamma^{-1}(t_1) f(\sigma(t_2, h_1)\gamma(t_3 h_2)m) \quad \text{by [87, Definition 7.1.1]} \\
&= \sum \gamma^{-1}(t_1) \sigma(t_2, h_1) f(\gamma(t_3 h_2)m) \\
&= \sum \gamma(h_1) \gamma^{-1}(t_1 h_2) f(\gamma(t_2 h_3)m) \\
&= \sum \gamma(h) \gamma^{-1}(t_1) f(\gamma(t_2)m) \quad \text{since } \sum h_1 \otimes t_1 h_2 \otimes t_2 h_3 = \sum h \otimes t_1 \otimes t_2 \\
&= \gamma(h) \bar{f}(m)
\end{aligned}$$

Thus $\bar{f}$ is an $R\#_\sigma H$ -module homomorphism. $\square$

In fact, we can obtain a functor by Lemma 10.1.4. Let $_{R\#_\sigma H}\overline{\mathcal{M}}$ denote the full subcategory of $_{R\#_\sigma H}\mathcal{M}$, its objects are all the left $R\#_\sigma H$ -modules and its morphisms from M to N are all the R -module homomorphisms from M to N . For any $M, N \in ob_{R\#_\sigma H}\overline{\mathcal{M}}$ and R -module homomorphism f from M to N , we define that

$$F : _{R\#_\sigma H}\overline{\mathcal{M}} \longrightarrow _{R\#_\sigma H}\mathcal{M}$$

such that

$$F(M) = M \quad \text{and} \quad F(f) = \bar{f},$$

where $\bar{f}$ is defined in Lemma 10.1.4. It is clear that F is a functor.

Lemma 10.1.5. *Let H be a finite-dimensional semisimple Hopf algebra, and let M and N be right $R\#_\sigma H$ -modules. If f is an R -module homomorphism from M to N , and*

$$\bar{f}(m) = \sum f(m\gamma^{-1}(t_1))\gamma(t_2)$$

for any $m \in M$, then $\bar{f}$ is an $R\#_\sigma H$ -module homomorphism from M to N , where $t \in \int_H^r$ with $\epsilon(t) = 1$, γ is a map from H to $R\#_\sigma H$ sending h to $1\#h$.

Proof. (See the proof of [87, Theorem 7.4.2].) For any $a \in R$ and $h \in H, m \in M$, we see that

$$\begin{aligned}
\bar{f}(ma) &= \sum f(ma\gamma^{-1}(t_1))\gamma(t_2) \\
&= \sum f(m\gamma^{-1}(t_1)(t_2 \cdot a))\gamma(t_3) \\
&= \sum f(m\gamma^{-1}(t_1))(t_2 \cdot a)\gamma(t_3) \\
&= \sum f(m\gamma^{-1}(t_1))\gamma(t_2)a \\
&= \bar{f}(m)a
\end{aligned}$$

and

$$\begin{aligned}
\bar{f}(m\gamma(h)) &= \sum f(m\gamma(h)\gamma^{-1}(t_1))\gamma(t_2) \\
&= \sum f(m\gamma(h_1)\gamma^{-1}(t_1h_2))\gamma(t_2h_3) \quad \text{since } \sum h_1 \otimes t_1h_2 \otimes t_2h_3 = \sum h \otimes t_1 \otimes t_2 \\
&= \sum f(m\gamma^{-1}(t_1)\sigma(t_2, h_1))\gamma(t_3h_2) \quad \text{by [87, Definition 7.1.1]} \\
&= \sum f(m\gamma^{-1}(t_1))\sigma(t_2, h_1)\gamma(t_3h_2) \\
&= \sum f(m\gamma^{-1}(t_1))\gamma(t_2)\gamma(h) \\
&= \bar{f}(m)\gamma(h)
\end{aligned}$$

Thus $\bar{f}$ is an $R\#_\sigma H$ -module homomorphism. $\square$

Proposition 10.1.6. *Let H be a finite-dimensional semisimple Hopf algebra.*

(i) *If P is a left (right) $R\#_\sigma H$ -modules and a projective left (right) R -module, then P is a projective left (right) $R\#_\sigma H$ -module;*

(ii) *If E is a left (right) $R\#_\sigma H$ -modules and an injective left (right) R -module, then E is an injective left (right) $R\#_\sigma H$ -module;*

(iii) *If F is a left (right) $R\#_\sigma H$ -modules and a flat left (right) R -module, then F is a flat left (right) $R\#_\sigma H$ -module.*

Proof. (i) Let

$$X \xrightarrow{f} Y \rightarrow 0$$

be an exact sequence of left (right) $R\#_\sigma H$ -modules and g be a $R\#_\sigma H$ -module homomorphism from P to Y . Since P is a projective left (right) R -module, we have that there exists an R -module homomorphism φ from P to X such that

$$f\varphi = g.$$

By Lemma 10.1.4 and 10.1.5, there exists an $R\#_\sigma H$ -module homomorphism $\bar{\varphi}$ from P to X such that

$$f\bar{\varphi} = g.$$

Thus P is a projective left (right) $R\#_\sigma H$ -module.

Similarly, we can obtain the proof of part (ii).

(iii) Since F is a flat left (right) R -module, we have that characteristic module $\text{Hom}_{\mathcal{Z}}(F, \mathcal{Q}/\mathcal{Z})$ of F is an injective left (right) R -module by [119, Theorem 2.3.6]. Obviously, $\text{Hom}_{\mathcal{Z}}(F, \mathcal{Q}/\mathcal{Z})$ is a left (right) $R\#_\sigma H$ -module. By part (ii), $\text{Hom}_{\mathcal{Z}}(F, \mathcal{Q}/\mathcal{Z})$ is an injective left (right) $R\#_\sigma H$ -module. Thus F is a flat left (right) $R\#_\sigma H$ -module. $\square$

Proposition 10.1.7. *Let H be a finite-dimensional semisimple Hopf algebra. Then for left (right) $R\#_\sigma H$ -modules M and N ,*

$$\text{Ext}_{R\#_\sigma H}^n(M, N) \subseteq \text{Ext}_R^n(M, N),$$

where n is any natural number.

Proof. We view the $Ext^n(M, N)$ as the equivalent classes of n -extensions of M and N (see [119, Definition 3.3.7]). We only prove this result for the case $n = 1$. For other cases, we can prove them similarly. We denote the equivalent classes in $Ext_{R\#_\sigma H}^1(M, N)$ and $Ext_R^1(M, N)$ by $[E]$ and $[F]'$ respectively, where E is an extension of $R\#_\sigma H$ -modules M and N , and F is an extension of R -modules M and N . We define a map

$$\Psi : Ext_{R\#_\sigma H}^1(M, N) \rightarrow Ext_R^1(M, N), \quad \text{by sending } [E] \text{ to } [E]'.$$

Obviously, Ψ is a map. Now we show that Ψ is injective. Let

$$0 \rightarrow M \xrightarrow{f} E \xrightarrow{g} N \rightarrow 0 \quad \text{and} \quad 0 \rightarrow M \xrightarrow{f'} E' \xrightarrow{g'} M \rightarrow 0$$

are two extensions of $R\#_\sigma H$ -modules M and N , and they are equivalent in $Ext_R^1(M, N)$. Thus there exists an R -module homomorphism φ from E to E' such that

$$\varphi f = f' \quad \text{and} \quad \varphi g = g'.$$

By lemma 10.1.4, there exists an $R\#_\sigma H$ -module homomorphism $\bar{\varphi}$ from E to E' such that

$$\bar{\varphi} f = f' \quad \text{and} \quad \bar{\varphi} g = g'.$$

Thus E and E' are equivalent in $Ext_{R\#_\sigma H}^1(M, N)$, which implies that Ψ is injective. $\square$

Lemma 10.1.8. *For any $M \in \mathcal{M}_{R\#_\sigma H}$ and $N \in {}_{R\#_\sigma H}\mathcal{M}$, there exists an additive group homomorphism*

$$\xi : M \otimes_R N \rightarrow M \otimes_{R\#_\sigma H} N$$

by sending $(m \otimes n)$ to $m \otimes n$, where $m \in M, n \in N$.

Proof. It is trivial. $\square$

Proposition 10.1.9. *If M is a right $R\#_\sigma H$ -modules and N is a left $R\#_\sigma H$ -module, then there exists an additive group homomorphism*

$$\xi_* : Tor_n^R(M, N) \longrightarrow Tor_n^{R\#_\sigma H}(M, N)$$

such that $\xi_([z_n]) = [\xi(z_n)]$, where ξ is the same as in Lemma 10.1.8.*

Proof. Let

$$\mathcal{P}_M : \quad \cdots P_n \xrightarrow{d_n} P_{n-1} \cdots \rightarrow P_0 \xrightarrow{d_0} M \rightarrow 0$$

be a projective resolution of right $R\#_\sigma H$ -module M , and set

$$T = - \otimes_{R\#_\sigma H} N \quad \text{and} \quad T^R = - \otimes_R N.$$

We have that

$$T\mathcal{P}_{\hat{M}} : \quad \cdots T(P_n) \xrightarrow{Td_n} T(P_{n-1}) \cdots \rightarrow T(P_1) \xrightarrow{Td_1} TP_0 \rightarrow 0$$

and

$$T^R\mathcal{P}_{\hat{M}} : \quad \cdots T^R(P_n) \xrightarrow{T^Rd_n} T^R(P_{n-1}) \cdots \rightarrow T^R(P_1) \xrightarrow{T^Rd_1} T^R(P_0) \rightarrow 0$$

are complexes. Thus ξ is a complex homomorphism from $T^R\mathcal{P}_{\hat{M}}$ to $T\mathcal{P}_{\hat{M}}$, which implies that ξ_* is an additive group homomorphism. $\square$

10.2 The global dimensions and weak dimensions of crossed products

In this section we give the relation between homological dimensions of R and $R\#_\sigma H$.

Lemma 10.2.1. *If R and R' are Morita equivalent rings, then*

- (i) $rgD(R) = rgD(R')$;
- (ii) $lgD(R) = lgD(R')$;
- (iii) $wD(R) = wD(R')$.

Proof. It is an immediate consequence of [2, Proposition 21.6, Exercise 22.12]. $\square$

Theorem 10.2.2. *If H is a finite-dimensional semisimple Hopf algebra, then*

- (i) $rgD(R\#_\sigma H) \leq rgD(R)$;
- (ii) $lgD(R\#_\sigma H) \leq lgD(R)$;
- (iii) $wD(R\#_\sigma H) \leq wD(R)$.

Proof. (i) When $lgD(R)$ is infinite, obviously part (i) holds. Now we assume $lgD(R) = n$. For any left $R\#_\sigma H$ -module M , and a projective resolution of left $R\#_\sigma H$ -module M :

$$\mathcal{P}_M : \quad \cdots P_n \xrightarrow{d_n} P_{n-1} \cdots \rightarrow P_0 \xrightarrow{d_0} M \rightarrow 0,$$

we have that $\mathcal{P}_M$ is also a projective resolution of left R -module M by Lemma 10.1.1. Let $K_n = \ker d_n$ be syzygy n of $\mathcal{P}_M$. Since $lgD(R) = n$, $Ext_R^{n+1}(M, N) = 0$ for any left R -module N by [119, Corollary 3.3.6]. Thus $Ext_R^1(K_n, N) = 0$, which implies K_n is a projective R -module. By Lemma 10.1.6 (i), K_n is a projective $R\#_\sigma H$ -module and $Ext_{R\#_\sigma H}^{n+1}(M, N) = 0$ for any $R\#_\sigma H$ -module N . Consequently,

$$lgD(R\#_\sigma H) \leq n = lgD(R) \quad \text{by [119, Corollary 3.3.6]} .$$

We complete the proof of part (i).

We can show part (ii) and part (iii) similarly. $\square$

Theorem 10.2.3. *Let H be a finite-dimensional semisimple and cosemisimple Hopf algebra. Then*

- (i) $rgD(R) = rgD(R\#_\sigma H)$;
- (ii) $lgD(R) = lgD(R\#_\sigma H)$;
- (iii) $wD(R) = wD(R\#_\sigma H)$.

Proof. (i) By the duality theorem (see [87, Corollary 9.4.17]), we have that $(R\#_\sigma H)\#H^*$ and R are Morita equivalent algebras. Thus $lgD(R) = lgD((R\#_\sigma H)\#H^*)$ by Lemma 10.2.1 (i). Considering Theorem 10.2.2 (i), we have that

$$lgD((R\#_\sigma H)\#H^*) \leq lgD(R\#_\sigma H) \leq lgD(R).$$

Consequently,

$$lgD(R) = lgD(R\#_\sigma H).$$

Similarly, we can prove (ii) and (iii) . $\square$

Corollary 10.2.4. *Let H be a finite-dimensional semisimple Hopf algebra.*

- (i) *If R is a left (right) semi-hereditary, then so is $R\#_{\sigma}H$;*
- (ii) *If R is von Neumann regular, then so is $R\#_{\sigma}H$.*

Proof. (i) It follows from Theorem 10.2.2 and [119, Theorem 2.2.9] .

(ii) It follows from Theorem 10.2.2 and [119, Theorem 3.4.13]. $\square$

By the way, part (ii) of Corollary 10.2.4 gives one case about the semiprime question in [87, Question 7.4.9]. That is, if H is a finite-dimensional semisimple Hopf algebra and R is a von Neumann regular algebra (notice that every von Neumann regular algebra is semiprime), then $R\#_{\sigma}H$ is semiprime.

Corollary 10.2.5. *Let H be a finite-dimensional semisimple and cosemisimple Hopf algebra. Then*

- (i) *R is semisimple artinian iff $R\#_{\sigma}H$ is semisimple artinian;*
- (ii) *R is left (right) semi-hereditary iff $R\#_{\sigma}H$ is left (right) semi-hereditary;*
- (iii) *R is von Neumann regular iff $R\#_{\sigma}H$ is von Neumann regular.*

Proof. (i) It follows from Theorem 10.2.3 and [119, Theorem 2.2.9].

(ii) It follows from Theorem 10.2.3 and [119, Theorem 2.2.9].

(iii) It follows from Theorem 10.2.3 and [119, Theorem 3.4.13]. $\square$

If H is commutative or cocommutative, then $S^2 = id_H$ by [114]. Consequently, by [104, Proposition 2 (c)], H is semisimple and cosemisimple iff the character $chark$ of k does not divides $dimH$. Considering Theorem 10.2.3 and Corollary 10.2.5, we have

Corollary 10.2.6. *Let H be a finite-dimensional commutative or cocommutative Hopf algebra. If the character $chark$ of k does not divides $dimH$, then*

- (i) $rgD(R) = rgD(R\#_{\sigma}H)$;
- (ii) $rgD(R) = rgD(R\#_{\sigma}H)$;
- (iii) $wD(R) = wD(R\#_{\sigma}H)$;
- (iv) *R is semisimple artinian iff $R\#_{\sigma}H$ is semisimple artinian;*
- (v) *R is left (right) semi-hereditary iff $R\#_{\sigma}H$ is left (right) semi-hereditary;*
- (vi) *R is von Neumann regular iff $R\#_{\sigma}H$ is von Neumann regular.*

Since group algebra kG is a cocommutative Hopf algebra, we have that

$$rgD(R) = rgD(R * G).$$

Thus Corollary 10.2.6 implies [89, Theorem 7.5.6].

Chapter 11

The Radicals of Hopf Module Algebras

Remark ¹

In this chapter, the characterization of H -prime radical is given in many ways. Meantime, the relations between the radical of smash product $R\#H$ and the H -radical of Hopf module algebra R are obtained.

In this chapter, let k be a commutative associative ring with unit, H be an algebra with unit and comultiplication Δ (i.e. Δ is a linear map: $H \rightarrow H \otimes H$), R be an algebra over k (R may be without unit) and R be an H -module algebra.

We define some necessary concept as follows.

If there exists a linear map $\begin{cases} H \otimes R & \longrightarrow R \\ h \otimes r & \mapsto h \cdot r \end{cases}$ such that

$$h \cdot rs = \sum (h_1 \cdot r)(h_2 \cdot s) \quad \text{and} \quad 1_H \cdot r = r$$

for all $r, s \in R, h \in H$, then we say that H weakly acts on R . For any ideal I of R , set

$$(I : H) := \{x \in R \mid h \cdot x \in I \text{ for all } h \in H\}.$$

I is called an H -ideal if $h \cdot I \subseteq I$ for any $h \in H$. Let I_H denote the maximal H -ideal of R in I . It is clear that $I_H = (I : H)$. An H -module algebra R is called an H -simple module algebra if R has not any non-trivial H -ideals and $R^2 \neq 0$. R is said to be H -semiprime if there are no non-zero nilpotent H -ideals in R . R is said to be H -prime if $IJ = 0$ implies $I = 0$ or $J = 0$ for any H -ideals I and J of R . An H -ideal I is called an H -(semi)prime ideal of R if R/I is H -(semi)prime. $\{a_n\}$ is called an H - m -sequence in R with beginning a if there exist $h_n, h'_n \in H$ such that $a_1 = a \in R$ and $a_{n+1} = (h_n \cdot a_n)b_n(h'_n \cdot a_n)$ for any natural number n . If every H - m -sequence $\{a_n\}$ with $a_{7.1.1} = a$, there exists a natural number k such that $a_k = 0$, then a is called an H - m -nilpotent element. Set

$$W_H(R) = \{a \in R \mid a \text{ is an } H\text{-}m\text{-nilpotent element}\}.$$

¹This chapter can be omitted

R is called an H -module algebra if the following conditions hold:

- (i) R is a unital left H -module (i.e. R is a left H -module and $1_H \cdot a = a$ for any $a \in R$);
- (ii) $h \cdot ab = \sum (h_1 \cdot a)(h_2 \cdot b)$ for any $a, b \in R$, $h \in H$, where $\Delta(h) = \sum h_1 \otimes h_2$.

H -module algebra is sometimes called a Hopf module.

If R is an H -module algebra with a unit 1_R , then

$$\begin{aligned} h \cdot 1_R &= \sum_h (h_1 \cdot 1_R)(h_2 S(h_3) \cdot 1_R) \\ &= \sum_h h_1 \cdot (1_R(S(h_2) \cdot 1_R)) = \sum_h h_1 S(h_2) \cdot 1_R = \epsilon(h)1_R, \end{aligned}$$

i.e. $h \cdot 1_R = \epsilon(h)1_R$

for any $h \in H$.

An H -module algebra R is called a unital H -module algebra if R has a unit 1_R such that $h \cdot 1_R = \epsilon(h)1_R$ for any $h \in H$. Therefore, every H -module algebra with unit is a unital H -module algebra. A left R -module M is called an R - H -module if M is also a left unital H -module with $h(am) = \sum (h_1 \cdot a)(h_2 m)$ for all $h \in H, a \in R, m \in M$. An R - H -module M is called an R - H -irreducible module if there are no non-trivial R - H -submodules in M and $RM \neq 0$. An algebra homomorphism $\psi : R \rightarrow R'$ is called an H -homomorphism if $\psi(h \cdot a) = h \cdot \psi(a)$ for any $h \in H, a \in R$. Let r_b, r_j, r_l, r_{bm} denote the Baer radical, the Jacobson radical, the locally nilpotent radical, the Brown-MacCoy radical of algebras respectively. Let $I \triangleleft_H R$ denote that I is an H -ideal of R .

11.1 The H -special radicals for H -module algebras

J.R. Fisher [48] built up the general theory of H -radicals for H -module algebras. We can easily give the definitions of the H -upper radical and the H -lower radical for H -module algebras as in [116]. In this section, we obtain some properties of H -special radicals for H -module algebras.

Lemma 11.1.1. (1) If R is an H -module algebra and E is a non-empty subset of R , then $(E) = H \cdot E + R(H \cdot E) + (H \cdot E)R + R(H \cdot E)R$, where (E) denotes the H -ideal generated by E in R .

(2) If B is an H -ideal of R and C is an H -ideal of B , then $(C)^3 \subseteq C$, where (C) denotes the H -ideal generated by C in R .

Proof. It is trivial. $\square$

Proposition 11.1.2. (1) R is H -semiprime iff $(H \cdot a)R(H \cdot a) = 0$ always implies $a = 0$ for any $a \in R$.

(2) R is H -prime iff $(H \cdot a)R(H \cdot b) = 0$ always implies $a = 0$ or $b = 0$ for any $a, b \in R$.

Proof. If R is an H -prime module algebra and $(H \cdot a)R(H \cdot b) = 0$ for $a, b \in R$, then $(a)^2(b)^2 = 0$, where (a) and (b) are the H -ideals generated by a and b in R respectively. Since R is H -prime, $(a) = 0$ or $(b) = 0$. Conversely, if B and C are H -ideals of R and $BC = 0$, then

$(H \cdot a)R(H \cdot b) = 0$ and $a = 0$ or $b = 0$ for any $a \in B, b \in C$, which implies that $B = 0$ or $C = 0$, i.e. R is an H -prime module algebra.

Similarly, part (1) holds. $\square$

Proposition 11.1.3. *If $I \triangleleft_H R$ and I is an H -semiprime module algebra, then*

(1) $I \cap I^ = 0$; (2) $I_r = I_l = I^*$; (3) $I^* \triangleleft_H R$, where $I_r = \{a \in R \mid I(H \cdot a) = 0\}$, $I_l = \{a \in R \mid (H \cdot a)I = 0\}$, $I^* = \{a \in R \mid (H \cdot a)I = 0 = I(H \cdot a)\}$.*

Proof . For any $x \in I^* \cap I$, we have that $I(H \cdot x) = 0$ and $(H \cdot x)I(H \cdot x) = 0$. Since I is an H -semiprime module algebra, $x = 0$, i.e. $I \cap I^* = 0$.

To show $I^* = I_r$, we only need to show that $(H \cdot x)I = 0$ for any $x \in I_r$. For any $y \in I, h \in H$, let $z = (h \cdot x)y$. It is clear that $(H \cdot z)I(H \cdot z) = 0$. Since I is an H -semiprime module algebra, $z = 0$, i.e. $(H \cdot x)I = 0$. Thus $I^* = I_r$. Similarly, we can show that $I_l = I^*$.

Obviously, I^* is an ideal of R . For any $x \in I^*, h \in H$, we have $(H \cdot (h \cdot x))I = 0$. Thus $h \cdot x \in I^*$ by part (2), i.e. I^* is an H -ideal of R . $\square$

Definition 11.1.4. $\mathcal{K}$ is called an H -(weakly)special class if

(S1) $\mathcal{K}$ consists of H -(semiprime)prime module algebras.

(S2) For any $R \in \mathcal{K}$, if $0 \neq I \triangleleft_H R$ then $I \in \mathcal{K}$.

(S3) If R is an H -module algebra and $B \triangleleft_H R$ with $B \in \mathcal{K}$, then $R/B^* \in \mathcal{K}$, where $B^* = \{a \in R \mid (H \cdot a)B = 0 = B(H \cdot a)\}$.

It is clear that (S3) may be replaced by one of the following conditions:

(S3') If B is an essential H -ideal of R (i.e. $B \cap I \neq 0$ for any non-zero H -ideal I of R) and $B \in \mathcal{K}$, then $R \in \mathcal{K}$.

(S3'') If there exists an H -ideal B of R with $B^* = 0$ and $B \in \mathcal{K}$, then $R \in \mathcal{K}$.

It is easy to check that if $\mathcal{K}$ is an H -special class, then $\mathcal{K}$ is an H -weakly special class.

Theorem 11.1.5. *If $\mathcal{K}$ is an H -weakly special class, then $r^{\mathcal{K}}(R) = \cap \{I \triangleleft_H R \mid R/I \in \mathcal{K}\}$, where $r^{\mathcal{K}}$ denotes the H -upper radical determined by $\mathcal{K}$.*

Proof. If I is a non-zero H -ideal of R and $I \in \mathcal{K}$, then $R/I^* \in \mathcal{K}$ by (S3) in Definition 11.1.4 and $I \not\subseteq I^*$ by Proposition 11.1.3. Consequently, it follows from [48, Proposition 5] that

$$r^{\mathcal{K}}(R) = \cap \{I \mid I \text{ is an } H\text{-ideal of } R \text{ and } R/I \in \mathcal{K}\} . \quad \square$$

Definition 11.1.6. *If r is a hereditary H -radical (i.e. if R is an r - H -module algebra and B is an H -ideal of R , then so is B) and any nilpotent H -module algebra is an r - H -module algebra, then r is called a supernilpotent H -radical.*

Proposition 11.1.7. *r is a supernilpotent H -radical, then r is H -strongly hereditary, i.e. $r(I) = r(R) \cap I$ for any $I \triangleleft_H R$.*

Proof. It follows from [48, Proposition 4] . $\square$

Theorem 11.1.8. *If $\mathcal{K}$ is an H -weakly special class, then $r^{\mathcal{K}}$ is a supernilpotent H -radical.*

Proof. Let $r = r^{\mathcal{K}}$. Since every non-zero H -homomorphic image R' of a nilpotent H -module algebra R is nilpotent and is not H -semiprime, we have that R is an r - H -module algebra by Theorem 11.1.5. It remains to show that any H -ideal I of r - H -module algebra R is an r - H -ideal. If I is not an r - H -module algebra, then there exists an H -ideal J of I such that $0 \neq I/J \in \mathcal{K}$. By (S3), $(R/J)/(I/J)^* \in \mathcal{K}$. Let $Q = \{x \in R \mid (H \cdot x)I \subseteq J \text{ and } I(H \cdot x) \subseteq J\}$. It is clear that J and Q are H -ideals of R and $Q/J = (I/J)^*$. Since $R/Q \cong (R/J)/(Q/J) = (R/J)/(I/J)^*$ and R/Q is an r - H -module algebra, we have $(R/J)/(I/J)^*$ is an r - H -module algebra. Thus $R/Q = 0$ and $I^2 \subseteq J$, which contradicts that I/J is a non-zero H -semiprime module algebra. Thus I is an r - H -ideal. $\square$

Proposition 11.1.9. *R is H -semiprime iff for any $0 \neq a \in R$, there exists an H - m -sequence $\{a_n\}$ in R with $a_{7.1.1} = a$ such that $a_n \neq 0$ for all n .*

Proof. If R is H -semiprime, then for any $0 \neq a \in R$, there exist $b_1 \in R$, h_1 and $h'_1 \in H$ such that $0 \neq a_2 = (h_1 \cdot a_1)b_1(h'_1 \cdot a_1) \in (H \cdot a_1)R(H \cdot a_1)$ by Proposition 11.1.2, where $a_1 = a$. Similarly, for $0 \neq a_2 \in R$, there exist $b_2 \in R$ and h_2 and $h'_2 \in H$ such that $0 \neq a_3 = (h_2 \cdot a_2)b_2(h'_2 \cdot a_2) \in (H \cdot a_2)R(H \cdot a_2)$, which implies that there exists an H - m -sequence $\{a_n\}$ such that $a_n \neq 0$ for any natural number n . Conversely, it is trivial. $\square$

11.2 H -Baer radical

In this section, we give the characterization of H -Baer radical(H -prime radical) in many ways.

Theorem 11.2.1. *We define a property r_{Hb} for H -module algebras as follows: R is an r_{Hb} - H -module algebra iff every non-zero H -homomorphic image of R contains a non-zero nilpotent H -ideal; then r_{Hb} is an H -radical property.*

Proof. It is clear that every H -homomorphic image of r_{Hb} - H -module algebra is an r_{Hb} - H -module algebra. If every non-zero H -homomorphic image B of H -module algebra R contains a non-zero r_{Hb} - H -ideal I , then I contains a non-zero nilpotent H -ideal J . It is clear that (J) is a non-zero nilpotent H -ideal of B , where (J) denotes the H -ideal generated by J in B . Thus R is an r_{Hb} - H -module algebra. Consequently, r_{Hb} is an H -radical property. $\square$

r_{Hb} is called H -prime radical or H -Baer radical.

Theorem 11.2.2. *Let*

$$\mathcal{E} = \{R \mid R \text{ is a nilpotent } H\text{-module algebra}\},$$

then $r_{\mathcal{E}} = r_{Hb}$, where $r_{\mathcal{E}}$ denotes the H -lower radical determined by $\mathcal{E}$.

Proof. If R is an r_{Hb} - H -module algebra, then every non-zero H -homomorphic image B of R contains a non-zero nilpotent H -ideal I . By the definition of the lower H -radical, I is an $r_{\mathcal{E}}$ - H -module algebra. Consequently, R is an $r_{\mathcal{E}}$ - H -module algebra. Conversely, since every nilpotent H -module algebra is an r_{Hb} - H -module algebra, $r_{\mathcal{E}} \leq r_{Hb}$. $\square$

Proposition 11.2.3. *R is H -semiprime if and only if $r_{Hb}(R) = 0$.*

Proof. If R is H -semiprime with $r_{Hb}(R) \neq 0$, then there exists a non-zero nilpotent H -ideal I of $r_{Hb}(R)$. It is clear that H -ideal (I) , which the H -ideal generated by I in R , is a non-zero nilpotent H -ideal of R . This contradicts that R is H -semiprime. Thus $r_{Hb}(R) = 0$. Conversely, if R is an H -module algebra with $r_{Hb}(R) = 0$ and there exists a non-zero nilpotent H -ideal I of R , then $I \subseteq r_{Hb}(R)$. We get a contradiction. Thus R is H -semiprime if $r_{Hb}(R) = 0$. $\square$

Theorem 11.2.4. *If $\mathcal{K} = \{R \mid R \text{ is an } H\text{-prime module algebra}\}$, then $\mathcal{K}$ is an H -special class and $r_{Hb} = r^{\mathcal{K}}$.*

Proof. Obviously, (S1) holds. If I is a non-zero H -ideal of an H -prime module algebra R and $BC = 0$ for H -ideals B and C of I , then $(B)^3(C)^3 = 0$ where (B) and (C) denote the H -ideals generated by B and C in R respectively. Since R is H -prime, $(B) = 0$ or $(C) = 0$, i.e. $B = 0$ or $C = 0$. Consequently, (S2) holds. Now we shows that (S3) holds. Let B be an H -prime module algebra and be an H -ideal of R . If $JI \subseteq B^*$ for H -ideals I and J of R , then $(BJ)(IB) = 0$, where $B^* = \{x \in R \mid (H \cdot x)B = 0 = B(H \cdot x)\}$. Since B is an H -prime module algebra, $BJ = 0$ or $IB = 0$. Considering I and J are H -ideals, we have that $B(H \cdot J) = 0$ or $(H \cdot I)B = 0$. By Proposition 11.1.3, $J \subseteq B^*$ or $I \subseteq B^*$, which implies that R/B^* is an H -prime module algebra. Consequently, (S3) holds and so $\mathcal{K}$ is an H -special class.

Next we show that $r_{Hb} = r^{\mathcal{K}}$. By Proposition 11.1.5, $r^{\mathcal{K}}(R) = \cap\{I \mid I \text{ is an } H\text{-ideal of } R \text{ and } R/I \in \mathcal{K}\}$. If R is a nilpotent H -module algebra, then R is an $r^{\mathcal{K}}$ - H -module algebra. It follows from Theorem 11.2.2 that $r_{Hb} \leq r^{\mathcal{K}}$. Conversely, if $r_{Hb}(R) = 0$, then R is an H -semiprime module algebra by Proposition 11.2.3. For any $0 \neq a \in R$, there exist $b_1 \in R$, $h_1, h'_1 \in H$ such that $0 \neq a_2 = (h_1 \cdot a_1)b_1(h'_1 \cdot a_1) \in (H \cdot a_1)R(H \cdot a_1)$, where $a_1 = a$. Similarly, for $0 \neq a_2 \in R$, there exist $b_2 \in R$ and $h_2, h'_2 \in H$ such that $0 \neq a_3 = (h_2 \cdot a_2)b_2(h'_2 \cdot a_2) \in (H \cdot a_2)R(H \cdot a_2)$. Thus there exists an H - m -sequence $\{a_n\}$ such that $a_n \neq 0$ for any natural number n . Let

$$\mathcal{F} = \{I \mid I \text{ is an } H\text{-ideal of } R \text{ and } I \cap \{a_1, a_2, \dots\} = \emptyset\}.$$

By Zorn's Lemma, there exists a maximal element P in $\mathcal{F}$. If I and J are H -ideals of R and $I \not\subseteq P$ and $J \not\subseteq P$, then there exist natural numbers n and m such that $a_n \in I$ and $a_m \in J$. Since $0 \neq a_{n+m+1} = (h_{n+m} \cdot a_{n+m})b_{n+m}(h'_{n+m} \cdot a_{n+m}) \in IJ$, which implies that $IJ \not\subseteq P$ and so P is an H -prime ideal of R . Obviously, $a \notin P$, which implies that $a \notin r^{\mathcal{K}}(R)$ and $r^{\mathcal{K}}(R) = 0$. Consequently, $r^{\mathcal{K}} = r_{Hb}$. $\square$

Theorem 11.2.5. $r_{Hb}(R) = W_H(R)$.

Proof. If $0 \neq a \notin W_H(R)$, then there exists an H -prime ideal P such that $a \notin P$ by the proof of Theorem 11.2.4. Thus $a \notin r_{Hb}(R)$, which implies that $r_{Hb}(R) \subseteq W_H(R)$. Conversely, for any $x \in W_H(R)$, let $\bar{R} = R/r_{Hb}(R)$. Since $r_{Hb}(\bar{R}) = 0$, $\bar{R}$ is an H -semiprime module algebra by Proposition 11.2.3. By the proof of Theorem 11.2.4, $W_H(\bar{R}) = 0$. For an H - m -sequence $\{\bar{a}_n\}$ with $\bar{a}_1 = \bar{x}$ in $\bar{R}$, there exist $\bar{b}_n \in \bar{R}$ and $h_n, h'_n \in H$ such that

$$\bar{a}_{n+1} = (h_n \cdot \bar{a}_n)\bar{b}_n(h'_n \cdot \bar{a}_n)$$

for any natural number n . Thus there exists $a'_n \in R$ such that $a'_1 = x$ and $a'_{n+1} = (h_n \cdot a'_n)b_n(h'_n \cdot a'_n)$ for any natural number n . Since $\{a'_n\}$ is an H - m -sequence with $a'_1 = x$ in R , there exists a natural number k such that $a'_k = 0$. It is easy to show that $\bar{a}_n = \bar{a}'_n$ for any natural number n by induction. Thus $\bar{a}_k = 0$ and $\bar{x} \in W_H(\bar{R})$. Considering $W_H(\bar{R}) = 0$, we have $x \in r_{Hb}(R)$ and $W_H(R) \subseteq r_{Hb}(R)$. Therefore $W_H(R) = r_{Hb}(R)$. $\square$

Definition 11.2.6. We define an H -ideal N_α in H -module algebra R for every ordinal number α as follows:

(i) $N_0 = 0$.

Let us assume that N_α is already defined for $\alpha \prec \beta$.

(ii) If $\beta = \alpha + 1$, N_β/N_α is the sum of all nilpotent H -ideals of R/N_α

(iii) If β is a limit ordinal number, $N_\beta = \sum_{\alpha \prec \beta} N_\alpha$.

By set theory, there exists an ordinal number τ such that $N_\tau = N_{\tau+1}$.

Theorem 11.2.7. $N_\tau = r_{Hb}(R) = \cap\{I \mid I \text{ is an } H\text{-semiprime ideal of } R\}$.

Proof. Let $D = \cap\{I \mid I \text{ is an } H\text{-semiprime ideal of } R\}$. Since R/N_τ has not any non-zero nilpotent H -ideal, we have that $r_{Hb}(R) \subseteq N_\tau$ by Proposition 11.2.3. Obviously, $D \subseteq r_{Hb}(R)$. Using transfinite induction, we can show that $N_\alpha \subseteq I$ for every H -semiprime ideal I of R and every ordinal number α (see the proof of [127, Theorem 3.7]). Thus $N_\tau \subseteq D$, which completes the proof. $\square$

Definition 11.2.8. Let $\emptyset \neq L \subseteq H$. An H - m -sequence $\{a_n\}$ in R is called an L - m -sequence with beginning a if $a_{7.1.1} = a$ and $a_{n+1} = (h_n \cdot a_n)b_n(h'_n \cdot a_n)$ such that $h_n, h'_n \in L$ for all n . For every L - m -sequence $\{a_n\}$ with $a_{7.1.1} = a$, there exists a natural number k such that $a_k = 0$, then R is called an L - m -nilpotent element, written as $W_L(R) = \{a \in R \mid a \text{ is an } L\text{-}m\text{-nilpotent element}\}$.

Similarly, we have

Proposition 11.2.9. If $L \subseteq H$ and $H = kL$, then

(i) R is H -semiprime iff $(L.a)R(L.a) = 0$ always implies $a = 0$ for any $a \in R$.

(ii) R is H -prime iff $(L.a)R(L.b) = 0$ always implies $a = 0$ or $b = 0$ for any $a, b \in R$.

(iii) R is H -semiprime if and only if for any $0 \neq a \in R$, there exists an L - m -sequence $\{a_n\}$ with $a_1 = a$ such that $a_n \neq 0$ for all n .

(iv) $W_H(R) = W_L(R)$.

11.3 The H -module theoretical characterization of H -special radicals

If V is an algebra over k with unit and $x \otimes 1_V = 0$ always implies that $x = 0$ for any right k -module M and for any $x \in M$, then V is called a faithful algebra to tensor. For example, if k is a field, then V is faithful to tensor for any algebra V with unit.

In this section, we need to add the following condition: H is faithful to tensor.

We shall characterize H -Baer radical r_{Hb} , H -locally nil radical r_{HL} , H -Jacobson radical r_H and H -Brown-McCoy radical r_{Hbm} by R - H -modules.

We can view every H -module algebra R as a sub-algebra of $R\#H$ since H is faithful to tensor. By computation, we have that

$$h \cdot a = \sum (1\#h_1)a(1\#S(h_2))$$

for any $h \in H, a \in R$, where S is the antipode of H .

Definition 11.3.1. An R - H -module M is called an R - H -prime module if for M the following conditions are fulfilled:

- (i) $RM \neq 0$;
- (ii) If x is an element of M and I is an H -ideal of R , then $I(Hx) = 0$ always implies $x = 0$ or $I \subseteq (0 : M)_R$, where $(0 : M)_R = \{a \in R \mid aM = 0\}$.

Definition 11.3.2. We associate to every H -module algebra R a class $\mathcal{M}_R$ of R - H -modules. Then the class $\mathcal{M} = \cup \mathcal{M}_R$ is called an H -special class of modules if the following conditions are fulfilled:

- (M1) If $M \in \mathcal{M}_R$, then M is an R - H -prime module.
- (M2) If I is an H -ideal of R and $M \in \mathcal{M}_I$, then $IM \in \mathcal{M}_R$.
- (M3) If $M \in \mathcal{M}_R$ and I is an H -ideal of R with $IM \neq 0$, then $M \in \mathcal{M}_I$.
- (M4) Let I be an H -ideal of R and $\bar{R} = R/I$. If $M \in \mathcal{M}_R$ and $I \subseteq (0 : M)_R$, then $M \in \mathcal{M}_{\bar{R}}$. Conversely, if $M \in \mathcal{M}_{\bar{R}}$, then $M \in \mathcal{M}_R$.

Let $\mathcal{M}(R)$ denote $\cap \{(0 : M)_R \mid M \in \mathcal{M}_R\}$, or R when $\mathcal{M}_R = \emptyset$.

Lemma 11.3.3. (1) If M is an R - H -module, then M is an $R\#H$ -module. In this case, $(0 : M)_{R\#H} \cap R = (0 : M)_R$ and $(0 : M)_R$ is an H -ideal of R ;

(2) R is a non-zero H -prime module algebra iff there exists a faithful R - H -prime module M ;

(3) Let I be an H -ideal of R and $\bar{R} = R/I$. If M is an R - H -(resp. prime, irreducible) module and $I \subseteq (0 : M)_R$, then M is an $\bar{R}$ - H -(resp. prime, irreducible) module (defined by $h \cdot (a + I) = h \cdot a$ and $(a + I)x = ax$). Conversely, if M is an $\bar{R}$ - H -(resp. prime irreducible) module, then M is an R - H -(resp. prime, irreducible) module (defined by $h \cdot a = h \cdot (a + I)$ and $ax = (a + I)x$). In the both cases, it is always true that $R/(0 : M)_R \cong \bar{R}/(0 : M)_{\bar{R}}$;

(4) I is an H -prime ideal of R with $I \neq R$ iff there exists an R - H -prime module M such that $I = (0 : M)_R$;

(5) If I is an H -ideal of R and M is an I - H -prime module, then IM is an R - H -prime module with $(0 : M)_I = (0 : IM)_R \cap I$;

(6) If M is an R - H -prime module and I is an H -ideal of R with $IM \neq 0$, then M is an I - H -prime module;

(7) If R is an H -semiprime module algebra with one side unit, then R has a unit.

Proof. (1) Obviously, $(0 : M)_R = (0 : M)_{R\#H} \cap R$. For any $h \in H, a \in (0 : M)_R$, we see that $(h \cdot a)M = \sum (1\#h_1)a(1\#S(h_2))M \subseteq \sum (1\#h_1)aM = 0$ for any $h \in H, a \in R$. Thus $h \cdot a \in (0 : M)_R$, which implies $(0 : M)_R$ is an H -ideal of R .

(2) If R is an H -prime module algebra, view $M = R$ as an R - H -module. Obviously, M is faithful. If $I(H \cdot x) = 0$ for $0 \neq x \in M$ and an H -ideal I of R , then $I(x) = 0$ and $I = 0$, where (x) denotes the H -ideal generated by x in R . Consequently, M is a faithful R - H -prime module. Conversely, let M be a faithful R - H -prime module. If $IJ = 0$ for two H -ideals I and J of R with $J \neq 0$, then $JM \neq 0$ and there exists $0 \neq x \in JM$ such that $I(Hx) = 0$. Since M is a faithful R - H -prime module, $I = 0$. Consequently, R is H -prime.

(3) If M is an R - H -module, then it is clear that M is a (left) $\overline{R}$ -module and $h(\overline{a}x) = h(ax) = \sum (h_1 \cdot a)(h_2x) = \sum \overline{(h_1 \cdot a)}(h_2x) = \sum (h_1 \cdot \overline{a})(h_2x)$ for any $h \in H$, $a \in R$ and $x \in M$. Thus M is an $\overline{R}$ - H -module. Conversely, if M is an $\overline{R}$ - H -module, then M is an (left) R -module and

$$h(ax) = h(\overline{a}x) = \sum (h_1 \cdot \overline{a})(h_2x) = \sum \overline{h_1 \cdot a}(h_2x) = \sum (h_1 \cdot a)(h_2x)$$

for any $h \in H$, $a \in R$ and $x \in M$. This shows that M is an R - H -module.

Let M be an R - H -prime module and I be an H -ideal of R with $I \subseteq (0 : M)_R$. If $\overline{J}(Hx) = 0$ for $0 \neq x \in M$ and an H -ideal J of R , then $J(Hx) = 0$ and $J \subseteq (0 : M)_R$. This shows that $\overline{J} \subseteq (0 : M)_{\overline{R}}$. Thus M is an R - H -prime module. Similarly, we can show the other assert.

(4) If I is an H -prime ideal of R with $R \neq I$, then $\overline{R} = R/I$ is an H -prime module algebra. By Part (2), there exists a faithful $\overline{R}$ - H -prime module M . By part (3), M is an R - H -prime module with $(0 : M)_R = I$. Conversely, if there exists a R - H -prime M with $I = (0 : M)_R$, then M is a faithful $\overline{R}$ - H -prime module by part (3) and I is an H -prime ideal of R by part (2).

(5) First, we show that IM is an R -module. We define

$$a(\sum_i a_i x_i) = \sum_i (aa_i)x_i \quad (11.1)$$

for any $a \in R$ and $\sum_i a_i x_i \in IM$, where $a_i \in I$ and $x_i \in M$. If $\sum_i a_i x_i = \sum_i a'_i x'_i$ with $a_i, a'_i \in R$, $x_i, x'_i \in M$, let $y = \sum_i (aa_i)x_i - \sum_i (aa'_i)x'_i$. For any $b \in I$ and $h \in H$, we see that

$$\begin{aligned} b(hy) &= \sum_i b\{h[(aa_i)x_i - (aa'_i)x'_i]\} \\ &= \sum_i \sum_{(h)} b\{[(h_1 \cdot (aa_i))(h_2x_i) - [h_1 \cdot (aa'_i)](h_2x'_i)]\} \\ &= \sum_{(h)} \sum_i \{b[(h_1 \cdot a)(h_2 \cdot a_i)](h_3x_i) - b[(h_1 \cdot a)(h_2 \cdot a'_i)](h_3x'_i)\} \\ &= \sum_{(h)} \sum_i b(h_1 \cdot a)[h_2(a_i x_i) - h_2(a'_i x'_i)] \\ &= \sum_{(h)} b(h_1 \cdot a)h_2 \sum_i [a_i x_i - a'_i x'_i] = 0. \end{aligned}$$

Thus $I(Hy) = 0$. Since M is an I - H -prime module and $IM \neq 0$, we have that $y = 0$. Thus this

definition in (11.1) is well-defined. It is easy to check that IM is an R -module. We see that

$$\begin{aligned}
h(a \sum_i a_i x_i) &= \sum_i h[(aa_i)x_i] \\
&= \sum_i \sum_h [h_1 \cdot (aa_i)][h_2 x_i] \\
&= \sum_i \sum_h [(h_1 \cdot a)(h_2 \cdot a_i)](h_3 x) \\
&= \sum_h (h_1 \cdot a) \sum_i (h_2 \cdot a_i)(h_3 x_i) \\
&= \sum_h (h_1 \cdot a)[h_2 \sum_i (a_i x_i)]
\end{aligned}$$

for any $h \in H$ and $\sum_i a_i x_i \in IM$. Thus IM is an R - H -module.

Next, we show that $(0 : M)_I = (0 : IM)_R \cap I$. If $a \in (0 : M)_I$, then $aM = 0$ and $aIM = 0$, i.e. $a \in (0 : IM)_A \cap I$. Conversely, if $a \in (0 : IM)_R \cap I$, then $aIM = 0$. By part (1), $(0 : IM)_R$ is an H -ideal of R . Thus $(H \cdot a)IM = 0$ and $(H \cdot a)I \subseteq (0 : M)_I$. Since $(0 : M)_I$ is an H -prime ideal of I by part (4), $a \in (0 : M)_I$. Consequently, $(0 : M)_I = (0 : IM)_R \cap I$.

Finally, we show that IM is an R - H -prime module. If $RIM = 0$, then $RI \subseteq (0 : M)_R$ and $I \subseteq (0 : M)_R$, which contradicts that M is an I - H -prime module. Thus $RIM \neq 0$. If $J(Hx) = 0$ for $0 \neq x \in IM$ and an H -ideal J of R , then $JI(Hx) \subseteq J(Hx) = 0$. Since M is an I - H -prime module, $JI \subseteq (0 : M)_I$ and $J(IM) = 0$. Consequently, IM is an R - H -prime module.

(6) Obviously, M is an I - H -module. If $J(Hx) = 0$ for $0 \neq x \in M$ and an H -ideal J of I , then $(J)^3(Hx) = 0$ and $(J)^3 \subseteq (0 : M)_R$, where (J) denotes the H -ideal generated by J in R . Since $(0 : M)_R$ is an H -prime ideal of R , $(J) \subseteq (0 : M)_R$ and $J \subseteq (0 : M)_I$. Consequently, M is an I - H -prime module.

(7) We can assume that u is a right unit of R . We see that

$$(h \cdot (au - a))b = \sum (1 \# h_1)(au - a)(1 \# S(h_2))b = 0$$

for any $a, b \in R, h \in H$. Therefore $(H \cdot (au - a))R = 0$ and $au = a$, which implies that R has a unit. $\square$

Theorem 11.3.4. (1) If $\mathcal{M}$ is an H -special class of modules and $\mathcal{K} = \{ R \mid \text{there exists a faithful } R\text{-}H\text{-module } M \in \mathcal{M}_R \}$, then $\mathcal{K}$ is an H -special class and $r^{\mathcal{K}}(R) = \mathcal{M}(R)$.

(2) If $\mathcal{K}$ is an H -special class and $\mathcal{M}_R = \{ M \mid M \text{ is an } R\text{-}H\text{-prime module and } R/(0 : M)_R \in \mathcal{K} \}$, then $\mathcal{M} = \cup \mathcal{M}_R$ is an H -special class of modules and $r^{\mathcal{K}}(R) = \mathcal{M}(R)$.

Proof. (1) By Lemma 11.3.3(2), (S1) is satisfied. If I is a non-zero H -ideal of R and $R \in \mathcal{K}$, then there exists a faithful R - H -prime module $M \in \mathcal{M}_R$. Since M is faithful, $IM \neq 0$ and $M \in \mathcal{M}_I$ with $(0 : M)_I = (0 : M)_R \cap I = 0$ by (M3). Thus $I \in \mathcal{K}$ and (S2) is satisfied. Now we show that (S3) holds. If I is an H -ideal of R with $I \in \mathcal{K}$, then there exists a faithful I - H -prime module $M \in \mathcal{M}_I$. By (M2) and Lemma 11.3.3(5), $IM \in \mathcal{M}_R$ and $0 = (0 : M)_I = (0 : IM)_R \cap I$. Thus $(0 : IM)_R \subseteq I^*$. Obviously, $I^* \subseteq (0 : IM)_R$. Thus $I^* = (0 : IM)_R$. Using (M4), we have

that $IM \in \mathcal{M}_{\overline{R}}$ and IM is a faithful $\overline{R}$ - H -module with $\overline{R} = R/I^*$. Thus $R/I^* \in \mathcal{K}$. Therefore $\mathcal{K}$ is an H -special class.

It is clear that

$$\{I \mid I \text{ is an } H\text{-ideal of } R \text{ and } R/I \in \mathcal{K}\} = \{(0 : M)_R \mid M \in \mathcal{M}_R\}.$$

Thus $r^{\mathcal{K}}(R) = \mathcal{M}(R)$.

(2) It is clear that (M1) is satisfied. If I is an H -ideal of R with $M \in \mathcal{M}_I$, then M is an I - H -prime module with $I/(0 : M)_I \in \mathcal{K}$. By Lemma 11.3.3(5), IM is an R - H -prime module with $(0 : M)_I = (0 : IM)_R \cap I$. It is clear that

$$(0 : IM)_R = \{a \in R \mid (H \cdot a)I \subseteq (0 : M)_I \text{ and } I(H \cdot a) \subseteq (0 : M)_I\}$$

and

$$(0 : IM)_R / (0 : M)_I = (I / (0 : M)_I)^*.$$

Thus $R / (0 : IM)_R \cong (R / (0 : M)_I) / ((0 : IM)_R / (0 : M)_I) = (R / (0 : M)_I) / (I / (0 : M)_I)^* \in \mathcal{K}$, which implies that $IM \in \mathcal{M}_R$ and (M2) holds. Let $M \in \mathcal{M}_R$ and I be an H -ideal of R with $IM \neq 0$. By Lemma 11.3.3(6), M is an I - H -prime module and $I / (0 : M)_I = I / ((0 : M)_R \cap I) \cong (I + (0 : M)_R) / (0 : M)_R$. Since $R / (0 : M)_R \in \mathcal{K}$, $I / (0 : M)_I \in \mathcal{K}$ and $M \in \mathcal{M}_I$. Thus (M3) holds. It follows from Lemma 11.3.3(3) that (M4) holds.

It is clear that

$$\{I \mid I \text{ is an } H\text{-ideal of } R \text{ and } 0 \neq R/I \in \mathcal{K}\} = \{(0 : M)_R \mid M \in \mathcal{M}_R\}.$$

Thus $r^{\mathcal{K}}(R) = \mathcal{M}(R)$. $\square$

Theorem 11.3.5. *Let $\mathcal{M}_R = \{ M \mid M \text{ is an } R\text{-}H\text{-prime module} \}$ for any H -module algebra R and $\mathcal{M} = \cup \mathcal{M}_R$. Then $\mathcal{M}$ is an H -special class of modules and $\mathcal{M}(R) = r_{Hb}(R)$.*

Proof. It follows from Lemma 11.3.3(3)(5)(6) that $\mathcal{M}$ is an H -special class of modules. By Lemma 11.3.3(2),

$$\{R \mid R \text{ is an } H\text{-prime module algebra with } R \neq 0\} =$$

$$\{R \mid \text{there exists a faithful } R\text{-}H\text{-prime module}\}.$$

Thus $r_{Hb}(R) = \mathcal{M}(R)$ by Theorem 11.2.4(1). $\square$

Theorem 11.3.6. *Let $\mathcal{M}_R = \{ M \mid M \text{ is an } R\text{-}H\text{-irreducible module} \}$ for any H -module algebra R and $\mathcal{M} = \cup \mathcal{M}_R$. Then $\mathcal{M}$ is an H -special class of modules and $\mathcal{M}(R) = r_{Hj}(R)$, where r_{Hj} is the H -Jacobson radical of R defined in [48].*

Proof. If M is an R - H -irreducible module and $J(Hx) = 0$ for $0 \neq x \in M$ and an H -ideal J of R , let $N = \{m \in M \mid J(Hm) = 0\}$. Since $J(h(am)) = J(\sum_h (h_1 \cdot a)(h_2 m)) = 0$, $am \in N$ for any $m \in N, h \in H, a \in R$, we have that N is an R -submodule of M . Obviously, N is an

H -submodule of M . Thus N is an R - H -submodule of M . Since $N \neq 0$, we have that $N = M$ and $JM = 0$, i.e. $J \subseteq (0 : M)_R$. Thus M is an R - H -prime module and (M1) is satisfied. If M is an I - H -irreducible module and I is an H -ideal, then IM is an R - H -module. If N is an R - H -submodule of IM , then N is also an I - H -submodule of M , which implies that $N = 0$ or $N = M$. Thus (M2) is satisfied. If M is an R - H -irreducible module and I is an H -ideal of R with $IM \neq 0$, then $IM = M$. If N is a non-zero I - H -submodule of M , then IN is an R - H -submodule of M by Lemma 11.3.3(5) and $IN = 0$ or $IN = M$. If $IN = 0$, then $I \subseteq (0 : M)_R$ by the above proof and $IM = 0$. We get a contradiction. If $IN = M$, then $N = M$. Thus M is an I - H -irreducible module and (M3) is satisfied.

It follows from Lemma 11.3.3(3) that (M4) holds. By Theorem 11.3.4(1), $\mathcal{M}(R) = r_{Hj}(R)$.

□

J.R. Fisher [48, Proposition 2] constructed an H -radical r_H by a common hereditary radical r for algebras, i.e. $r_H(R) = (r(R) : H) = \{a \in R \mid h \cdot a \in r(R) \text{ for any } h \in H\}$. Thus we can get H -radicals $r_{bH}, r_{lH}, r_{jH}, r_{bmH}$.

Definition 11.3.7. An R - H -module M is called an R - H -BM-module, if for M the following conditions are fulfilled:

- (i) $RM \neq 0$;
- (ii) If I is an H -ideal of R and $I \not\subseteq (0 : M)_R$, then there exists an element $u \in I$ such that $m = um$ for all $m \in M$.

Theorem 11.3.8. Let $\mathcal{M}_R = \{ M \mid M \text{ is an } R\text{-}H\text{-BM-module} \}$ for every H -module algebra R and $\mathcal{M} = \cup \mathcal{M}_R$. Then $\mathcal{M}$ is an H -special class of modules.

Proof. It is clear that M satisfies (M_1) and (M_4) . To prove (M_2) we exhibit: if $I \triangleleft_H R$ and $M \in \mathcal{M}_I$, then M is an I - H -prime module and IM is an R - H -prime module. If J is an H -ideal of R with $J \not\subseteq (0 : M)_R$, then JI is an H -ideal of I with $JI \not\subseteq (0 : M)_I$. Thus there exists an element $u \in JI \subseteq J$ such that $um = m$ for every $m \in M$. Hence $IM \in \mathcal{M}_R$.

To prove (M_3) , we exhibit: if $M \in \mathcal{M}_R$ and I is an H -ideal of R with $IM \neq 0$. If J is an H -ideal of I with $J \not\subseteq (0 : M)_I$, then $(J) \not\subseteq (0 : M)_R$, where (J) is the H -ideal generated by J in R . Thus there exists an element $u \in (J)$ such that $um = m$ for every $m \in M$. Moreover,

$$m = um = uum = uuum = u^3m$$

and $u^3 \in J$. Thus $M \in \mathcal{M}_I$. □

Proposition 11.3.9. If M is an R - H -BM-module, then $R/(0 : M)_R$ is an H -simple module algebra with unit.

Proof. Let I be any H -ideal of R with $I \not\subseteq (0 : M)_R$. Since M is an R - H -BM-module, there exists an element $u \in I$ such that $uam = am$ for every $m \in M, a \in R$. It follows that $a - ua \in (0 : M)_R$, whence $R = I + (0 : M)_R$. Thus $(0 : M)_R$ is a maximal H -ideal of R . Therefore $R/(0 : M)_R$ is an H -simple module algebra.

Next we shall show that $R/(0 : M)_R$ has a unit. Now $R \not\subseteq (0 : M)_R$, since $RM \neq 0$. By the above proof, there exists an element $u \in R$ such that $a - ua \in (0 : M)_R$ for any $a \in R$. Hence $R/(0 : M)_R$ has a left unit. Furthermore, by Lemma 11.3.7 (7) it has a unity element. □

Proposition 11.3.10. *If R is an H -simple-module algebra with unit, then there exists a faithful R - H -BM-module.*

Proof. Let $M = R$. It is clear that M is a faithful R - H -BM-module. $\square$

Theorem 11.3.11. *Let $\mathcal{M}_R = \{ M \mid M \text{ is an } R\text{-}H\text{-BM-module} \}$ for every H -module algebra R and $\mathcal{M} = \cup \mathcal{M}_R$. Then $r_{Hbm}(R) = \mathcal{M}(R)$, where r_{Hbm} denotes the H -upper radical determined by $\{R \mid R \text{ is an } H\text{-simple module algebra with unit} \}$.*

Proof. By Theorem 11.3.8, $\mathcal{M}$ is an H -special class of modules. Let

$$\mathcal{K} = \{R \mid \text{there exists a faithful } R\text{-}H\text{-BM-module} \}.$$

By Theorem 11.3.4(1), $\mathcal{K}$ is an H -special class and $r^{\mathcal{K}}(R) = \mathcal{M}(R)$. Using Proposition 11.3.9 and 11.3.10, we have that

$$\mathcal{K} = \{R \mid R \text{ is an } H\text{-simple module algebra with unit} \}.$$

Therefore $\mathcal{M}(R) = r_{Hbm}(R)$. $\square$

Assume that H is a finite-dimensional semisimple Hopf algebra with $t \in \int_H^l$ and $\epsilon(t) = 1$. Let

$$G_t(a) = \{z \mid z = x + (t \cdot a)x + \sum (x_i(t \cdot a)y_i + x_i y_i) \text{ for all } x_i, y_i, x \in R\}.$$

R is called an r_{gt} - H -module algebra, if $a \in G_t(a)$ for all $a \in R$.

Theorem 11.3.12. *r_{gr} is an H -radical property of H -module algebra and $r_{gt} = r_{Hbm}$.*

Proof. It is clear that any H -homomorphic image of r_{gt} - H -module algebra is an r_{gt} - H -module algebra. Let

$$N = \sum \{I \triangleleft_H \mid I \text{ is an } r_{gt}\text{-}H\text{-ideal of } R\}.$$

Now we show that N is an r_{gt} - H -ideal of R . In fact, we only need to show that $I_1 + I_2$ is an r_{gt} - H -ideal for any two r_{gt} - H -ideals I_1 and I_2 . For any $a \in I_1, b \in I_2$, there exist $x, x_i, y_i \in R$ such that

$$a = x + (t \cdot a)x + \sum_i (x_i(t \cdot a)y_i + x_i y_i).$$

Let

$$c = x + (t \cdot (a + b))x + \sum x_i(t \cdot (a + b))y_i + x_i y_i \in G_t(a + b).$$

Obviously,

$$a + b - c = b - (t \cdot b)x - \sum x_i(t \cdot b)y_i \in I_2.$$

Thus there exist $w, u_j, v_j \in R$ such that

$$a + b - c = w + (t \cdot (a + b - c))w + \sum_j (u_j(t \cdot (a + b - c))v_j + u_j v_j).$$

Let $d = (t \cdot (a + b))w + w + \sum_j (u_j(t \cdot (a + b))v_j + u_j v_j)$ and $e = c - \sum_j u_j(t \cdot c)v_j - (t \cdot c)w$. By computation, we have that

$$a + b = d + e.$$

Since $c \in G_t(a+b)$ and $d \in G_t(a+b)$, we get that $e \in G_t(a+b)$ and $a+b \in G_t(a+b)$, which implies that $I_1 + I_2$ is an r_{gt} - H -ideal.

Let $\bar{R} = R/N$ and $\bar{B}$ be an r_{gt} - H -ideal of $\bar{R}$. For any $a \in B$, there exist $x, x_i, y_i \in R$ such that

$$\bar{a} = \bar{x} + (t \cdot \bar{a})\bar{x} + \sum (\bar{x}_i(t \cdot \bar{a})\bar{y}_i + \bar{x}_i\bar{y}_i)$$

and

$$x + (t \cdot a)x + \sum (x_i(t \cdot a)y_i + x_iy_i) - a \in N.$$

Let

$$c = x + (t \cdot a)x + \sum (x_i(t \cdot a)y_i + x_iy_i) \in G_t(a).$$

Thus there exist $w, u_j, v_j \in R$ such that

$$a - c = (t \cdot (a - c))w + w + \sum (u_j(t \cdot (a - c))v_j + u_jv_j)$$

and

$$a = (t \cdot a)w + w + \sum u_j(t \cdot a)v_j + u_jv_j + c - (t \cdot c)w - \sum u_j(t \cdot c)v_j \in G_t(a),$$

which implies that B is an r_{gt} - H -ideal and $\bar{B} = 0$. Therefore r_{gt} is an H -radical property. $\square$

Proposition 11.3.13. *If R is an H -simple module algebra, then $r_{gr}(R) = 0$ iff R has a unit.*

Proof. If R is an H -simple module algebra with unit 1, then $-1 \notin G_t(-1)$ since

$$x + (t \cdot (-1))x + \sum (x_i(t \cdot (-1))y_i + x_iy_i) = 0$$

for any $x, x_i, y_i \in R$. Thus R is r_{gt} - H -semisimple. Conversely, if $r_{gt}(R) = 0$, then there exists $0 \neq a \notin G_t(a)$ and $G_t(a) = 0$, which implies that $ax + x = 0$ for any $x \in R$. It follows from Lemma 9.3.3 (7) that R has a unit. $\square$

Theorem 11.3.14. $r_{gt} = r_{Hbm}$.

Proof. By Proposition 11.3.13, $r_{gt}(R) \subseteq r_{Hbm}(R)$ for any H -module algebra R . It remains to show that if $a \notin r_{gt}(R)$ then $a \notin r_{Hbm}(R)$. Obviously, there exists $b \in (a)$ such that $b \notin G_t(b)$, where (a) denotes the H -ideal generated by a in R . Let

$$\mathcal{E} = \{I \triangleleft_H R \mid G_t(b) \subseteq I, b \notin I\}.$$

By Zorn's Lemma, there exists a maximal element P in $\mathcal{E}$. P is a maximal H -ideal of R , for, if Q is an H -ideal of R with $P \subseteq Q$ and $P \neq Q$, then $b \in Q$ and $x = -bx + (bx + x) \in Q$ for any $x \in R$. Consequently, R/P is an H -simple module algebra with $r_{gt}(R/P) = 0$. It follows from Proposition 11.3.13 that R/P is an H -simple module algebra with unit and $r_{Hbm}(R) \subseteq P$. Therefore $b \notin r_{Hbm}(R)$ and so $a \notin r_{Hbm}(R)$. $\square$

Definition 11.3.15. Let I be an H -ideal of H -module algebra R , N be an R - H -submodule of R - H -module M . (N, I) are said to have "L-condition", if for any finite subset $F \subseteq I$, there exists a positive integer k such that $F^k N = 0$.

Definition 11.3.16. An R - H -module M is called an R - H - L -module, if for M the following conditions are fulfilled:

(i) $RM \neq 0$.

(ii) For every non-zero R - H -submodule N of M and every H -ideal I of R , if (N, I) has “ L -condition”, then $I \subseteq (0 : M)_R$.

Proposition 11.3.17. If M is an R - H - L -module, then $R/(0 : M)_R$ is an r_{lH} - H -semisimple and H -prime module algebra.

Proof. If M is an R - H - L -module, let $\bar{R} = R/(0 : M)_R$. Obviously, $\bar{R}$ is H -prime. If $\bar{B}$ is an r_{lH} - H -ideal of $\bar{R}$, then (M, B) has “ L -condition” in R - H -module M , since for any finite subset F of B , there exists a natural number n such that $F^n \subseteq (0 : M)_R$ and $F^n M = 0$. Consequently, $B \subseteq (0 : M)_R$ and $\bar{R}$ is r_{lH} -semisimple. $\square$

Proposition 11.3.18. R is a non-zero r_{lH} - H -semisimple and H -prime module algebra iff there exists a faithful R - H - L -module.

Proof. If R is a non-zero r_{lH} - H -semisimple and H -prime module algebra, let $M = R$. Since R is an H -prime module algebra, $(0 : M)_R = 0$. If (N, B) has “ L -condition” for non-zero R - H -submodule of M and H -ideal B , then, for any finite subset F of B , there exists a natural number n , such that $F^n N = 0$ and $F^n(NR) = 0$, which implies that $F^n = 0$ and B is an r_{lH} - H -ideal, i.e. $B = 0 \subseteq (0 : M)_R$. Consequently, M is a faithful R - H - L -module.

Conversely, if M is a faithful R - H - L -module, then R is an H -prime module algebra. If I is an r_{lH} - H -ideal of R , then (M, I) has “ L -condition”, which implies $I = 0$ and R is an r_{lH} - H -semisimple module algebra. $\square$

Theorem 11.3.19. Let $\mathcal{M}_R = \{ M \mid M \text{ is an } R\text{-}H\text{-}L\text{-module} \}$ for any H -module algebra R and $\mathcal{M} = \cup \mathcal{M}_R$. Then $\mathcal{M}$ is an H -special class of modules and $\mathcal{M}(R) = r_{Hl}(R)$, where $\mathcal{K} = \{ R \mid R \text{ is an } H\text{-prime module algebra with } r_{lH}(R) = 0 \}$ and $r_{Hl} = r^{\mathcal{K}}$.

Proof. Obviously, (M1) holds. To show that (M2) holds, we only need to show that if I is an H -ideal of R and $M \in \mathcal{M}_I$, then $IM \in \mathcal{M}_R$. By Lemma 11.3.3(5), IM is an R - H -prime module. If (N, B) has the “ L -condition” for non-zero R - H -submodule N of IM and H -ideal B of R , i.e. for any finite subset F of B , there exists a natural number n such that $F^n N = 0$, then (N, BI) has “ L -condition” in I - H -module M . Thus $BI \subseteq (0 : M)_I = (0 : IM)_R \cap I$. Considering $(0 : IM)_R$ is an H -prime ideal of R , we have that $B \subseteq (0 : IM)_R$ or $I \subseteq (0 : IM)_R$. If $I \subseteq (0 : IM)_R$, then $I^2 \subseteq (0 : M)_I$ and $I \subseteq (0 : M)_I$, which contradicts $IM \neq 0$. Therefore $B \subseteq (0 : IM)_R$ and so IM is an R - H - L -module.

To show that (M3) holds, we only need to show that if $M \in \mathcal{M}_R$ and $I \triangleleft_H R$ with $IM \neq 0$, then $M \in \mathcal{M}_I$. By Lemma 11.3.3(6), M is an I - H -prime module. If (N, B) has the “ L -condition” for non-zero I - H -submodule N of M and H -ideal B of I , then IN is an R - H -prime module and $(IN, (B))$ has “ L -condition” in R - H -module M , since for any finite subset F of (B) , $F^3 \subseteq B$ and there exists a natural number n such that $F^{3n} IN \subseteq F^{3n} N = 0$, where (B) is the H -ideal generated by B in R . Therefore, $(B) \subseteq (0 : M)_R$ and $B \subseteq (0 : M)_I$, which implies $M \in \mathcal{M}_I$.

Finally, we show that (M4) holds. Let $I \triangleleft_H R$ and $\bar{R} = R/I$. If $M \in \mathcal{M}_R$ and $I \subseteq (0 : M)_R$, then M is an $\bar{R}$ - H -prime module. If $(N, \bar{B})$ has "L-condition" for H -ideal $\bar{B}$ of $\bar{R}$ and $\bar{R}$ - H -submodule N of M , then subset $F \subseteq B$ and there exists a natural number n such that $F^n N = (\bar{F})^n N = 0$. Consequently, $M \in \mathcal{M}_{\bar{R}}$. Conversely, if $M \in \mathcal{M}_{\bar{R}}$, we can similarly show that $M \in \mathcal{M}_R$.

The second claim follows from Proposition 11.3.18 and Theorem 11.3.4(1). $\square$

Theorem 11.3.20. $r_{Hl} = r_{lH}$.

Proof. Obviously, $r_{lH} \leq r_{Hl}$. It remains to show that $r_{Hl}(R) \neq R$ if $r_{lH}(R) \neq R$. There exists a finite subset F of R such that $F^n \neq 0$ for any natural number n . Let

$$\mathcal{F} = \{I \mid I \text{ is an } H\text{-ideal of } R \text{ with } F^n \not\subseteq I \text{ for any natural number } n\}.$$

By Zorn's lemma, there exists a maximal element P in $\mathcal{F}$. It is clear that P is an H -prime ideal of R . Now we show that $r_{lH}(R/P) = 0$. If $0 \neq B/P$ is an H -ideal of R/P , then there exists a natural number m such that $F^m \subseteq B$. Since $(F^m + P)^n \neq 0 + P$ for any natural number n , we have that B/P is not locally nilpotent and $r_{lH}(R/P) = 0$. Consequently, $r_{Hl}(R) \neq R$. $\square$

In fact, all of the results hold in braided tensor categories determined by (co)quasitriangular structure.

Chapter 12

The H -Radicals of Twisted H -Module Algebras

Remark ¹

J.R. Fisher [48] built up the general theory of H -radicals for H -module algebras. He studied H -Jacobson radical and obtained

$$r_j(R\#H) \cap R = r_{Hj}(R) \quad (12.1)$$

for any irreducible Hopf algebra H ([48, Theorem 4]). J.R. Fisher [48] asked when is

$$r_j(R\#H) = r_{Hj}(R)\#H \quad (12.2)$$

and asked if

$$r_j(R\#H) \subseteq (r_j(R) : H)\#H \quad (12.3)$$

R.J. Blattner, M. Cohen and S. Montgomery in [9] asked whether $R\#_\sigma H$ is semiprime with a finite-dimensional semisimple Hopf algebra H when R is semiprime, which is called the semiprime problem.

If H is a finite-dimensional semisimple Hopf algebra and R is semiprime, then $R\#_\sigma H$ is semiprime in the following five cases:

- (i) k is a perfect field and H is cocommutative;
- (ii) H is irreducible cocommutative;
- (iii) The weak action of H on R is inner;
- (iv) $H = (kG)^*$, where G is a finite group;
- (v) H is cocommutative.

Part (i) (ii) are due to W. Chin [23, Theorem 2, Corollary 1]. Part (iii) is due to B.J. Blattner and S. Montgomery [19, Theorem 2.7]. Part (iv) is due to M. Cohen and S. Montgomery [29, Theorem 2.9]. Part (v) is due to S. Montgomery and H.J. Schneider [90, Corollary 7.13].

If $H = (kG)^*$, then relation (12.2) holds, due to M. Cohen and S. Montgomery [29, Theorem 4.1]

¹This chapter can be omitted

In this chapter we obtain the relation between H -radical of H -module algebra R and radical of $R\#H$. We give some sufficient conditions for (12.2) and (12.3) and the formulae, which are similar to (12.1), (12.2) and (12.3) for H -prime radical respectively. We show that (12.1) holds for any Hopf algebra H . Using radical theory and the conclusions in [90], we also obtain that if H is a finite-dimensional semisimple, cosemisimple and either commutative or cocommutative Hopf algebra, then R is H -semiprime iff R is semiprime iff $R\#_\sigma H$ is semiprime.

In this chapter, unless otherwise stated, let k be a field, R be an algebra with unit over k , H be a Hopf algebra over k and H^* denote the dual space of H .

R is called a twisted H -module algebra if the following conditions are satisfied:

- (i) H weakly acts on R ;
- (ii) R is a twisted H -module, that is, there exists a linear map $\sigma \in \text{Hom}_k(H \otimes H, R)$ such that $h \cdot (k \cdot r) = \sum \sigma(h_1, k_1)(h_2 k_2 \cdot r) \sigma^{-1}(h_3, k_3)$ for all $h, k \in H$ and $r \in R$.

It is clear that if σ is trivial, then twisted H -module algebra R is an H -module algebra.

Set

$$\text{Spec}(R) = \{I \mid I \text{ is a prime ideal of } R\};$$

$$H\text{-Spec}(R) = \{I \mid I \text{ is an } H\text{-prime ideal of } R\}.$$

12.1 The Baer radical of twisted H -module algebras

In this section, let k be a commutative associative ring with unit, H be an algebra with unit and comultiplication Δ , R be an algebra over k (R may be without unit) and R be a twisted H -module algebra.

Definition 12.1.1. $r_{Hb}(R) := \cap \{I \mid I \text{ is an } H\text{-semiprime ideal of } R\};$

$$r_{bH}(R) := (r_b(R) : H)$$

$r_{Hb}(R)$ is called the H -Baer radical (or H -prime radical) of twisted H -module algebra R .

Lemma 12.1.2. (1) If E is a non-empty subset of R , then $(E) = (H \cdot E) + R(H \cdot E) + (H \cdot E)R + R(H \cdot E)R$, where (E) denotes the H -ideal generated by E in R ;

(2) If I is a nilpotent H -ideal of R , then $I \subseteq r_{Hb}(R)$.

Proof. (1) It is trivial.

(2) If I is a nilpotent H -ideal and P is an H -semiprime ideal, then $(I + P)/P$ is nilpotent simply because $(I + P)/P \cong I/(I \cap P)$ (as algebras). Thus $I \subseteq P$ and $I \subseteq r_{Hb}(R)$. $\square$

Proposition 12.1.3. (1) $r_{Hb}(R) = 0$ iff R is H -semiprime;

(2) $r_{Hb}(R/r_{Hb}(R)) = 0$;

(3) R is H -semiprime iff $(H \cdot a)R(H \cdot a) = 0$ always implies $a = 0$ for any $a \in R$;

R is H -prime iff $(H \cdot a)R(H \cdot b) = 0$ always implies $a = 0$ or $b = 0$ for any $a, b \in R$;

(4) If R is H -semiprime, then $W_H(R) = 0$.

Proof. (1) If $r_{Hb}(R) = 0$, then R is H -semiprime by Lemma 12.1.2 (2). Conversely, if R is H -semiprime, then 0 is an H -semiprime ideal and so $r_{Hb}(R) = 0$ by Definition 12.1.1.

(2) If $B/r_{Hb}(R)$ is a nilpotent H -ideal of $R/r_{Hb}(R)$, then $B^k \subseteq r_{Hb}(R)$ for some natural number k and so $B \subseteq r_{Hb}(R)$, which implies that $R/r_{Hb}(R)$ is H -semiprime. Thus $r_{Hb}(R/r_{Hb}(R)) = 0$ by part (1).

(3) If R is H -prime and $(H \cdot a)R(H \cdot b) = 0$ for a and $b \in R$, then $(a)^2(b)^2 = 0$ by Lemma 12.1.2 (1), where (a) and (b) are the H -ideals generated by a and b in R respectively. Since R is H -prime, $a = 0$ or $b = 0$. Conversely, if both B and C are H -ideals of R and $BC = 0$, then $(H \cdot a)R(H \cdot b) = 0$ and $a = 0$ or $b = 0$ for any $a \in B$ and $b \in C$, which implies that $B = 0$ or $C = 0$. Thus R is an H -prime. Similarly, the other assertion holds.

(4) For any $0 \neq a \in R$, there exist $b_1 \in R$ and $h_1, h'_1 \in H$ such that $0 \neq a_2 = (h_1 \cdot a_1)b_1(h'_1 \cdot a_1) \in (H \cdot a_1)R(H \cdot a_1)$ by part (3), where $a_1 = a$. Similarly, for $0 \neq a_2 \in R$, there exist $b_2 \in A$, $h_2, h'_2 \in H$ such that $0 \neq a_3 = (h_2 \cdot a_2)b_2(h'_2 \cdot a_2) \in (H \cdot a_2)R(H \cdot a_2)$, which implies that there exists an H - m -sequence $\{a_n\}$ such that $a_n \neq 0$ for any natural number n . Thus $W_H(R) = 0$. $\square$

Theorem 12.1.4. $r_{Hb}(R) = W_H(R) = \cap \{I \mid I \text{ is an } H\text{-prime ideal of } R\}$.

Proof. Let $D = \cap \{I \mid I \text{ is an } H\text{-prime ideal of } R\}$. Obviously, $r_{Hb}(R) \subseteq D$.

If $0 \neq a \notin W_H(R)$, then there exists an m -sequence $\{a_i\}$ in R with $a_1 = a$ and $a_{n+1} = (h_n \cdot a_n)b_n(h'_n \cdot a_n) \neq 0$ for $n = 1, 2, \dots$. Let $\mathcal{F} = \{I \mid I \text{ is an } H\text{-ideal of } R \text{ and } I \cap \{a_1, a_2, \dots\} = \emptyset\}$. By Zorn's Lemma, there exists a maximal P in $\mathcal{F}$. If both I and J are H -ideals of R with $I \not\subseteq P$ and $J \not\subseteq P$ such that $IJ \subseteq P$, then there exist natural numbers n and m such that $a_n \in I + P$ and $a_m \in J + P$. Since $a_{n+m+1} = (h_{n+m} \cdot a_{n+m})b_{n+m}(h'_{n+m} \cdot a_{n+m}) \in (I + P)(J + P) \subseteq P$, we get a contradiction. Thus P is an H -prime ideal of R . Obviously, $a \notin P$, which implies that $a \notin D$. Therefore $D \subseteq W_H(R)$.

For any $x \in W_H(R)$, let $\bar{R} = R/r_{Hb}(R)$. It follows from Proposition 12.1.3 (1) (2) (4) that $W_H(\bar{R}) = 0$. For any H - m -sequence $\{\bar{a}_n\}$ with $\bar{a}_1 = \bar{x}$ in $\bar{R}$, there exist $\bar{b}_n \in \bar{R}$ and $h_n, h'_n \in H$ such that $\bar{a}_{n+1} = (h_n \cdot \bar{a}_n)\bar{b}_n(h'_n \cdot \bar{a}_n)$ for any natural number n . Let $a'_1 = x$ and $a'_{n+1} = (h_n \cdot a'_n)b_n(h'_n \cdot a'_n)$ for any natural number n . Since $\{a'_n\}$ is an H - m -sequence with $a'_1 = x$ in R , there exists a natural number k such that $a'_k = 0$. It is clear that $\bar{a}_n = \overline{a'_n}$ for any natural number n by induction. Thus $\bar{a}_k = 0$ and $\bar{x} \in W_H(\bar{R})$. Considering $W_H(\bar{R}) = 0$, we have $x \in r_{Hb}(R)$, which implies that $W_H(R) \subseteq r_{Hb}(R)$. Therefore $W_H(R) = r_{Hb}(R) = D$. $\square$

12.2 The Baer and Jacobson radicals of crossed products

By [87, Lemma 7.1.2], if $R \#_\sigma H$ is crossed product defined in [87, Definition 7.1.1], then R is a twisted H -module algebra.

Let R be an algebra and $M_{m \times n}(R)$ be the algebra of $m \times n$ matrices with entries in R . For $i = 1, 2, \dots, m$ and $j = 1, 2, \dots, n$. Let $(e_{ij})_{m \times n}$ denote the matrix in $M_{m \times n}(R)$, where (i, j) -entry is 1_R and the others are zero. Set

$$\mathcal{I}(R) = \{I \mid I \text{ is an ideal of } R\}.$$

$$r_{Hj}(R) := r_j(R \#_\sigma H) \cap R;$$

$$r_{jH}(R) := (r_j(R) : H).$$

Lemma 12.2.1. *Let M_R be a free R -module with finite rank and $R' = \text{End}(M_R)$. Then there exists a unique bijective map*

$$\Phi : \mathcal{I}(R) \longrightarrow \mathcal{I}(R')$$

such that $\Phi(I)M = MI$ and

- (1) Φ is a map preserving containments, finite products, and infinite intersections;*
- (2) $\Phi(I) \cong M_{n \times n}(I)$ for any ideal I of R ;*
- (3) I is a (semi)prime ideal of R iff $\Phi(I)$ is a (semi)prime ideal of R' ;*
- (4) $r_b(R') = \Phi(r_b(R))$;*
- (5) $r_j(R') = \Phi(r_j(R))$.*

Proof Since M_R is a free R -module with rank n , we can assume $M = M_{n \times 1}(R)$. Thus $R' = \text{End}(M_R) = M_{n \times n}(R)$ and the module operation of M over R becomes the matrix operation. Set $M' = M_{1 \times n}(R)$. Obviously, $M'M = R$. Since $(e_{i1})_{n \times 1}(e_{1j})_{1 \times n} = (e_{ij})_{n \times n}$ for $i, j = 1, 2, \dots, n$, $MM' = M_{n \times n}(R) = R'$. Define

$$\Phi(I) = MIM'$$

for any ideal I of R . By simple computation, we have that $\Phi(I)$ is an ideal of R' and $\Phi(I)M = MI$. If J is an ideal of R' such that $JM = MI$, then $JM = \Phi(I)M$ and $J = \Phi(I)$, which implies Φ is unique. In order to show that Φ is a bijection from $\mathcal{I}(R)$ onto $\mathcal{I}(R')$, we define a map Ψ from $\mathcal{I}(R')$ to $\mathcal{I}(R)$ sending I' to $M'I'M$ for any ideal I' of R' . Since $\Phi\Psi(I') = MM'I'MM' = I'$ and $\Psi\Phi(I) = M'MIM'M = I$ for any ideal I' of R' and ideal I of R , we have that Φ is bijective.

(1) Obviously Φ preserves containments. We see that

$$\Phi(IJ) = MIJM' = (MIM')(MJM') = \Phi(I)\Phi(J)$$

for any ideals I and J of R . Thus Φ preserves finite products. To show that Φ preserves infinite intersections, we first show that

$$M(\cap\{I_\alpha \mid \alpha \in \Omega\}) = \cap\{MI_\alpha \mid \alpha \in \Omega\} \tag{12.4}$$

for any $\{I_\alpha \mid \alpha \in \Omega\} \subseteq \mathcal{I}(R)$. Obviously, the right side of relation (12.4) contains the left side of relation (12.4). Let $\{u_1, u_2, \dots, u_n\}$ be a basis of M over R . For any $x \in \cap\{MI_\alpha \mid \alpha \in \Omega\}$, any $\alpha, \alpha' \in \Omega$, there exist $r_i \in I_\alpha$ and $r'_i \in I_{\alpha'}$ such that $x = \sum u_i r_i = \sum u_i r'_i$. Since $\{u_i\}$ is a basis, $r_i = r'_i$, which implies $x \in M(\cap\{I_\alpha \mid \alpha \in \Omega\})$. Thus the relation (12.4) holds. It follows from relation (12.4) that

$$\Phi(\cap\{I_\alpha \mid \alpha \in \Omega\})M = \cap\{\Phi(I_\alpha)M \mid \alpha \in \Omega\}$$

Since $\Phi(\cap\{I_\alpha \mid \alpha \in \Omega\})M = \cap\{\Phi(I_\alpha)M \mid \alpha \in \Omega\} \supseteq (\cap\{\Phi(I_\alpha) \mid \alpha \in \Omega\})M$, we have that

$$\Phi(\cap\{I_\alpha \mid \alpha \in \Omega\}) \supseteq \cap\{\Phi(I_\alpha) \mid \alpha \in \Omega\}.$$

Obviously,

$$\Phi(\cap\{I_\alpha \mid \alpha \in \Omega\}) \subseteq \cap\{\Phi(I_\alpha) \mid \alpha \in \Omega\}.$$

Thus

$$\Phi(\cap\{I_\alpha \mid \alpha \in \Omega\}) = \cap\{\Phi(I_\alpha) \mid \alpha \in \Omega\}.$$

(2) Obviously, $\Phi(I) = MIM' = M_{n \times 1}(R)IM_{1 \times n}(R) \subseteq M_{n \times n}(I)$. Since $a(e_{ij})_{n \times n} = (e_{i1})_{n \times 1}a(e_{1j})_{1 \times n} \in MIM'$ for all $a \in I$ and $i, j = 1, 2, \dots, n$,

$$\Phi(I) = MIM' = M_{n \times 1}(R)IM_{1 \times n}(R) \supseteq M_{n \times n}(I).$$

Thus part (2) holds.

(3) Since bijection Φ preserves products, part (3) holds.

(4) We see that

$$\begin{aligned} \Phi(r_b(R)) &= \Phi(\cap\{I \mid I \text{ is a prime ideal of } R\}) \\ &= \cap\{\Phi(I) \mid I \text{ is a prime ideal of } R\} \text{ by part (1)} \\ &= \cap\{\Phi(I) \mid \Phi(I) \text{ is a prime ideal of } R'\} \text{ by part (3)} \\ &= \cap\{I' \mid I' \text{ is a prime ideal of } R'\} \text{ since } \Phi \text{ is surjective} \\ &= r_b(R') \end{aligned}$$

(5) We see that

$$\begin{aligned} \Phi(r_j(R)) &= M_{n \times n}(r_j(R)) \text{ by part (2)} \\ &= r_j(M_{n \times n}(R)) \text{ by [116, Theorem 30.1]} \\ &= r_j(R') . \quad \square \end{aligned}$$

Let H be a finite-dimensional Hopf algebra and $A = R \#_\sigma H$. Then A is a free right R -module with finite rank by [87, Proposition 7.2.11] and $End(A_R) \cong (R \#_\sigma H) \# H^*$ by [87, Corollary 9.4.17]. By part (a) in the proof of [90, Theorem 7.2], it follows that Φ in Lemma 1.2 is the same as in [90, Theorem 7.2].

Lemma 12.2.2. *Let H be a finite-dimensional Hopf algebra and $A = R \#_\sigma H$. Then*

(1) *If P is an H^* -ideal of A , then $P = (P \cap R) \#_\sigma H$.*

(2) *$\Phi(I) = (I \#_\sigma H) \# H^*$ for every H -ideal I of R ;*

(3)

$$\{P \mid P \text{ is an } H\text{-ideal of } A \# H^*\} = \{(I \#_\sigma H) \# H^* \mid I \text{ is an } H\text{-ideal of } R\} \quad (12.5)$$

$$\{P \mid P \text{ is an } H^*\text{-ideal of } A\} = \{I \#_\sigma H \mid I \text{ is an } H\text{-ideal of } R\} \quad (12.6)$$

$$\{P \mid P \text{ is an } H\text{-prime ideal of } A \# H^*\}$$

$$= \{(I \#_\sigma H) \# H^* \mid I \text{ is an } H\text{-prime ideal of } R\} \quad (12.7)$$

(4) *$H\text{-Spec}(R) = \{(I : H) \mid I \in \text{Spec}(R)\}$;*

(5)

$$(\cap\{I_\alpha \mid \alpha \in \Omega\} : H) = \cap\{(I_\alpha : H) \mid \alpha \in \Omega\} \quad (12.8)$$

where I_α is an ideal of R for all $\alpha \in \Omega$;

(6)

$$(\cap\{I_\alpha \mid \alpha \in \Omega\})\#_\alpha H = \cap\{(I_\alpha\#_\sigma H) \mid \alpha \in \Omega\} \quad (12.9)$$

where I_α is an H -ideal of R for all $\alpha \in \Omega$;

$$(7) \quad \Phi(r_b(R)) = r_b(A\#H^*);$$

$$(8) \quad \Phi(r_j(R)) = r_j(A\#H^*);$$

$$(9) \quad \Phi(r_{Hb}(R)) = r_{Hb}(A\#H^*) = (r_{Hb}(R)\#_\sigma H)\#H^*.$$

Proof (1) By [87, Corollary 8.3.11], we have that $P = (P \cap R)A = (P \cap R)\#_\sigma H$.

(2) By the part (b) in the proof of [90, Theorem 7.2], it follows that

$$\Phi(I) = (I\#_\sigma H)\#H^*$$

for every H -ideal I of R .

(3) Obviously, the left side of relation (12.5) contains the right side of relation (12.5). If P is an H -ideal of $A\#H^*$, then $P = (P \cap A)\#H^* = (((P \cap A) \cap R)\#_\sigma H)\#H^*$ by part (1), which implies that the right side of relation (12.5) contains the left side. Thus relation (12.5) holds. Similarly, relation (12.6) holds. Now, we show that relation (12.7) holds. If P is an H -prime ideal of $A\#H^*$, there exists an H -ideal I of R such that $P = (I\#_\sigma H)\#H^*$ by relation (12.5). For any H -ideals J and J' of R with $JJ' \subseteq I$, since $\Phi(JJ') = \Phi(J)\Phi(J') \subseteq \Phi(I) = P$ by Lemma 12.2.1 (1), we have that $\Phi(J) \subseteq \Phi(I)$ or $\Phi(J') \subseteq \Phi(I)$, which implies that $J \subseteq I$ or $J' \subseteq I$ by Lemma 12.2.1. Thus I is an H -prime ideal of R . Conversely, if I is an H -prime ideal of R and $P = (I\#_\sigma H)\#H^*$, we claim that P is an H -prime of $A\#H^*$. For any H -ideals Q and Q' of $A\#H^*$ with $QQ' \subseteq P$, there exist two H -ideals J and J' of R such that $(J\#_\sigma H)\#H^* = Q$ and $(J'\#_\sigma H)\#H^* = Q'$ by relation (12.5). Since $\Phi(JJ') = \Phi(J)\Phi(J') = QQ' \subseteq P = \Phi(I)$, $JJ' \subseteq I$, which implies $J \subseteq I$, or $J' \subseteq I$, and so $Q \subseteq P$ or $Q' \subseteq P$. Thus P is an H -prime ideal of $A\#H^*$. Consequently, relation (12.7) holds.

(4) It follows from [90, Lemma 7.3 (1) (2)].

(5) Obviously, the right side of relation (12.8) contains the left side. Conversely, if $x \in \cap\{(I_\alpha : H) \mid \alpha \in \Omega\}$, then $x \in (I_\sigma : H)$ and $h \cdot x \in I_\alpha$ for all $\alpha \in \Omega, h \in H$, which implies that $h \cdot x \in \cap\{I_\alpha \mid \alpha \in \Omega\}$ and $x \in (\cap\{I_\alpha \mid \alpha \in \Omega\} : H)$. Thus relation (12.8) holds.

(6) Let $\{h^{(1)}, \dots, h^{(n)}\}$ be a basis of H . Obviously, the right side of relation (12.9) contains the left side of relation (12.9). Conversely, for $u \in \cap\{(I_\alpha\#_\sigma H) \mid \alpha \in \Omega\}$ and $\alpha, \alpha' \in \Omega$, there exist $r_i \in I_\alpha$ and $r'_i \in I_{\alpha'}$ such that $u = \sum_{i=1}^n r_i\#h^{(i)} = \sum_{i=1}^n r'_i\#h^{(i)}$. Since $\{h^{(1)}, \dots, h^{(n)}\}$ is linearly independent, we have that $r_i = r'_i$, which implies that $u \in (\cap\{I_\alpha \mid \alpha \in \Omega\})\#_\alpha H$. Thus relation (12.9) holds.

(7) and (8) follow from Lemma 12.2.1(4)(5).

(9) We see that

$$\begin{aligned}
r_{Hb}(A\#H^*) &= \cap\{P \mid P \text{ is an } H\text{-prime ideal of } A\#H^*\} \quad \text{by Theorem 12.1.4} \\
&= \cap\{(I\#_\sigma H)\#H^* \mid I \text{ is an } H\text{-prime ideal of } R\} \quad \text{by relation (12.7)} \\
&= (\cap\{I\#_\sigma H \mid I \text{ is an } H\text{-prime ideal of } R\})\#H^* \quad \text{by part (6)} \\
&= ((\cap\{I \mid I \text{ is an } H\text{-prime ideal of } R\})\#_\sigma H)\#H^* \quad \text{by part (6)} \\
&= (r_{Hb}(R)\#_\sigma H)\#H^* \quad \text{by Theorem 12.1.4} \\
&= \Phi(r_{Hb}(R)) \quad \text{by part (2)}.
\end{aligned}$$

(10) If H is cosemisimple, then H is semisimple by [87, Theorem 2.5.2]. Conversely, if H is semisimple, then H^* is cosemisimple. By [87, Theorem 2.5.2], H^* is semisimple. Thus H is cosemisimple. $\square$

Proposition 12.2.3. (1) $r_{Hb}(R) \subseteq r_b(R\#_\sigma H) \cap R \subseteq r_{bH}(R)$;
(2) $r_{Hb}(R)\#_\sigma H \subseteq r_b(R\#_\sigma H)$.

Proof. (1) If P is a prime ideal of $R\#_\sigma H$, then $P \cap R$ is an H -prime ideal of R by [24, Lemma 1.6]. Thus $r_b(R\#_\sigma H) \cap R = \cap\{P \cap R \mid P \text{ is prime ideal of } R\#_\sigma H\} \supseteq r_{Hb}(R)$. For any $a \in r_b(R\#_\sigma H) \cap R$ and any m -sequence $\{a_i\}$ in R with $a_1 = a$, it is easy to check that $\{a_i\}$ is also an m -sequence in $R\#_\sigma H$. Thus $a_n = 0$ for some natural n , which implies $a \in r_b(R)$. Thus $r_b(R\#_\sigma H) \cap R \subseteq r_{bH}(R)$ by [19, Lemma 1.6]

(2) We see that

$$\begin{aligned}
r_{Hb}(R)\#_\sigma H &= (r_{Hb}(R)\#_\sigma 1)(1\#_\sigma H) \\
&\subseteq r_b(R\#_\sigma H)(1\#_\sigma H) \quad \text{by part (1)} \\
&\subseteq r_b(R\#_\sigma H). \quad \square
\end{aligned}$$

Proposition 12.2.4. Let H be finite-dimensional Hopf algebra and $A = R\#_\sigma H$. Then

- (1) $r_{H^*b}(R\#_\sigma H) = r_{Hb}(R)\#_\sigma H$;
(2) $r_{Hb}(R) = r_{bH}(R) = r_b(R\#_\sigma H) \cap R$.

Proof

(1) We see that

$$\begin{aligned}
r_{H^*b}(R\#_\sigma H) &= \cap\{P \mid P \text{ is an } H^*\text{-prime ideal of } A\} \\
&= \cap\{I\#_\sigma H \mid I \text{ is an } H\text{-prime ideal of } R\} \quad (\text{by [90, Lemma 7.3 (4)]}) \\
&= (\cap\{I \mid I \text{ is an } H\text{-prime ideal of } R\})\#_\sigma H \quad (\text{by Lemma 12.2.2 (6)}) \\
&= r_{Hb}(R)\#_\sigma H.
\end{aligned}$$

(2) We see that

$$\begin{aligned}
r_{Hb}(R) &= \cap\{P \mid P \text{ is an } H\text{-prime ideal of } R\} \\
&= \cap\{(I : H) \mid I \in \text{Spec}(R)\} \text{ by Lemma 12.2.2 part (4)} \\
&= (\cap\{I \mid I \in \text{Spec}(R)\} : H) \text{ by Lemma 12.2.2 part (5)} \\
&= (r_b(R) : H) \\
&= r_{bH}(R).
\end{aligned}$$

Thus it follows from Proposition 12.2.3(1) that $r_{Hb}(R) = r_b(R\#_\sigma H) \cap R = r_{bH}(R)$. $\square$

Theorem 12.2.5. . *Let H be a finite-dimensional Hopf algebra and the weak action of H be inner. Then*

- (1) $r_{Hb}(R) = r_b(R) = r_{bH}(R)$;
- Moreover, if H is semisimple, then
- (2) $r_b(R\#_\sigma H) = r_{Hb}(R)\#_\sigma H$.

Proof (1) Since the weak action is inner, every ideal of R is an H -ideal, which implies that $r_{Hb}(R) = r_b(R) = r_{bH}(R)$ by Proposition 12.2.4 (2).

(2) Considering Proposition 12.2.3(2), it suffices to show $r_b(R\#_\sigma H) \subseteq r_{Hb}(R)\#_\sigma H$. It is clear that

$$(R\#_\sigma H)/(r_{Hb}(R)\#_\sigma H) \cong (R/r_{Hb}(R))\#_\sigma H \quad (\text{as algebras}). \quad (12.10)$$

It follows by [87, Theorem 7.4.7] that $(R/r_{Hb}(R))\#_\sigma H$ is semiprime. Therefore

$$r_b(R\#_\sigma H) \subseteq r_{Hb}(R)\#_\sigma H. \quad \square$$

Theorem 12.2.6. *Let H be a finite-dimensional, semisimple and either commutative or cocommutative Hopf algebra and let $A = R\#_\sigma H$. Then*

- (1) $r_b(R\#_\sigma H) = r_{Hb}(R)\#_\sigma H$;
- (2) R is H semiprime iff $R\#_\sigma H$ is semiprime.

Moreover, if H is cosemisimple, or $\text{char } k$ does not divide $\dim H$, then both part (3) and part (4) hold:

- (3) $r_{Hb}(R) = r_{bH}(R) = r_b(R)$;
- (4) R is H -semiprime iff R is semiprime iff $R\#_\sigma H$ is semiprime.

Proof (1) Considering Proposition 12.2.3(2), it suffices to show

$$r_b(R\#_\sigma H) \subseteq r_{Hb}(R)\#_\sigma H.$$

It follows by [90, Theorem 7.12 (3)] that $(R/r_{Hb}(R))\#_\sigma H$ is semiprime. Using relation (12.10), we have that $r_b(R\#_\sigma H) \subseteq r_{Hb}(R)\#_\sigma H$.

(2) It follows from part (1) and Proposition 12.1.3 (1).

(3) By [67, Theorem 4.3 (1)], we have that H is semisimple and cosemisimple.

We see that

$$\begin{aligned}
\Phi(r_b(R)) &= r_b(A \# H^*) \quad \text{by Lemma 12.2.2 (7)} \\
&= r_{H^*b}(A) \# H^* \quad \text{by part (1)} \\
&= (r_{Hb}(R) \#_\sigma H) \# H^* \quad \text{by Proposition 12.2.4 (1)} \\
&= \Phi(r_{Hb}(R)) \quad \text{by Lemma 12.2.2 (2)}.
\end{aligned}$$

Thus $r_b(R) = r_{Hb}(R)$.

(4) It immediately follows from part (2) and part (3). $\square$

We now provide an example to show that the Baer radical $r_b(R)$ of R is not H -stable when H is not cosemisimple.

Example: Let k be a field of characteristic $p > 0$ and $R = k[x]/(x^p)$. Then we can define a derivation d on R by sending x to $x + 1$. Then $d^2(x) = d(x + 1) = d(x)$ and then, by induction, $d^p(x) = d(x)$. It follows that $d^p = d$ on all of R . Thus $H = u(kd)$, the restricted enveloping algebra, is semisimple by [87, Theorem 2.3.3]. clearly H acts on R , but H does not stabilize the Baer radical of R which is the principal ideal generated by x . Note also that H is commutative and cocommutative.

Proposition 12.2.7. *If R is an H -module algebra, then*

$$r_{Hj}(R) = \cap \{(0 : M)_R \mid M \text{ is an irreducible } R\text{-}H\text{-module}\}.$$

That is, $r_{Hj}(R)$ is the H -Jacobson radical of the H -module algebra R defined in [48].

Proof. It is easy to show that M is an irreducible $R\text{-}H$ -module iff M is an irreducible $R \# H$ -module by [48, Lemma 1]. Thus

$$\begin{aligned}
r_{Hj}(R) &= r_j(R \# H) \cap R \quad \text{by definition 12.1.1} \\
&= (\cap \{(0 : M)_{R \# H} \mid M \text{ is an irreducible } R \# H\text{-module}\}) \cap R \\
&= \cap \{(0 : M)_R \mid M \text{ is an irreducible } R\text{-}H\text{-module}\}. \square
\end{aligned}$$

Proposition 12.2.8. (1) $r_j(R \#_\sigma H) \cap R = r_{Hj}(R) \subseteq r_{jH}(R)$;

(2) $r_{Hj}(R) \#_\sigma H \subseteq r_j(R \#_\sigma H)$.

Proof. (1) For any $a \in r_j(R \#_\sigma H) \cap R$, there exists $u = \sum_i a_i \# h_i \in R \#_\sigma H$ such that

$$a + u + au = 0.$$

Let $(id \otimes \epsilon)$ act on the above equation. We get that $a + \sum a_i \epsilon(h_i) + a(\sum a_i \epsilon(h_i)) = 0$, which implies that a is a right quasi-regular element in R . Thus $r_j(R \#_\sigma H) \cap R \subseteq r_{jH}(R)$.

(2) It is similar to the proof of Proposition 12.2.3 (2). $\square$

Proposition 12.2.9. *Let H be a finite-dimensional Hopf algebra and $A = R \#_\sigma H$. Then*

(1) $r_{jH}(R) \#_\sigma H = r_{H^*j}(R \#_\sigma H)$;

(2) $r_{Hj}(R) = r_{jH}(R)$;

(3) $r_{Hj}(A \# H^*) = (r_{Hj}(R) \#_\sigma H) \# H^*$.

Proof (1) We see that

$$\begin{aligned}
(r_{jH}(R) \#_{\sigma} H) \# H^* &= \Phi(r_{jH}(R)) \\
&= (\Phi(r_j(R)) \cap A) \# H^* \quad \text{by [90, Theorem 7.2]} \\
&= (r_j(A \# H^*) \cap A) \# H^* \quad \text{by Lemma 12.2.2 (8)} \\
&= r_{H^*j}(A) \# H^* \quad \text{by Definition 12.1.1 .}
\end{aligned}$$

Thus $r_{H^*j}(A) = r_{jH}(R) \#_{\sigma} H$.

(2) We see that

$$\begin{aligned}
r_{Hj}(R) &= r_j(A) \cap R \\
&\supseteq r_{H^*j}(A) \cap R \quad \text{by Proposition 12.2.8 (1)} \\
&= r_{jH}(R) \quad \text{by part (1) .}
\end{aligned}$$

It follows by Proposition 12.2.1 (1) that $r_{Hj}(R) = r_{jH}(R)$.

(3) It immediately follows from part (1) (2). $\square$

By Proposition 12.2.8 and 12.2.9, it is clear that if H is a finite-dimensional Hopf algebra, then relation (12.2) holds iff relation (12.3) holds.

Theorem 12.2.10. *Let H be a finite-dimensional Hopf algebra and the weak action of H be inner. Then*

$$(1) \quad r_{Hj}(R) = r_j(R) = r_{jH}(R).$$

Moreover, if H is semisimple, then

$$(2) \quad r_j(R \#_{\sigma} H) = r_{Hj}(R) \#_{\sigma} H.$$

Proof (1) Since the weak action is inner, every ideal of R is an H -ideal and $r_j(R) = r_{jH}(R)$.

It follows from Proposition 12.2.2(2) that $r_{Hj}(R) = r_{jH}(R) = r_j(R)$.

(2) Considering Proposition 12.2.8(2), it suffices to show

$$r_j(R \#_{\sigma} H) \subseteq r_{Hj}(R) \#_{\sigma} H.$$

It is clear that

$$(R \#_{\sigma} H) / (r_{Hj}(R) \#_{\sigma} H) \cong (R / r_{Hj}(R)) \#_{\sigma} H \quad (\text{as algebras}).$$

It follows by [87, Corollary 7.4.3] and part (1) that $(R / r_{Hj}(R)) \#_{\sigma} H$ is semiprimitive. Therefore $r_j(R \#_{\sigma} H) \subseteq r_{Hj}(R) \#_{\sigma} H$. $\square$

Theorem 12.2.11. *Let H be a finite-dimensional, semisimple Hopf algebra, let k be an algebraically closed field and let $A = R \#_{\sigma} H$. Assume H is cosemisimple or $\text{char } k$ does not divide $\dim H$.*

(1) *If H is cocommutative, then*

$$r_{Hj}(R) = r_{jH}(R) = r_j(R);$$

(2) *If H is commutative, then*

$$r_j(R \#_{\sigma} H) = r_{Hj}(R) \#_{\sigma} H.$$

Proof By [67, Theorem 4.3 (1)] , we have that H is semisimple and cosemisimple.

(1) If $g \in G(H)$, then the weak action of g on R is an algebraic homomorphism, which implies that $g \cdot r_j(R) \subseteq r_j(R)$. Let H_0 be the coradical of H , $H_1 = H_0 \wedge H_0$, $H_{i+1} = H_0 \wedge H_i$ for $i = 1, \dots, n$, where n is the dimension $\dim H$ of H . It is clear that $H_0 = kG$ with $G = G(H)$ by [114, Theorem 8.0.1 (c)] and $H = \cup H_i$. It is easy to show that if $k > i$, then

$$H_i \cdot (r_j(R))^k \subseteq r_j(R)$$

by induction for i . Thus

$$H \cdot (r_j(R))^{\dim H + 1} \subseteq r_j(R),$$

which implies that $(r_j(R))^{\dim H + 1} \subseteq r_{jH}(R)$.

We see that

$$\begin{aligned} r_j(R/r_{jH}(R)) &= r_j(R)/r_{jH}(R) \\ &= r_b(R/r_{jH}(R)) \text{ since } r_j(R)/r_{jH}(R) \text{ is nilpotent} \\ &= r_{bH}(R/r_{jH}(R)) \text{ by Theorem 12.2.6 (3)} \\ &\subseteq r_{jH}(R/r_{jH}(R)) \\ &= 0 . \end{aligned}$$

Thus $r_j(R) \subseteq r_{jH}(R)$, which implies that $r_j(R) = r_{jH}(R)$.

(2) It immediately follows from part (1) and Proposition 12.2.9(1) (2). $\square$

12.3 The general theory of H -radicals for twisted H -module algebras

In this section we give the general theory of H -radicals for twisted H -module algebras.

Definition 12.3.1. Let r be a property of H -ideals of twisted H -module algebras. An H -ideal I of twisted H -module algebra R is called an r - H -ideal of R if it is of the r -property. A twisted H -module algebra R is called an r -twisted H -module algebra if it is r - H -ideal of itself. A property r of H -ideals of twisted H -module algebras is called an H -radical property if the following conditions are satisfied:

(R1) Every twisted H -homomorphic image of r -twisted H -module algebra is an r twisted H -module algebra;

(R2) Every twisted H -module algebra R has the maximal r - H -ideal $r(R)$;

(R3) $R/r(R)$ has not any non-zero r - H -ideal.

We call $r(R)$ the H -radical of R .

Proposition 12.3.2. Let r be an ordinary hereditary radical property for rings. An H -ideal I of twisted H -module algebra R is called an r_H - H -ideal of R if I is an r -ideal of ring R . Then r_H is an H -radical property for twisted H -module algebras.

Proof. (R1). If (R, σ) is an r_H -twisted H -module algebra and $(R, \sigma) \stackrel{f}{\sim} (R', \sigma')$, then $r(R') = R'$ by ring theory. Consequently, R' is an r_H -twisted H -module algebra.

(R2). For any twisted H -module algebra R , $r(R)$ is the maximal r -ideal of R by ring theory. It is clear that $r(R)_H$ is the maximal r - H -ideal, which is an r_H - H -ideal of R . Consequently, $r_H(R) = r(R)_H$ is the maximal r_H - H -ideal of R .

(R3). If $I/r_H(R)$ is an r_H - H -ideal of $R/r_H(R)$, then I is an r -ideal of algebra R by ring theory. Consequently, $I \subseteq r(R)$ and $I \subseteq r_H(R)$. $\square$

Proposition 12.3.3. r_{Hb} is an H -radical property.

Proof. (R1). Let (R, σ) be an r_{Hb} -twisted H -module algebra and $(R, \sigma) \stackrel{f}{\sim} (R', \sigma')$. For any $x' \in R'$ and any H - m -sequence $\{a'_n\}$ in R' with $a'_1 = x'$, there exist $b'_n \in R'$ and $h_n, h'_n \in H$ such that $a'_{n+1} = (h_n \cdot a'_n)b'_n(h'_n \cdot a'_n)$ for any natural number n . Let $a_1, b_i \in R$ such that $f(a_1) = x'$ and $f(b_i) = b'_i$ for $i = 1, 2, \dots$. Set $a_{n+1} = (h_n \cdot a_n)b_n(h'_n \cdot a_n)$ for any natural number n . Since $\{a_n\}$ is an H - m -sequence in R , there exists a natural number k such that $a_k = 0$. It is clear that $f(a_n) = a'_n$ for any natural number n by induction. Thus $a'_k = 0$, which implies that x' is an H - m -nilpotent element. Consequently, R' is an r_{Hb} -twisted H -module algebra.

(R2). By [130, Theorem 1.5], $r_{Hb}(R) = W_H(R) = \{a \mid a \text{ is an } H\text{-}m\text{-nilpotent element in } R\}$. Thus $r_{Hb}(R)$ is the maximal r_{Hb} - H -ideal of R .

(R3). It immediately follows from [130, Proposition 1.4]. $\square$

12.4 The relations among radical of R , radical of $R \#_\sigma H$, and H -radical of R

In this section we give the relation among the Jacobson radical $r_j(R)$ of R , the Jacobson radical $r_j(R \#_\sigma H)$ of $R \#_\sigma H$, and H -Jacobson radical $r_{Hj}(R)$ of R .

In this section, let k be a field, R an algebra with unit, H a Hopf algebra over k and $R \#_\sigma H$ an algebra with unit. Let r be a hereditary radical property for rings which satisfies

$$r(M_{n \times n}(R)) = M_{n \times n}(r(R))$$

for any twisted H -module algebra R .

Example. r_j , r_{bm} and r_n satisfy the above conditions by [116]. Using [130, Lemma 2.1 (2)], we can easily prove that r_b and r_l also satisfy the above conditions.

Definition 12.4.1. $\bar{r}_H(R) := r(R \#_\sigma H) \cap R$ and $r_H(R) := (r(R) : H)$.

If H is a finite-dimensional Hopf algebra and $M = R \#_\sigma H$, then M is a free right R -module with finite rank by [87, Proposition 7.2.11] and $End(M_R) \cong (R \#_\sigma H) \# H^*$ by [87, Corollary 9.4.17]. It follows from part (a) in the proof of [90, Theorem 7.2] that there exists a unique bijective map

$$\Phi : \mathcal{I}(R) \longrightarrow \mathcal{I}(R')$$

such that $\Phi(I)M = MI$, where $R' = (R \#_\sigma H) \# H^*$ and

$$\mathcal{I}(R) = \{I \mid I \text{ is an ideal of } R\}.$$

Lemma 12.4.2. *If H is a finite-dimensional Hopf algebra, then $\Phi(r(R)) = r((R\#_\sigma H)\#H^*)$.*

Proof. It is similar to the proof of [130, lemma 2.1 (5)]. $\square$

Proposition 12.4.3. $\bar{r}_H(R)\#_\sigma H \subseteq r_{H^*}(R\#_\sigma H) \subseteq r(R\#_\sigma H)$.

Proof. We see that

$$\begin{aligned} \bar{r}_H(R)\#_\sigma H &= (\bar{r}_H(R)\#_\sigma 1)(1\#_\sigma H) \\ &\subseteq r(R\#_\sigma H)(1\#_\sigma H) \\ &\subseteq r(R\#_\sigma H). \end{aligned}$$

Thus $\bar{r}_H(R)\#_\sigma H \subseteq r_{H^*}(R\#_\sigma H)$ since $\bar{r}_H(R)\#_\sigma H$ is an H^* -ideal of $R\#_\sigma H$. $\square$

Proposition 12.4.4. *If H is a finite-dimensional Hopf algebra, then*

(1) $r_H(R)\#_\sigma H = \bar{r}_{H^*}(R\#_\sigma H)$;

Furthermore, if $\bar{r}_H \leq r_H$, then

(2) $\bar{r}_H = r_H$ and $r_H(R)\#_\sigma H \subseteq r(R\#_\sigma H)$;

(3) $R\#_\sigma H$ is r -semisimple for any r_H -semisimple R iff

$$r(R\#_\sigma H) = r_H(R)\#_\sigma H.$$

Proof. Let $A = R\#_\sigma H$.

(1) We see that

$$\begin{aligned} (r_H(R)\#_\sigma H)\#H^* &= \Phi(r_H(R)) \\ &= (\Phi(r(R)) \cap A)\#H^* \quad \text{by [90, Theorem 7.2]} \\ &= (r(A\#H^*) \cap A)\#H^* \quad \text{by Lemma 12.4.2} \\ &= \bar{r}_{H^*}(A)\#H^* \quad \text{by Definition 12.4.1 .} \end{aligned}$$

Thus $\bar{r}_{H^*}(A) = r_H(R)\#_\sigma H$.

(2) We see that

$$\begin{aligned} \bar{r}_H(R) &= r(A) \cap R \\ &\supseteq \bar{r}_{H^*}(A) \cap R \quad \text{by assumption} \\ &= r_H(R) \quad \text{by part (1) .} \end{aligned}$$

Thus $\bar{r}_H(R) = r_H(R)$ by assumption.

(3) Sufficiency is obvious. Now we show the necessity. Since

$$r((R\#_\sigma H)/(r_H(R)\#_\sigma H)) \cong r(R/r_H(R)\#_{\sigma'} H) = 0,$$

we have $r(R\#_\sigma H) \subseteq r_H(R)\#_\sigma H$. Considering part (2), we have

$$r(R\#_\sigma H) = r_H(R)\#_\sigma H. \quad \square$$

Corollary 12.4.5. *Let r denote r_b, r_l, r_j, r_{bm} and r_n . Then*

$$(1) \bar{r}_H \leq r_H;$$

Furthermore, if H is a finite-dimensional Hopf algebra, then

$$(2) \bar{r}_H = r_H;$$

$$(3) R \#_\sigma H \text{ is } r\text{-semisimple for any } r_H\text{-semisimple } R \text{ iff } r(R \#_\sigma H) = r_H(R) \#_\sigma H;$$

$$(4) R \#_\sigma H \text{ is } r_j\text{-semisimple for any } r_{H_j}\text{-semisimple } R \text{ iff } r_j(R \#_\sigma H) = r_{H_j}(R) \#_\sigma H.$$

Proof. (1) When $r = r_b$ or $r = r_j$, it has been proved in [130, Proposition 2.3 (1) and 3.2 (1)] and in the preceding sections. The others can similarly be proved.

(2) It follows from Proposition 12.4.4 (2).

(3) and (4) follow from part (1) and Proposition 12.4.4 (3). $\square$

Proposition 12.4.6. *If $H = kG$ or the weak action of H on R is inner, then*

$$(1). r_H(R) = r(R);$$

$$(2) \text{ If, in addition, } H \text{ is a finite-dimensional Hopf algebra and } \bar{r}_H \leq r_H, \text{ then } r_H(R) = \bar{r}_H(R) = r(R).$$

Proof. (1) It is trivial.

(2) It immediately follows from part (1) and Proposition 12.4.1 (1) (2). $\square$

Theorem 12.4.7. *Let G be a finite group and $|G|^{-1} \in k$. If $H = kG$ or $H = (kG)^*$, then*

$$(1) r_j(R) = r_{H_j}(R) = r_{jH}(R);$$

$$(2) r_j(R \#_\sigma H) = r_{H_j}(R) \#_\sigma H.$$

Proof. (1) Let $H = kG$. We can easily check $r_j(R) = r_{jH}(R)$ using the method similar to the proof of [129, Proposition 4.6]. By [130, Proposition 3.3 (2)], $r_{H_j}(R) = r_{jH}(R)$. Now, we only need to show that

$$r_j(R) = r_{H^*j}(R).$$

We see that

$$\begin{aligned} r_j((R \#_\sigma H^*) \# H) &= r_{H^*j}((R \#_\sigma H^*) \# H) \quad \text{by [29, Theorem 4.4 (3)]} \\ &= r_{H_j}(R \#_\sigma H^*) \# H \quad \text{by [130, Proposition 3.3 (1)]} \\ &= (r_{H^*j}(R) \#_\sigma H^*) \# H \quad \text{by [130, Proposition 3.3 (1)]}. \end{aligned}$$

On the one hand, by [130, Lemma 2.2 (8)], $\Phi(r_j(R)) = r_j((R \#_\sigma H^*) \# H)$. On the other hand, we have that $\Phi(r_{H^*j}(R)) = (r_{H^*j}(R) \#_\sigma H^*) \# H$ by [130, Lemma 2.2 (2)]. Consequently, $r_j(R) = r_{H^*j}(R)$.

(2) It immediately follows from part (1) and [130, Proposition 3.3 (1) (2)]. $\square$

Corollary 12.4.8. *Let H be a semisimple and cosemisimple Hopf algebra over algebraically closed field k . If H is commutative or cocommutative, then*

$$r_j(R) = r_{H_j}(R) = r_{jH}(R) \quad \text{and} \quad r_j(R \#_\sigma H) = r_{H_j}(R) \#_\sigma H.$$

Proof. It immediately follows from Theorem 12.4.7 and [114, Lemma 8.0.1 (c)]. $\square$

We give an example to show that conditions in Corollary 12.4.8 can not be omitted.

Example 12.4.9. (see [130, Example P20]) Let k be a field of characteristic $p > 0$, $R = k[x]/(x^p)$. We can define a derivation on R by sending x to $x + 1$. Set $H = u(kd)$, the restricted enveloping algebra, and $A = R \# H$. Then

- (1) $r_b(A \# H^*) \neq r_{H^*b}(A) \# H^*$;
- (2) $r_j(A \# H^*) \neq r_{H^*j}(A) \# H^*$;
- (3) $r_j(A \# H^*) \not\subseteq r_{jH^*}(A) \# H^*$.

Proof. (1) By [130, Example P20], we have $r_b(R) \neq 0$ and $r_{bH}(R) = 0$. Since $\Phi(r_b(R)) = r_b(A \# H^*) \neq 0$ and $\Phi(r_{bH}(R)) = r_{bH^*}(A) \# H^* = 0$, we have that part (1) holds.

(3) We see that $r_j(A \# H^*) = \Phi(r_j(R))$ and $r_{Hj}(A) \# H^* = \Phi(r_{Hj}(R))$. Since R is commutative, $r_j(R) = r_b(R)$. Thus $r_{Hj}(R) = r_{jH}(R) = r_{bH}(R) = 0$ and $r_j(R) = r_b(R) \neq 0$, which implies $r_j(A \# H^*) \not\subseteq r_{jH^*}(A) \# H^*$.

(2) It follows from part (3). $\square$

This example also answer the question J.R. Fisher asked in [48] :

$$\text{Is } r_j(R \# H) \subseteq r_{jH}(R) \# H \quad ?$$

If F is an extension field of k , we write R^F for $R \otimes_k F$ (see [90, P49]) .

Lemma 12.4.10. If F is an extension field of k , then

- (1) H is a semisimple Hopf algebra over k iff H^F is a semisimple Hopf algebra over F ;
- (2) Furthermore, if H is a finite-dimensional Hopf algebra, then H is a cosemisimple Hopf algebra over k iff H^F is a cosemisimple Hopf algebra over F .

Proof. (1) It is clear that $\int_H^l \otimes F = \int_{H^F}^l$. Thus H is a semisimple Hopf algebra over k iff H^F is a semisimple Hopf algebra over F .

(2) $(H \otimes F)^* = H^* \otimes F$ since $H^* \otimes F \subseteq (H \otimes F)^*$ and $\dim_F(H \otimes F) = \dim_F(H^* \otimes F) = \dim_k H$. Thus we can obtain part (2) by Part (1). $\square$

By the way, if H is a semisimple Hopf algebra, then H is a separable algebra by Lemma 12.4.10 (see [95, P284]).

Proposition 12.4.11. Let F be an algebraic closure of k , R an algebra over k and

$$r(R \otimes_k F) = r(R) \otimes_k F.$$

If H is a finite-dimensional Hopf algebra with cocommutative coradical over k , then

$$r(R)^{\dim H} \subseteq r_H(R).$$

Proof. It is clear that H^F is a finite-dimensional Hopf algebra over F and $\dim H = \dim H^F = n$. Let H_0^F be the coradical of H^F , $H_1^F = H_0^F \wedge H_0^F$, $H_{i+1}^F = H_0^F \wedge H_i^F$ for $i = 1, \dots, n-1$.

Notice $H_0^F \subseteq H_0 \otimes F$. Thus H_0^F is cocommutative. It is clear that $H_0^F = kG$ by [114, Lemma 8.0.1 (c)] and $H^F = \cup H_i^F$. It is easy to show that if $k > i$, then

$$H_i^F \cdot (r(R^F))^k \subseteq r(R^F)$$

by induction for i . Thus

$$H^F \cdot (r(R^F))^{\dim H} \subseteq r(R^F),$$

which implies that $(r(R^F))^{\dim H} \subseteq r(R^F)_{H^F}$. By assumption, we have that $(r(R) \otimes F)^{\dim H} \subseteq (r(R) \otimes F)_{H^F}$. It is clear that $(I \otimes F)_{H^F} = I_H \otimes F$ for any ideal I of R . Consequently, $(r(R))^{\dim H} \subseteq r(R)_H$. $\square$

Theorem 12.4.12. *Let H be a semisimple, cosemisimple and either commutative or cocommutative Hopf algebra over k . If there exists an algebraic closure F of k such that*

$$r_j(R \otimes F) = r_j(R) \otimes F \quad \text{and} \quad r_j((R \#_\sigma H) \otimes F) = r_j(R \#_\sigma H) \otimes F,$$

then

- (1) $r_j(R) = r_{Hj}(R) = r_{jH}(R)$;
- (2) $r_j(R \#_\sigma H) = r_{Hj}(R) \#_\sigma H$.

Proof. (1). By Lemma 12.4.10, H^F is semisimple and cosemisimple. Considering Corollary 12.4.8, we have that $r_j(R^F) = r_{H^Fj}(R^F) = r_{jH^F}(R^F)$. On the one hand, by assumption, $r_j(R^F) = r_j(R) \otimes F$. On the other hand, $r_{jH^F}(R^F) = (r_j(R) \otimes F)_{H^F} = r_{jH}(R) \otimes F$. Thus $r_j(R) = r_{jH}(R)$.

(2). It immediately follows from part (1). $\square$

Considering Theorem 12.4.12 and [95, Theorem 7.2.13], we have

Corollary 12.4.13. *Let H be a semisimple, cosemisimple and either commutative or cocommutative Hopf algebra over k . If there exists an algebraic closure F of k such that F/k is separable and algebraic, then*

- (1) $r_j(R) = r_{Hj}(R) = r_{jH}(R)$;
- (2) $r_j(R \#_\sigma H) = r_{Hj}(R) \#_\sigma H$.

Lemma 12.4.14. (Szász [116])

$$r_j(R) = r_k(R)$$

holds in the following three cases:

- (1) *Every element in R is algebraic over k ([116, Proposition 31.2]);*
- (2) *The cardinality of k is strictly greater than the dimension of R and k is infinite ([116, Theorem 31.4]);*
- (3) *k is uncountable and R is finitely generated ([116, Proposition 31.5]).*

Proposition 12.4.15. *Let F be an extension of k . Then $r(R) \otimes F \subseteq r(R \otimes F)$, where r denotes r_b, r_k, r_l, r_n .*

Proof. When $r = r_n$, for any $x \otimes a \in r_n(R) \otimes F$ with $a \neq 0$, there exists $y \in R$ such that $x = xyx$. Thus $x \otimes a = (x \otimes a)(y \otimes a^{-1})(x \otimes a)$, which implies $r_n(R) \otimes F \subseteq r_n(R \otimes F)$.

Similarly, we can obtain the others. $\square$

Corollary 12.4.16. *Let H be a semisimple, cosemisimple and commutative or cocommutative Hopf algebra. If there exists an algebraic closure F of k such that F/k is a pure transcendental extension and one of the following three conditions holds:*

- (i) *every element in $R \#_\sigma H$ is algebraic over k ;*
- (ii) *the cardinality of k is strictly greater than the dimension of R and k is infinite;*
- (iii) *k is uncountable and R is finitely generated;*

then

- (1) $r_j(R) = r_{Hj}(R) = r_{jH}(R)$;
- (2) $r_j(R \#_\sigma H) = r_{Hj}(R) \#_\sigma H$;
- (3) $r_j(R) = r_k(R)$ and $r_j(R \#_\sigma H) = r_k(R \#_\sigma H)$.

Proof. First, we have that part (3) holds by Lemma 12.4.14. We next see that

$$\begin{aligned} r_j(R \otimes F) &\subseteq r_j(R) \otimes F \quad [95, \text{Theorem 7.3.4}] \\ &= r_k(R) \otimes F \quad \text{part (3)} \\ &\subseteq r_k(R \otimes F) \quad \text{proposition 12.4.15} \\ &\subseteq r_j(R \otimes F). \end{aligned}$$

Thus $r_j(R \otimes F) = r_j(R) \otimes F$. Similarly, we can show that $r_j((R \#_\sigma H) \otimes F) = r_j(R \#_\sigma H) \otimes F$.

Finally, using Theorem 12.4.12, we complete the proof. $\square$

12.5 The H -Von Neumann regular radical

In this section, we construct the H -von Neumann regular radical for H -module algebras and show that it is an H -radical property.

Definition 12.5.1. *Let $a \in R$. If $a \in (H \cdot a)R(H \cdot a)$, then a is called an H -von Neumann regular element, or an H -regular element in short. If every element of R is an H -regular, then R is called an H -regular module algebra, written as r_{Hn} - H -module algebra. I is an H -ideal of R and every element in I is H -regular, then I is called an H -regular ideal.*

Lemma 12.5.2. *If I is an H -ideal of R and $a \in I$, then a is H -regular in I iff a is H -regular in H .*

Proof. The necessity is clear.

Sufficiency: If $a \in (H \cdot a)R(H \cdot a)$, then there exist $h_i, h'_i \in H, b_i \in R$, such that

$$a = \sum (h_i \cdot a)b_i(h'_i \cdot a).$$

We see that

$$\begin{aligned}
a &= \sum_{i,j} [h_i \cdot ((h_j \cdot a)b_j(h'_j \cdot a))] b_i(h'_i \cdot a) \\
&= \sum_{i,j} [((h_i)_1 \cdot (h_j \cdot a))((h_i)_2 \cdot b_j)((h_i)_3 \cdot (h'_j \cdot a))] b_i(h'_i \cdot a) \\
&\in (H \cdot a)I(H \cdot a).
\end{aligned}$$

Thus a is an H -regular in I . $\square$

Lemma 12.5.3. *If $x - \sum_i (h_i \cdot x)b_i(h'_i \cdot x)$ is H -regular, then x is H -regular, where $x, b_i \in R, h_i, h'_i \in H$.*

Proof. Since $x - \sum_i (h_i \cdot x)b_i(h'_i \cdot x)$ is H -regular, there exist $g_i, g'_i \in H, c_i \in R$ such that

$$x - \sum_i (h_i \cdot x)b_i(h'_i \cdot x) = \sum_j (g_j \cdot (x - \sum_i (h_i \cdot x)b_i(h'_i \cdot x)))c_j(g'_j \cdot (x - \sum_i (h_i \cdot x)b_i(h'_i \cdot x))).$$

Consequently, $x \in (H \cdot x)R(H \cdot x)$. $\square$

Definition 12.5.4.

$$r_{Hn}(R) := \{a \in R \mid \text{the } H\text{-ideal } (a) \text{ generated by } a \text{ is } H\text{-regular}\}.$$

Theorem 12.5.5. $r_{Hn}(R)$ is an H -ideal of R .

Proof. We first show that $Rr_{Hn}(R) \subseteq r_{Hn}(R)$. For any $a \in r_{Hn}(R), x \in R$, we have that (xa) is H -regular since $(xa) \subseteq (a)$. We next show that $a - b \in r_{Hn}(R)$ for any $a, b \in r_{Hn}(R)$. For any $x \in (a - b)$, since $(a - b) \subseteq (a) + (b)$, we have that $x = u - v$ and $u \in (a), v \in (b)$. Say $u = \sum_i (h_i \cdot u)c_i(h'_i \cdot u)$ and $h_i, h'_i \in H, c_i \in R$. We see that

$$\begin{aligned}
x &= \sum_i (h_i \cdot x)c_i(h'_i \cdot x) \\
&= (u - v) - \sum_i (h_i \cdot (u - v))c_i(h'_i \cdot (u - v)) \\
&= -v - \sum_i [(h_i \cdot u)c_i(h'_i \cdot v) - (h_i \cdot v)c_i(h'_i \cdot u) + (h_i \cdot v)c_i(h'_i \cdot v)] \\
&\in (v).
\end{aligned}$$

Thus $x - \sum_i (h_i \cdot x)c_i(h'_i \cdot x)$ is H -regular and x is H -regular by Lemma 12.5.3. Therefore $a - b \in r_{Hn}(R)$. Obviously, $r_{Hn}(R)$ is H -stable. Consequently, $r_{Hn}(R)$ is an H -ideal of R . $\square$

Theorem 12.5.6. $r_{Hn}(R/r_{Hn}(R)) = 0$.

Proof. Let $\bar{R} = R/r_{Hn}(R)$ and $\bar{b} = b + r_{Hn}(R) \in r_{Hn}(R/r_{Hn}(R))$. It is sufficient to show that $b \in r_{Hn}(R)$. For any $a \in (b)$, it is clear that $\bar{a} \in (\bar{b})$. Thus there exist $h_i, h'_i \in H, \bar{c}_i \in \bar{R}$ such that

$$\bar{a} = \sum_i (h_i \cdot \bar{a})\bar{c}_i(h'_i \cdot \bar{a}) = \sum_i \overline{(h_i \cdot a)c_i(h'_i \cdot a)}.$$

Thus $a - \sum_i (h_i \cdot a)c_i(h'_i \cdot a) \in r_{Hn}(R)$, which implies that a is H -regular. Consequently, $b \in r_{Hn}(R)$. Namely, $\bar{b} = 0$ and $r_{Hn}(R) = 0$. $\square$

Corollary 12.5.7. r_{Hn} is an H -radical property for H -module algebras and $r_{nH} \leq r_{Hn}$.

Proof. (R1) If $R \stackrel{f}{\sim} R'$ and R is an r_{Hn} - H -module algebra, then for any $f(a) \in R'$, $f(a) \in (H \cdot f(a))R'(H \cdot f(a))$. Thus R' is also an r_{Hn} - H -module algebra.

(R2) If I is an r_{Hn} - H -ideal of R and $r_{Hn}(R) \subseteq I$ then, for any $a \in I$, (a) is H -regular since $(a) \subseteq I$. Thus $I \subseteq r_{Hn}(R)$.

(R3) It follows from Theorem 12.5.8.

Consequently r_{Hn} is an H -radical property for H -module algebras. It is straightforward to check $r_{nH} \leq r_{Hn}$. $\square$

r_{Hn} is called the H -von Neumann regular radical.

Theorem 12.5.8. If I is an H -ideal of R , then $r_{Hn}(I) = r_{Hn}(R) \cap I$. Namely, r_{Hn} is a strongly hereditary H -radical property.

Proof. By Lemma 12.5.2, $r_{Hn}(R) \cap I \subseteq r_{Hn}(I)$. Now, it is sufficient to show that $(x)_I = (x)_R$ for any $x \in r_{Hn}(I)$, where $(x)_I$ and $(x)_R$ denote the H -ideals generated by x in I and R respectively. Let $x = \sum (h_i \cdot x)b_i(h'_i \cdot x)$, where $h_i, h'_i \in H, b_i \in I$. We see that

$$\begin{aligned} R(H \cdot x) &= R(H \cdot (\sum (h_i \cdot x)b_i(h'_i \cdot x))) \\ &\subseteq R(H \cdot x)I(H \cdot x) \\ &\subseteq I(H \cdot x). \end{aligned}$$

Similarly,

$$(H \cdot x)R \subseteq (H \cdot x)I.$$

Thus $(x)_I = (x)_R$. $\square$

A graded algebra R of type G is said to be Gr-regular if for every homogeneous $a \in R_g$ there exists $b \in R$ such that $a = aba$ (see [92] P258). Now, we give the relations between Gr-regularity and H -regularity.

Theorem 12.5.9. If G is a finite group, R is a graded algebra of type G , and $H = (kG)^*$, then R is Gr-regular iff R is H -regular.

Proof. Let $\{p_g \mid g \in G\}$ be the dual base of base $\{g \mid g \in G\}$. If R is Gr-regular for any $a \in R$, then $a = \sum_{g \in G} a_g$ with $a_g \in R_g$. Since R is Gr-regular, there exist $b_{g^{-1}} \in R_{g^{-1}}$ such that $a_g = a_g b_{g^{-1}} a_g$ and

$$a = \sum_{g \in G} a_g = \sum_{g \in G} a_g b_{g^{-1}} a_g = \sum_{g \in G} (p_g \cdot a) b_{g^{-1}} (p_g \cdot a).$$

Consequently, R is H -regular.

Conversely, if R is H -regular, then for any $a \in R_g$, there exists $b_{x,y} \in R$ such that

$$a = \sum_{x,y \in G} (p_x \cdot a) c_{x,y} (p_y \cdot a).$$

Considering $a \in R_g$, we have that $a = ab_{g,g}a$. Thus R is Gr-regular. $\square$

12.6 About J.R. Fisher's question

In this section, we answer the question J.R. Fisher asked in [48]. Namely, we give a necessary and sufficient condition for validity of relation (2) .

Throughout this section, let k be a commutative ring with unit, R an H - module algebra and H a Hopf algebra over k .

Theorem 12.6.1. *Let $\mathcal{K}$ be an ordinary special class of rings and closed with respect to isomorphism. Set $r = r^{\mathcal{K}}$ and $\bar{r}_H(R) = r(R\#H) \cap R$ for any H -module algebra R . Then $\bar{r}_H$ is an H -radical property of H -module algebras. Furthermore, it is an H -special radical.*

Proof. Let $\bar{\mathcal{M}}_R = \{M \mid M \text{ is an } R\text{-prime module and } R/(0 : M)_R \in \mathcal{K}\}$ for any ring R and $\bar{\mathcal{M}} = \cup \bar{\mathcal{M}}_R$. Set $\mathcal{M}_R = \{M \mid M \in \bar{\mathcal{M}}_{R\#H}\}$ for any H - module algebra R and $\mathcal{M} = \cup \mathcal{M}_R$. It is straightforward to check that $\bar{\mathcal{M}}$ satisfies the conditions of [129, Proposition 4.3]. Thus $\mathcal{M}$ is an H -special module by [129, Proposition 4.3]. It is clear that $\mathcal{M}(R) = \bar{\mathcal{M}}(R\#H) \cap R = r(R\#H) \cap R$ for any H -module algebra R . Thus $\bar{r}_H$ is an H -special radical by [129, Theorem 3.1]. $\square$

Using the Theorem 12.6.1, we have that $\bar{r}_{bH}, \bar{r}_{lH}, \bar{r}_{kH}, \bar{r}_{jH}, \bar{r}_{bmH}$ are all H -special radicals.

Proposition 12.6.2. *Let $\mathcal{K}$ be a special class of rings and closed with respect to isomorphism. Set $r = r^{\mathcal{K}}$. Then*

- (1) $\bar{r}_H(R)\#H \subseteq r(R\#H)$;
- (2) $\bar{r}_H(R)\#H = r(R\#H)$ iff there exists an H -ideal I of R such that $r(R\#H) = I\#H$;
- (3) R is an $\bar{r}_H$ - H -module algebra iff $r(R\#H) = R\#H$;
- (4) I is an $\bar{r}_H$ - H -ideal of R iff $r(I\#H) = I\#H$;
- (5) $r(\bar{r}_H(R)\#H) = \bar{r}_H(R)\#H$;
- (6) $r(R\#H) = \bar{r}_H(R)\#H$ iff $r(\bar{r}_H(R)\#H) = r(R\#H)$.

Proof. (1) It is similar to the proof of Proposition 12.4.3.

(2) It is a straightforward verification.

(3) If R is an $\bar{r}_H$ -module algebra, then $R\#H \subseteq r(R\#H)$ by part (1). Thus $R\#H = r(R\#H)$. The sufficiency is obvious.

(4), (5) and (6) immediately follow from part (3) . $\square$

Theorem 12.6.3. *If R is an algebra over field k with unit and H is a Hopf algebra over field k , then*

- (1) $\bar{r}_{jH}(R) = r_{Hj}(R)$ and $r_j(r_{Hj}(R)\#H) = r_{Hj}(R)\#H$;
- (2) $r_j(R\#H) = r_{Hj}(R)\#H$ iff $r_j(r_{Hj}(R)\#H) = r_j(R\#H)$ iff $r_j(r_j(R\#H) \cap R\#H) = r_j(R\#H)$;
- (3) Furthermore, if H is finite-dimensional, then $r_j(R\#H) = r_{Hj}(R)\#H$ iff $r_j(r_{jH}(R)\#H) = r_j(R\#H)$.

Proof. (1) By [130, Proposition 3.1], we have $\bar{r}_{jH}(R) = r_{Hj}(R)$. Consequently, $r_j(r_{Hj}(R)\#H) = r_{Hj}(R)\#H$ by Proposition 12.6.2 (5).

(2) It immediately follows from part (1) and Proposition 12.6.2 (6).

(3) It can easily be proved by part (2) and [130, Proposition 3.3 (2)]. $\square$

The theorem answers the question J.R. Fisher asked in [48] : When is $r_j(R\#H) = r_{Hj}(R)\#H$?

Proposition 12.6.4. *If R is an algebra over field k with unit and H is a finite-dimensional Hopf algebra over field k , then*

(1) $\bar{r}_{bH}(R) = r_{Hb}(R) = r_{bH}(R)$ and $r_b(r_{Hb}(R)\#H) = r_{Hb}(R)\#H$;

(2) $r_b(R\#H) = r_{Hb}(R)\#H$ iff $r_b(r_{Hb}(R)\#H) = r_b(R\#H)$ iff $r_b(r_{bH}(R)\#H) = r_b(R\#H)$ iff $r_b(r_b(R\#H) \cap R\#H) = r_b(R\#H)$.

Proof. (1) By [130, Proposition 2.4], we have $\bar{r}_{bH}(R) = r_{Hb}(R)$. Thus $r_b(r_{Hb}(R)\#H) = r_{Hb}(R)\#H$ by Proposition 12.6.2 (5).

(2) It follows from part (1) and Proposition 12.6.2 (6) . $\square$

In fact, if H is commutative or cocommutative, then $S^2 = id_H$ by [114, Proposition 4.0.1], and H is semisimple and cosemisimple iff the character $chark$ of k does not divides $dimH$ (see [104, Proposition 2 (c)]). It is clear that if H is a finite-dimensional commutative or cocommutative Hopf algebra and the character $chark$ of k does not divides $dimH$, then H is a finite-dimensional semisimple, cosemisimple, commutative or cocommutative Hopf algebra. Consequently, the conditions in Corollary 12.4.8, Theorem 12.4.12, Corollary 12.4.13 and 12.4.16 can be simplified

Chapter 13

Classical Yang-Baxter Equation And Low Dimensional Triangular Lie Algebras

The concept and structures of Lie coalgebras were introduced and studied by W. Michaelis in [85] [86]. V.G.Drinfel'd and A.A.Belavin in [42] [10] introduced the notion of triangular, coboundary Lie bialgebra L associated to a solution $r \in L \otimes L$ of the CYBE and gave a classification of solutions of CYBE with parameter for simple Lie algebras. W. Michaelis in [84] obtained the structure of a triangular, coboundary Lie bialgebra on any Lie algebra containing linearly independent elements a and b satisfying $[a, b] = \alpha b$ for some non-zero $\alpha \in k$ by setting $r = a \otimes b - b \otimes a$.

The Yang-Baxter equation first came up in a paper by Yang as factorization condition of the scattering S-matrix in the many-body problem in one dimension and in the work of Baxter on exactly solvable models in statistical mechanics. It has been playing an important role in mathematics and physics (see [10] , [125]). Attempts to find solutions of The Yang-Baxter equation in a systematic way have led to the theory of quantum groups. The Yang-Baxter equation is of many forms. The classical Yang-Baxter equation is one of these.

In many applications one needs to know the solutions of classical Yang-Baxter equation and know if a Lie algebra is a coboundary Lie bialgebra or a triangular Lie bialgebra. A systematic study of low dimensional Lie algebras, specially, of those Lie algebras that play a role in physics (as e.g. $sl(2, \mathbf{C})$, or the Heisenberg algebra), is very useful.

In this chapter, we obtain all solutions of constant classical Yang-Baxter equation (CYBE) in Lie algebra L with $\dim L \leq 3$ and give the sufficient and necessary conditions for $(L, [\], \Delta_r, r)$ to be a coboundary (or triangular) Lie bialgebra. We find the strongly symmetric elements in $L \otimes L$ and show they are all solutions of CYBE in L with $\dim L \leq 3$. Using these conclusions, we study the Lie algebra $sl(2)$.

In order to make the chapter somewhat self-contained, we begin by recalling the definition of Lie bialgebra.

Let k be a field with $\text{char } k \neq 2$ and L be a Lie algebra over k . If $r = \sum a_i \otimes b_i \in L \otimes L$ and

$x \in L$, we define

$$\begin{aligned}
[r^{12}, r^{13}] &:= \sum_{i,j} [a_i, a_j] \otimes b_i \otimes b_j \\
[r^{12}, r^{23}] &:= \sum_{i,j} a_i \otimes [b_i, a_j] \otimes b_j \\
[r^{13}, r^{23}] &:= \sum_{i,j} a_i \otimes a_j \otimes [b_i, b_j] \\
x \cdot r &:= \sum_i [x, a_i] \otimes b_i + a_i \otimes [x, b_i] \\
\Delta_r(x) &:= x \cdot r
\end{aligned}$$

and call

$$[r^{12}, r^{13}] + [r^{12}, r^{23}] + [r^{13}, r^{23}] = 0$$

the classical Yang-Baxter equation (CYBE).

Let

$$\tau : L \otimes L \longrightarrow L \otimes L$$

denote the natural twist map (defined by $x \otimes y \mapsto y \otimes x$), and let

$$\xi : L \otimes L \otimes L \longrightarrow L \otimes L \otimes L$$

the map defined by $x \otimes y \otimes z \mapsto y \otimes z \otimes x$ for any $x, y, z \in L$.

A vector space L is called a Lie coalgebra, if there exists a linear map

$$\Delta : L \longrightarrow L \otimes L$$

such that

- (i) $Im \Delta \subseteq Im(1 - \tau)$ and
- (ii) $(1 + \xi + \xi^2)(1 \otimes \Delta)\Delta = 0$

A vector space $(L, [\], \Delta)$ is called a Lie bialgebra if

- (i) $(L, [\])$ is a Lie algebra;
- (ii) (L, Δ) is a Lie coalgebra; and
- (iii) for all $x, y \in L$,

$$\Delta[x, y] = x \cdot \Delta(y) - y \cdot \Delta(x),$$

where, for all $x, a_i, b_i \in L$,

$(L, [\], \Delta, r)$ is called a coboundary Lie bialgebra, if $(L, [\], \Delta)$ is a Lie bialgebra and $r \in Im(1 - \tau) \subseteq L \otimes L$ such that

$$\Delta(x) = x \cdot r$$

for all $x \in L$. A coboundary Lie bialgebra $(L, [\], \Delta, r)$ is called triangular, if r is a solution of CYBE.

Jacobson gave a classification of Lie algebras with their dimension $dim L \leq 3$ in [59, P 11-14]. We now write their whole operations as follows:

(I) If L is an abelian Lie algebra, then its operation is trivial;

(II) If $\dim L = \dim L' = 3$, then there exists a base $\{e_1, e_2, e_3\}$ and $\alpha, \beta \in k$ with $\alpha\beta \neq 0$ such that $[e_1, e_2] = e_3, [e_2, e_3] = \alpha e_1, [e_3, e_1] = \beta e_2$;

(III) If $\dim L = 3$ and $L' \subseteq$ the center of L with $\dim L' = 1$, then there exist a base $\{e_1, e_2, e_3\}$ and $\alpha, \beta \in k$ with $\alpha = \beta = 0$ such that $[e_1, e_2] = e_3, [e_2, e_3] = \alpha e_1, [e_3, e_1] = \beta e_2$;

(IV) If k is an algebraically closed field and $\dim L = 3$ with $\dim L' = 2$, then there exists a base $\{e_1, e_2, e_3\}$ and $\beta, \delta \in k$ with $\delta \neq 0$ such that $[e_1, e_2] = 0, [e_1, e_3] = e_1 + \beta e_2, [e_2, e_3] = \delta e_2$;

(V) If $\dim L = 3$ and $L' \not\subseteq$ the center of L , then there exists a base $\{e_1, e_2, e_3\}$ and $\beta, \delta \in k$ with $\beta = \delta = 0$ such that $[e_1, e_2] = 0, [e_1, e_3] = e_1 + \beta e_2, [e_2, e_3] = \delta e_2$;

(VI) If $\dim L = 2$ with $\dim L' = 1$, then there exists a base $\{e, f\}$ such that $[e, f] = e$.

Remark: In Case (IV), the condition which k is an algebraically closed field is required. In other word, A lie algebra L with $\dim L = 3$ and $\dim L' = 2$, it is possible that there is not the basis in (IV). We shall consider the general case in Section 6–9.

13.1 The solutions of CYBE

In this section, we find the general solution of CYBE for Lie algebra L with $\dim L \leq 3$.

Definition 13.1.1. Let $\{e_1, e_2, \dots, e_n\}$ be a base of vector space V and $r = \sum_{i,j=1}^n k_{ij}(e_i \otimes e_j) \in V \otimes V$, where $k_{ij} \in k$, for $i, j = 1, 2, \dots, n$.

(i) If

$$k_{ij} = k_{ji} \quad k_{ij}k_{lm} = k_{il}k_{jm}$$

for $i, j, l, m = 1, 2, \dots, n$, then r is called strongly symmetric to the base $\{e_1, e_2, \dots, e_n\}$;

(ii) If $k_{ij} = -k_{ji}$, for $i, j = 1, 2, \dots, n$, then r is called skew symmetric.

(iii) Let $\dim V = 3$, $k_{11} = x, k_{22} = y, k_{33} = z, k_{12} = p, k_{21} = q, k_{13} = s, k_{31} = t, k_{23} = u, k_{32} = v, \alpha, \beta \in k$. If $p = -q, s = -t, u = -v, x = \alpha z, y = \beta z$ and

$$\alpha\beta z^2 + \beta s^2 + \alpha u^2 + p^2 = 0,$$

then r is called α, β -skew symmetric to the base $\{e_1, e_2, e_3\}$.

Obviously, skew symmetry does not depend on the particular choice of bases of V . We need to know if strong symmetry depends on the particular choice of bases of V .

Lemma 13.1.2. (I) If V is a finite-dimensional vector space, then the strong symmetry does not depend on the particular choice of bases of V ;

(II) Let $\{e_1, e_2, e_3\}$ be a base of vector space V , $p, q, s, t, u, v, x, y, z, \alpha, \beta \in k$ and $r = p(e_1 \otimes e_2) + q(e_2 \otimes e_1) + s(e_1 \otimes e_3) + t(e_3 \otimes e_1) + u(e_2 \otimes e_3) + v(e_3 \otimes e_2) + x(e_1 \otimes e_1) + y(e_2 \otimes e_2) + z(e_3 \otimes e_3)$.

Then r is strongly symmetric iff

$$xy = p^2, xz = s^2, yz = u^2, xu = sp, ys = pu, zp = su, p = q, s = t, u = v$$

iff

$$xy = p^2, xz = s^2, yz = u^2, xu = sp, p = q, s = t, u = v$$

iff

$$xy = p^2, xz = s^2, yz = u^2, ys = up, p = q, s = t, u = v$$

iff

$$xy = p^2, xz = s^2, yz = u^2, zp = su, p = q, s = t, u = v$$

iff

$$\begin{aligned} r &= z^{-1}su(e_1 \otimes e_2) + z^{-1}su(e_2 \otimes e_1) + s(e_1 \otimes e_3) + s(e_3 \otimes e_1) + u(e_2 \otimes e_3) + u(e_3 \otimes e_2) + \\ &z^{-1}s^2(e_1 \otimes e_1) + z^{-1}u^2(e_2 \otimes e_2) + z(e_3 \otimes e_3) \text{ with } z \neq 0; \text{ or} \\ r &= p(e_1 \otimes e_2) + p(e_2 \otimes e_1) + x(e_1 \otimes e_1) + x^{-1}p^2(e_2 \otimes e_2) \text{ with } x \neq 0; \text{ or} \\ r &= y(e_2 \otimes e_2). \end{aligned}$$

Proof. (I) Let $\{e_1, e_2, \dots, e_n\}$ and $\{e'_1, e'_2, \dots, e'_n\}$ are two bases of V and $r = \sum_{i,j=1}^n k_{ij}(e_i \otimes e_j)$ be strongly symmetric to the base $\{e_1, e_2, \dots, e_n\}$. It is sufficient to show that r is strongly symmetric to the base $\{e'_1, e'_2, \dots, e'_n\}$. Obviously, there exists $q_{ij} \in k$ such that $e_i = \sum_s e'_s q_{si}$ for $i = 1, 2, \dots, n$. By computation, we have that

$$r = \sum_{s,t} \left(\sum_{i,j} k_{ij} q_{si} q_{tj} \right) (e_s \otimes e_t).$$

If we set $k'_{st} = \sum_{i,j} k_{ij} q_{si} q_{tj}$, then for $l, m, u, v = 1, 2, \dots, n$, we have that

$$\begin{aligned} k'_{lm} k'_{uv} &= \sum_{i,j,s,t} k_{ij} k_{st} q_{li} q_{mj} q_{us} q_{vt} \\ &= \sum_{i,j,s,t} k_{is} k_{jt} q_{li} q_{mj} q_{us} q_{vt} \quad (\text{by } k_{ij} k_{st} = k_{is} k_{jt}) \\ &= k'_{lu} k'_{mv} \end{aligned}$$

and

$$k'_{lm} = k'_{ml},$$

which implies that r is strongly symmetric to the base $\{e'_1, e'_2, \dots, e'_n\}$.

(II) We only show that if

$$xy = p^2, xz = s^2, yz = u^2, xu = sp, p = q, s = t, u = v,$$

then

$$ys = pu, zp = su.$$

If $p \neq 0$, then $ysp = yxu = up^2$ and $usp = uxu = xyz = zp^2$, which implies $ys = up$ and $us = zp$. If $p = 0$, then $x = 0$ or $y = 0$, which implies $s = 0$ or $u = 0$. Consequently, $ys = up$ and $us = zp$. $\square$

Proposition 13.1.3. *Let L be a Lie algebra with $\dim L = 2$. Then r is a solution of CYBE iff r is strongly symmetric or skew symmetric.*

Proof. It is not hard since the tensor r has only 4 coefficients. $\square$

Proposition 13.1.4. *Let L be a Lie algebra with a basis $\{e_1, e_2, e_3\}$ such that $[e_1, e_2] = e_3$, $[e_2, e_3] = \alpha e_1$, $[e_3, e_1] = \beta e_2$, where $\alpha, \beta \in k$. Let $p, q, s, t, u, v, x, y, z \in k$.*

(I) If r is strongly symmetric or α, β -skew symmetric to basis $\{e_1, e_2, e_3\}$, then r is a solution of CYBE;

(II) If $\alpha \neq 0, \beta \neq 0$, then r is a solution of CYBE in L iff r is strongly symmetric or α, β -skew symmetric;

(III) If $\alpha = \beta = 0$, then r is a solution of CYBE in L iff

$$r = p(e_1 \otimes e_2) + p(e_2 \otimes e_1) + s(e_1 \otimes e_3) + t(e_3 \otimes e_1) + u(e_2 \otimes e_3) + v(e_3 \otimes e_2) + x(e_1 \otimes e_1) + y(e_2 \otimes e_2) + z(e_3 \otimes e_3)$$

with $p \neq 0, p^2 = xy, xu = sp, xv = tp, tu = vs$, or

$$r = s(e_1 \otimes e_3) + t(e_3 \otimes e_1) + u(e_2 \otimes e_3) + v(e_3 \otimes e_2) + x(e_1 \otimes e_1) + y(e_2 \otimes e_2) + z(e_3 \otimes e_3)$$

with $xy = xu = xv = ys = yt = 0$ and $tu = vs$.

Proof . Let $r = \sum_{i,j=1}^3 k_{ij}(e_i \otimes e_j) \in L \otimes L$ and $k_{ij} \in k$ with $i, j = 1, 2, 3$. It is clear that

$$\begin{aligned} [r^{12}, r^{13}] &= \sum_{i,j=1}^3 \sum_{s,t=1}^3 k_{ij} k_{st} [e_i, e_s] \otimes e_j \otimes e_t \\ [r^{12}, r^{23}] &= \sum_{i,j=1}^3 \sum_{s,t=1}^3 k_{ij} k_{st} e_i \otimes [e_j, e_s] \otimes e_t \\ [r^{13}, r^{23}] &= \sum_{i,j=1}^3 \sum_{s,t=1}^3 k_{ij} k_{st} e_i \otimes e_s \otimes [e_j, e_t]. \end{aligned}$$

By computation, for $i, j, n = 1, 2, 3$, we have that the coefficient of $e_j \otimes e_i \otimes e_i$ in $[r^{12}, r^{13}]$ is zero and $e_i \otimes e_i \otimes e_j$ in $[r^{13}, r^{23}]$ is zero.

We now see the coefficient of $e_i \otimes e_j \otimes e_n$

in $[r^{12}, r^{13}] + [r^{12}, r^{23}] + [r^{13}, r^{23}]$.

- (1) $e_1 \otimes e_1 \otimes e_1(\alpha k_{12} k_{31} - \alpha k_{13} k_{21})$;
- (2) $e_2 \otimes e_2 \otimes e_2(-\beta k_{21} k_{32} + \beta k_{23} k_{12})$;
- (3) $e_3 \otimes e_3 \otimes e_3(k_{31} k_{23} - k_{32} k_{13})$;
- (4) $e_1 \otimes e_2 \otimes e_3(\alpha k_{22} k_{33} - \alpha k_{32} k_{23} - \beta k_{11} k_{33} + \beta k_{13} k_{13} + k_{11} k_{22} - k_{12} k_{21})$;
- (5) $e_2 \otimes e_3 \otimes e_1(\beta k_{33} k_{11} - \beta k_{13} k_{31} - k_{22} k_{11} + k_{21} k_{21} + \alpha k_{22} k_{33} - \alpha k_{23} k_{32})$;
- (6) $e_3 \otimes e_1 \otimes e_2(k_{11} k_{22} - k_{21} k_{12} - \alpha k_{33} k_{22} + \alpha k_{32} k_{32} + \beta k_{33} k_{11} - \beta k_{31} k_{13})$;
- (7) $e_1 \otimes e_3 \otimes e_2(-\alpha k_{33} k_{22} + \alpha k_{23} k_{32} + k_{11} k_{22} - k_{12} k_{12} - \beta k_{11} k_{33} + \beta k_{13} k_{31})$;
- (8) $e_3 \otimes e_2 \otimes e_1(-k_{22} k_{11} + k_{12} k_{21} + \beta k_{33} k_{11} - \beta k_{31} k_{31} - \alpha k_{33} k_{22} + \alpha k_{32} k_{23})$;
- (9) $e_2 \otimes e_1 \otimes e_3(-\beta k_{11} k_{33} + \beta k_{31} k_{13} + \alpha k_{22} k_{33} - \alpha k_{23} k_{23} - k_{22} k_{11} + k_{21} k_{12})$;
- (10) $e_1 \otimes e_1 \otimes e_2(-\alpha k_{31} k_{22} + \alpha k_{21} k_{32} + \alpha k_{12} k_{32} - \alpha k_{13} k_{22})$;
- (11) $e_2 \otimes e_1 \otimes e_1(-\alpha k_{23} k_{21} + \alpha k_{22} k_{31} + \alpha k_{22} k_{13} - \alpha k_{23} k_{12})$;
- (12) $e_1 \otimes e_1 \otimes e_3(\alpha k_{21} k_{33} - \alpha k_{31} k_{23} + \alpha k_{12} k_{33} - \alpha k_{13} k_{23})$;
- (13) $e_3 \otimes e_1 \otimes e_1(\alpha k_{32} k_{31} - \alpha k_{33} k_{21} + \alpha k_{32} k_{13} - \alpha k_{33} k_{12})$;
- (14) $e_2 \otimes e_2 \otimes e_1(-\beta k_{12} k_{31} + \beta k_{32} k_{11} + \beta k_{23} k_{11} - \beta k_{21} k_{31})$;
- (15) $e_1 \otimes e_2 \otimes e_2(-\beta k_{11} k_{32} + \beta k_{13} k_{12} - \beta k_{11} k_{23} + \beta k_{13} k_{21})$;

- (16) $e_2 \otimes e_2 \otimes e_3(\beta k_{32}k_{13} - \beta k_{12}k_{33} - \beta k_{21}k_{33} + \beta k_{23}k_{13});$
- (17) $e_3 \otimes e_2 \otimes e_2(-\beta k_{31}k_{32} + \beta k_{33}k_{12} + \beta k_{33}k_{21} - \beta k_{31}k_{23});$
- (18) $e_3 \otimes e_3 \otimes e_1(k_{13}k_{21} - k_{23}k_{11} + k_{31}k_{21} - k_{32}k_{11});$
- (19) $e_3 \otimes e_3 \otimes e_2(-k_{23}k_{12} + k_{13}k_{22} - k_{32}k_{12} + k_{31}k_{22});$
- (20) $e_2 \otimes e_3 \otimes e_3(k_{21}k_{23} - k_{22}k_{13} + k_{21}k_{32} - k_{22}k_{31});$
- (21) $e_1 \otimes e_3 \otimes e_3(-k_{12}k_{13} + k_{11}k_{23} - k_{12}k_{31} + k_{11}k_{32});$
- (22) $e_1 \otimes e_3 \otimes e_1(\alpha k_{23}k_{31} - \alpha k_{33}k_{21} - k_{12}k_{11} + k_{11}k_{21} + \alpha k_{12}k_{33} - \alpha k_{13}k_{32});$
- (23) $e_1 \otimes e_2 \otimes e_1(-\alpha k_{32}k_{21} + \alpha k_{22}k_{31} - \beta k_{11}k_{31} + \beta k_{13}k_{11} - \alpha k_{13}k_{22} + \alpha k_{12}k_{23});$
- (24) $e_2 \otimes e_1 \otimes e_2(\beta k_{31}k_{12} - \beta k_{11}k_{32} + \alpha k_{22}k_{32} - \alpha k_{23}k_{22} - \beta k_{21}k_{13} + \beta k_{23}k_{11});$
- (25) $e_2 \otimes e_3 \otimes e_2(-\beta k_{13}k_{32} + \beta k_{33}k_{12} + k_{21}k_{22} - k_{22}k_{12} - \beta k_{21}k_{33} + \beta k_{23}k_{31});$
- (26) $e_3 \otimes e_2 \otimes e_3(k_{12}k_{23} - k_{22}k_{13} - \beta k_{31}k_{33} + \beta k_{33}k_{13} + k_{31}k_{22} - k_{32}k_{21});$
- (27) $e_3 \otimes e_1 \otimes e_3(-k_{21}k_{13} + k_{11}k_{23} + \alpha k_{32}k_{33} - \alpha k_{33}k_{23} + k_{31}k_{12} - k_{32}k_{11}).$

Let $k_{11} = x, k_{22} = y, k_{33} = z, k_{12} = p, k_{21} = q, k_{13} = s, k_{31} = t, k_{23} = u, k_{32} = v$.

It follows from (1)–(27) that

- (28) $\alpha pt = \alpha qs;$
- (29) $\beta qv = \beta pu;$
- (30) $tu = vs;$
- (31) $\alpha yz - \beta xz + xy - \alpha uv + \beta s^2 - pq = 0;$
- (32) $\beta zx - yx + \alpha yz - \beta st + q^2 - \alpha uv = 0;$
- (33) $xy - \alpha zy + \beta zx - pq + \alpha v^2 - \beta st = 0;$
- (34) $-\alpha zy + xy - \beta xz + \alpha uv - p^2 + \beta st = 0;$
- (35) $-xy + \beta xz - \alpha yz + pq - \beta t^2 + \alpha uv = 0;$
- (36) $-\beta xz + \alpha yz - yx + \beta st - \alpha u^2 + pq = 0;$
- (37) $\alpha(-ty + qv + pv - sy) = 0;$
- (38) $\alpha(-uq + yt + ys - up) = 0;$
- (39) $\alpha(qz - tu + pz - su) = 0;$
- (40) $\alpha(vt - zq + vs - zp) = 0;$
- (41) $\beta(-pt + vx + ux - qt) = 0;$
- (42) $\beta(-xv + sp - xu + sq) = 0;$
- (43) $\beta(vs - pz + us - qz) = 0;$
- (44) $\beta(-tv + zp + zq - tu) = 0;$
- (45) $sq - ux + tq - vx = 0;$
- (46) $-up + sy - vp + ty = 0;$
- (47) $qu - ys + qv - yt = 0;$
- (48) $-ps + xu - pt + xv = 0;$
- (49) $\alpha ut - \alpha zq - px + xq + \alpha pz - \alpha sv = 0;$
- (50) $-\alpha vq + \alpha yt - \beta xt + \beta sx - \alpha sy + \alpha pu = 0;$
- (51) $\beta tp - \beta xv + \alpha yv - \alpha uy - \beta qs + \beta ux = 0;$
- (52) $-\beta sv + \beta zp + qy - yp - \beta qz + \beta ut = 0;$
- (53) $pu - ys - \beta tz + \beta zs + ty - vq = 0;$
- (54) $-qs + xu + \alpha vz - \alpha zu + tp - vx = 0.$

It is clear that r is the solution of CYBE iff relations (28)–(54) hold.

(I) By computation, we have that if r is strongly symmetric or α, β -skew symmetric, then relations (28)–(54) hold and so r is a solution of CYBE;

(II) Let $\alpha \neq 0$ and $\beta \neq 0$. By part (I), we only need show that if r is a solution of CYBE, then r is strongly symmetric or α, β -skew symmetric.

By computation, we have

$$(55) \quad q^2 = p^2 \text{ (by (34) + (32));}$$

$$(56) \quad u^2 = v^2 \text{ (by (33) + (36));}$$

$$(57) \quad t^2 = s^2 \text{ (by (31) + (35)).}$$

(a) If $s = t \neq 0$, then we have that

$$p = q \text{ by (28);}$$

$$u = v \text{ by (30);}$$

$$(58) \quad yz = u^2 \text{ by (35) - (32);}$$

$$(59) \quad xz = s^2 \text{ by (32) + (33);}$$

$$(60) \quad xy = p^2 \text{ by (36) - (31);}$$

$$(61) \quad xu = sp \text{ by (41).}$$

Thus r is strongly symmetric.

Similarly, we can show that if $p = q \neq 0$, or $u = v \neq 0$, then r is strongly symmetric.

(b) If $s = -t \neq 0$, then we have that

$$p = -q \text{ by (28);}$$

$$u = -v \text{ by (30);}$$

$$y = \beta z \text{ by (53);}$$

$$x = \alpha z \text{ by (50);}$$

$$\alpha\beta z^2 + \alpha u^2 + \beta s^2 + p^2 = 0 \text{ by (31).}$$

Then r is α, β -skew symmetric.

Similarly, we have that if $p = -q \neq 0$ or $u = -v \neq 0$, then r is α, β -skew symmetric.

(c) If $s = t = u = v = p = q = 0$, then we have that

$$xz = 0 \text{ by (32) + (33);}$$

$$yz = 0 \text{ by (32) + (31);}$$

$$xy = 0 \text{ by (31).}$$

Thus r is strongly symmetric.

(III) Let $\alpha = \beta = 0$. It is clear that the system of equations (28)–(54) is equivalent to the

below

$$\begin{cases} tu = vs, q = p, p^2 = xy \\ (s+t)p - (u+v)x = 0 \\ (s+t)y - (u+v)p = 0 \\ (t-s)y + (u-v)p = 0 \\ (t-s)p + (u-v)x = 0 \end{cases}$$

It is easy to check that if r is one case in part (III), then the system of equations hold. Conversely, if r is a solution of CYBE, we shall show that r is one of the two cases in part (III). If $p \neq 0$, then r is the first case in part (III). If $p = 0$, then r is the second case in part (III). $\square$

In particular, Proposition 13.1.4 implies:

Example 13.1.5. *Let*

$$sl(2) := \{x \mid x \text{ is a } 2 \times 2 \text{ matrix with trace zero over } k\}$$

and

$$e_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, e_2 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, e_3 = \begin{pmatrix} 2 & 0 \\ 0 & -2 \end{pmatrix}.$$

Thus L is a Lie algebra (defined by $[x, y] = xy - yx$) with a basis $\{e_1, e_2, e_3\}$. It is clear that

$$[e_1, e_2] = e_3, [e_2, e_3] = 4e_1, [e_3, e_1] = -4e_2.$$

Consequently, r is a solution of CYBE iff r is strongly symmetric or 4-, -4-skew symmetric to the basis $\{e_1, e_2, e_3\}$.

Proposition 13.1.6. *Let L be a Lie algebra with a basis $\{e_1, e_2, e_3\}$ such that $[e_1, e_2] = 0, [e_1, e_3] = e_1 + \beta e_2, [e_2, e_3] = \delta e_2$, where $\beta, \delta \in k$. Let $p, q, s, t, u, v, x, y, z \in k$.*

(I) *If r is strongly symmetric, then r is a solution of CYBE.*

(II) *If $\beta = 0, \delta \neq 0$, then r is a solution of CYBE in L iff r is strongly symmetric, or*

$$r = p(e_1 \otimes e_2) + q(e_2 \otimes e_1) + s(e_1 \otimes e_3) - s(e_3 \otimes e_1) + u(e_2 \otimes e_3) - u(e_3 \otimes e_2) + x(e_1 \otimes e_1) + y(e_2 \otimes e_2)$$

with $xu = xs = ys = yu = (1 - \delta)us = (1 + \delta)s(q + p) = (1 + \delta)u(q + p) = 0$

(III) *If $\beta \neq 0$ and $\delta = 1$, then r is a solution of CYBE in L iff r is strongly symmetric, or*

$$r = p(e_1 \otimes e_2) + q(e_2 \otimes e_1) + u(e_2 \otimes e_3) - u(e_3 \otimes e_2) + x(e_1 \otimes e_1) + y(e_2 \otimes e_2)$$

$$\text{with } xu = yu = u(q + p) = 0$$

(IV) *If $\beta = \delta = 0$, then r is a solution of CYBE in L iff*

$$r = p(e_1 \otimes e_2) + q(e_2 \otimes e_1) + s(e_1 \otimes e_3) + s(e_3 \otimes e_1) + u(e_2 \otimes e_3) + v(e_3 \otimes e_2) + x(e_1 \otimes e_1) + y(e_2 \otimes e_2) + z(e_3 \otimes e_3)$$

$$\text{with } z \neq 0, zp = vs, zq = us, zx = s^2, \text{ or}$$

$$r = p(e_1 \otimes e_2) + q(e_2 \otimes e_1) + s(e_1 \otimes e_3) - s(e_3 \otimes e_1) + u(e_2 \otimes e_3) + v(e_3 \otimes e_2) + x(e_1 \otimes e_1) + y(e_2 \otimes e_2)$$

$$\text{with } us = vs = xs = xu = xv = 0, up = qv, s(p + q) = 0.$$

Proof. Let $r = \sum_{i,j=1}^3 k_{ij}(e_i \otimes e_j) \in L \otimes L$ and $k_{ij} \in k$ with $i, j = 1, 2, 3$. By computation, for all $i, j, n = 1, 2, 3$, we have that the coefficient of $e_j \otimes e_i \otimes e_i$ in $[r^{12}, r^{13}]$ is zero and $e_i \otimes e_i \otimes e_j$ in $[r^{13}, r^{23}]$ is zero.

We now see the coefficient of $e_i \otimes e_j \otimes e_n$

in $[r^{12}, r^{13}] + [r^{12}, r^{23}] + [r^{13}, r^{23}]$.

$$(1) e_1 \otimes e_1 \otimes e_1(-k_{13}k_{11} + k_{11}k_{31});$$

$$(2) e_2 \otimes e_2 \otimes e_2(-\beta k_{23}k_{12} + \beta k_{21}k_{32} - \delta k_{23}k_{22} + \delta k_{22}k_{32});$$

$$(3) e_3 \otimes e_3 \otimes e_3(0);$$

$$(4) e_1 \otimes e_2 \otimes e_3(-k_{32}k_{13} + k_{12}k_{33} - \beta k_{13}k_{13} + \beta k_{11}k_{33} - \delta k_{13}k_{23} + \delta k_{12}k_{33});$$

$$(5) e_2 \otimes e_3 \otimes e_1(-\beta k_{33}k_{11} + \beta k_{13}k_{31} - \delta k_{33}k_{21} + \delta k_{23}k_{31} - k_{23}k_{31} + k_{21}k_{33});$$

$$(6) e_3 \otimes e_1 \otimes e_2(-k_{33}k_{12} + k_{31}k_{32} - \beta k_{33}k_{11} + \beta k_{31}k_{13} - \delta k_{33}k_{12} + \delta k_{32}k_{13});$$

$$(7) e_1 \otimes e_3 \otimes e_2(-k_{33}k_{12} + k_{13}k_{32} - \beta k_{13}k_{32} + \beta k_{11}k_{33} - \delta k_{13}k_{32} + \delta k_{12}k_{33});$$

$$(8) e_3 \otimes e_2 \otimes e_1(-\beta k_{33}k_{11} + \beta k_{31}k_{31} - \delta k_{33}k_{21} + \delta k_{32}k_{31} - k_{33}k_{21} + k_{31}k_{23});$$

$$(9) e_2 \otimes e_1 \otimes e_3(-\beta k_{31}k_{13} + \beta k_{11}k_{33} - \delta k_{31}k_{23} + \delta k_{21}k_{33} - k_{23}k_{13} + k_{21}k_{33});$$

$$(10) e_1 \otimes e_1 \otimes e_2(-k_{31}k_{12} + k_{11}k_{32} - k_{13}k_{12} + k_{11}k_{32});$$

$$(11) e_2 \otimes e_1 \otimes e_1(-k_{23}k_{11} + k_{21}k_{31} - k_{23}k_{11} + k_{21}k_{13});$$

$$(12) e_1 \otimes e_1 \otimes e_3(-k_{31}k_{13} + k_{11}k_{33} - k_{13}k_{13} + k_{11}k_{33});$$

$$(13) e_3 \otimes e_1 \otimes e_1(-k_{33}k_{11} + k_{31}k_{31} - k_{33}k_{11} + k_{31}k_{13});$$

$$(14) e_2 \otimes e_2 \otimes e_1(-\beta k_{32}k_{11} + \beta k_{12}k_{31} - \delta k_{32}k_{21} + \delta k_{22}k_{31} - \beta k_{23}k_{11} + \beta k_{21}k_{31} - \delta k_{23}k_{21} + \delta k_{22}k_{31});$$

$$(15) e_1 \otimes e_2 \otimes e_2(-\beta k_{13}k_{12} + \beta k_{11}k_{32} - \delta k_{13}k_{22} + \delta k_{12}k_{32} - \beta k_{13}k_{21} + \beta k_{11}k_{23} - \delta k_{13}k_{22} + \delta k_{12}k_{23});$$

$$(16) e_2 \otimes e_2 \otimes e_3(-\beta k_{32}k_{13} + \beta k_{12}k_{33} - \delta k_{32}k_{23} + \delta k_{22}k_{33} - \beta k_{23}k_{13} + \beta k_{21}k_{33} - \delta k_{23}k_{23} + \delta k_{22}k_{33});$$

$$(17) e_3 \otimes e_2 \otimes e_2(-\beta k_{33}k_{12} + \beta k_{31}k_{32} - \delta k_{33}k_{22} + \delta k_{32}k_{32} - \beta k_{33}k_{21} + \beta k_{31}k_{23} - \delta k_{33}k_{22} + \delta k_{32}k_{23});$$

$$(18) e_3 \otimes e_3 \otimes e_1(0);$$

$$(19) e_1 \otimes e_3 \otimes e_3(0);$$

$$(20) e_3 \otimes e_3 \otimes e_2(0);$$

$$(21) e_2 \otimes e_3 \otimes e_3(0);$$

$$(22) e_1 \otimes e_3 \otimes e_1(-k_{33}k_{11} + k_{13}k_{31} - k_{13}k_{31} + k_{11}k_{33}) = e_1 \otimes e_3 \otimes e_1(0);$$

$$(23) e_1 \otimes e_2 \otimes e_1(-k_{32}k_{11} + k_{12}k_{31} - \beta k_{13}k_{11} + \beta k_{11}k_{31} - \delta k_{13}k_{21} + \delta k_{12}k_{31} - k_{13}k_{21} + k_{11}k_{23});$$

$$(24) e_2 \otimes e_1 \otimes e_2(-\beta k_{31}k_{12} + \beta k_{11}k_{32} - \delta k_{31}k_{22} + \delta k_{21}k_{32} - k_{23}k_{12} + k_{21}k_{32} - \beta k_{23}k_{11} + \beta k_{21}k_{13} - \delta k_{23}k_{12} + \delta k_{22}k_{13});$$

$$(25) e_2 \otimes e_3 \otimes e_2(-\beta k_{33}k_{12} + \beta k_{13}k_{32} - \delta k_{33}k_{22} + \delta k_{23}k_{32} - \beta k_{23}k_{31} + \beta k_{21}k_{33} + \delta k_{33}k_{22} - \delta k_{22}k_{33});$$

$$(26) e_3 \otimes e_2 \otimes e_3(-\beta k_{33}k_{13} + \beta k_{31}k_{33} - \delta k_{33}k_{23} + \delta k_{32}k_{33});$$

$$(27) e_3 \otimes e_1 \otimes e_3(-k_{33}k_{13} + k_{31}k_{33});$$

Let $k_{11} = x, k_{22} = y, k_{33} = z, k_{12} = p, k_{21} = q, k_{13} = s, k_{31} = t, k_{23} = u, k_{32} = v$.

It follows from (1)–(27) that

$$(28) -sx + xt = 0;$$

$$(29) -\beta up + \beta qv - \delta uy + \delta yv = 0;$$

$$(30) -vs + pz - \beta s^2 + \beta xz - \delta su + \delta zp = 0;$$

$$(31) -\beta xz + \beta st - \delta zq + \delta ut - ut + qz = 0;$$

$$(32) -zp + tv - \beta zx + \beta ts - \delta zp + \delta vs = 0;$$

$$(33) -zp + sv - \beta st + \beta xz - \delta sv + \delta zp = 0;$$

$$(34) -\beta zx + \beta tt - \delta zq + \delta vt - zq + tu = 0;$$

$$(35) -\beta st + \beta xz - \delta tu + \delta qz - us + qz = 0;$$

$$(36) -tp + xv - sp + vx = 0;$$

$$(37) -ux + qt - ux + qs = 0;$$

$$(38) -st + xz - s^2 + xz = 0;$$

$$(39) -zx + tt - zx + st = 0;$$

$$(40) -\beta vx + \beta pt - \delta vq + \delta yt - \beta ux + \beta qt - \delta uq + \delta yt = 0;$$

$$(41) -\beta sp + \beta xv - \delta sy + \delta pv - \beta sq + \beta xu - \delta sy + \delta pu = 0;$$

$$(42) -\beta vs + \beta pz - \delta vu + \delta yz - \beta su + \beta zq - \delta u^2 + \delta yz = 0;$$

$$(43) -\beta zp + \beta tv - \delta zy + \delta vv - \beta zq + \beta tu - \delta zy + \delta vu = 0;$$

$$(44) -vx + pt - \beta sx + \beta xt - \delta sq + \delta pt - sq + xu = 0;$$

$$(45) -\beta pt + \beta vx - \delta ty + \delta qv - up + qv - \beta ux + \beta qs - \delta up + \delta ys = 0;$$

$$(46) -\beta zp + \beta sv - \beta ut + \beta qz = 0;$$

$$(47) -\beta zs + \beta tz - \delta zu + \delta vz = 0;$$

$$(48) -zs + tz = 0.$$

It is clear that r is a solution of CYBE iff (28)–(48) hold.

(I) It is trivial.

(II) Let $\beta = 0$ and $\delta \neq 0$. We only show that if r is a solution of CYBE, then r is one of the two cases in part (II). If $z \neq 0$, we have that $u = v, s = t, yz = u^2, xz = s^2$ and $ux = sp$ by (47), (48), (42), (38) and (36) respectively. It follows from (30) and (33) that $pz = us$ and from (31) and (34) that $qz = us$. Thus $q = p$ and $xy = p^2$ since $xyz = xu^2 = spu = zp^2$. By Lemma 13.1.2, r is strongly symmetric.

If $z = 0$, we have that $u = -v$ and $s = -t$ by (43), (42), (38) and (39). It follows from (28), (29), (36), (40), (30), (44) and (45) that $sx = 0, yu = 0, xv = 0, ys = 0, (1 - \delta)us = 0, (1 + \delta)s(p + q) = 0$ and $(1 + \delta)u(p + q) = 0$ respectively. Consequently, r is the second case.

(III) Let $\beta \neq 0, \delta = 1$. We only show that if r is a solution of CYBE, then r is one case in part (III). If $z \neq 0$, we have that $t = s, u = v, q = p, xz = s^2, zp = su, sp = xu$ and $u^2 = yz$ by (48), (47), (46), (38), (32), (36) and (42) respectively. Since $xyz = xu^2 = spu = zp^2, xy = p^2$. Thus r is strongly symmetric by Lemma 13.1.2.

If $z = 0$, then $st = 0, t = 0$ and $s = 0$ by (31), (39) and (38) respectively. It follows from (42) and (43) that $u = -v$. By (36), (45) and (29), $xu = 0, u(p + q) = 0$ and $uy = 0$. Consequently, r is the second case.

(IV) Let $\beta = \delta = 0$. We only show that if r is a solution of CYBE, then r is one case in part (IV). If $z \neq 0$, we have that $s = t, s^2 = xz, zp = vs$ and $zq = us$ by (48), (38), (32) and (34) respectively. Consequently, r is the first case in part (IV). If $z = 0$,

then $s = -t$ by (38) and (39). It follows from (30), (31), (28), (37), (36), (45) and (44) that $vs = 0, us = 0, xs = 0, xu = 0, xv = 0, up = qv$ and $s(p + q) = 0$. Consequently, r is the second case in part (IV). $\square$

Corollary 13.1.7. *Let L be a Lie algebra with $\dim L \leq 3$ and $r \in L \otimes L$. If r is strongly symmetric, then r is a solution of CYBE in L .*

Proof . If k is algebraically closed, then r is a solution of CYBE by Proposition 13.1.4, 13.1.6 and 13.1.3. If k is not algebraically closed, let P be algebraic closure of k . We can construct a Lie algebra $L_P = P \otimes L$ over P , as in [59, section 8]. Set $\Psi : L \longrightarrow L_P$ by sending x

to $1 \otimes x$. It is clear that L_P is a Lie algebra over P and Ψ is homomorphic with $\ker \Psi = 0$ over k . Let

$$\bar{r} = (\Psi \otimes \Psi)(r).$$

Obviously, $\bar{r}$ is strongly symmetric. Therefore $\bar{r}$ is a solution of CYBE in L_P and so is r in L . $\square$

13.2 Coboundary Lie bialgebras

In this section, using the general solution, which are obtained in the preceding section, of CYBE in Lie algebra L with $\dim L \leq 3$, we give the sufficient and necessary conditions for $(L, [\], \Delta_r, r)$ to be a coboundary (or triangular) Lie bialgebra.

We now observe the connection between solutions of CYBE and triangular Lie bialgebra structures. It is clear that if $(L, [\], \Delta_r, r)$ is a triangular Lie bialgebra then r is a solution of CYBE. Conversely, if r is a solution of CYBE and r is skew symmetric with $r \in L \otimes L$, then $(L, [\], \Delta_r, r)$ is a triangular Lie bialgebra by [118, Proposition 1].

Theorem 13.2.1. *Let L be a Lie algebra with a basis $\{e_1, e_2, e_3\}$ such that $[e_1, e_2] = e_3, [e_2, e_3] = \alpha e_1, [e_3, e_1] = \beta e_2$, where $\alpha, \beta \in k$ and $\alpha\beta \neq 0$ or $\alpha = \beta = 0$. Let $p, u, s \in k$ and $r = p(e_1 \otimes e_2) - p(e_2 \otimes e_1) + s(e_1 \otimes e_3) - s(e_3 \otimes e_1) + u(e_2 \otimes e_3) - u(e_3 \otimes e_2)$. Then*

(I) $(L, [\], \Delta_r, r)$ is a coboundary Lie bialgebra iff r is skew symmetric;

(II) $(L, [\], \Delta_r, r)$ is a triangular Lie bialgebra iff

$$\beta s^2 + \alpha u^2 + p^2 = 0.$$

Proof. (I) It is sufficient to show that

$$(1 + \xi + \xi^2)(1 \otimes \Delta)\Delta(e_i) = 0$$

for $i = 1, 2, 3$. First, by computation, we have that

$$\begin{aligned} & (1 \otimes \Delta)\Delta(e_1) \\ &= \{(e_1 \otimes e_2 \otimes e_3)(\beta ps - \beta ps) + (e_3 \otimes e_1 \otimes e_2)(\beta ps) + (e_2 \otimes e_3 \otimes e_1)(-\beta sp)\} \\ &+ \{(e_1 \otimes e_3 \otimes e_2)(-\beta ps + \beta ps) + (e_3 \otimes e_2 \otimes e_1)(-\beta ps) + (e_2 \otimes e_1 \otimes e_3)(\beta sp)\} \\ &+ \{(e_1 \otimes e_1 \otimes e_2)(\alpha\beta su) + (e_1 \otimes e_2 \otimes e_1)(-\beta\alpha su) + (e_2 \otimes e_1 \otimes e_1)(0)\} \\ &+ \{(e_1 \otimes e_1 \otimes e_3)(-\alpha pu) + (e_1 \otimes e_3 \otimes e_1)(\alpha pu) + (e_3 \otimes e_1 \otimes e_1)(0)\} \\ &+ \{(e_2 \otimes e_1 \otimes e_2)(-\beta^2 s^2) + (e_1 \otimes e_2 \otimes e_2)(0) + (e_2 \otimes e_2 \otimes e_1)(\beta^2 s^2)\} \\ &+ \{(e_3 \otimes e_3 \otimes e_1)(p^2) + (e_3 \otimes e_1 \otimes e_3)(-p^2) + (e_1 \otimes e_3 \otimes e_3)(0)\}. \end{aligned}$$

Thus

$$(1 + \xi + \xi^2)(1 \otimes \Delta)\Delta(e_1) = 0.$$

Similarly, we have that

$$(1 + \xi + \xi^2)(1 \otimes \Delta)\Delta(e_2) = 0, \quad (1 + \xi + \xi^2)(1 \otimes \Delta)\Delta(e_3) = 0.$$

Thus $(L, [\cdot, \cdot], \Delta_r, r)$ is a coboundary Lie bialgebra.

(II) By part (I), it is sufficient to show that r is a solution of CYBE iff

$$\beta s^2 + \alpha u^2 + p^2 = 0.$$

(a) Let $\alpha \neq 0$ and $\beta \neq 0$. Considering r is skew symmetric, by Proposition 13.1.4 (II), we have that r is a solution of CYBE iff

$$\beta s^2 + \alpha u^2 + p^2 = 0.$$

(b) Let $\alpha = \beta = 0$. Considering r is skew symmetric, by Proposition 13.1.4 (III), we have that r is a solution of CYBE iff

$$\beta s^2 + \alpha u^2 + p^2 = 0.$$

□

Let $(L, [\cdot, \cdot])$ be a familiar Lie algebra, namely, Euclidean 3-space under vector cross product. By Theorem 13.2.1, $(L, [\cdot, \cdot], \Delta_r, r)$ is not a triangular Lie bialgebra for any $0 \neq r \in L \otimes L$. In fact, [84, Example 2.14] already contains the essence of this observation.

Example 13.2.2. Under Example 13.1.5, we have the following:

- (i) $(sl(2), [\cdot, \cdot], \Delta_r, r)$ is a coboundary Lie bialgebra iff r is skew symmetric;
- (ii) $(sl(2), [\cdot, \cdot], \Delta_r, r)$ is a triangular Lie bialgebra iff

$$-4s^2 + 4u^2 + p^2 = 0,$$

where $r = p(e_1 \otimes e_2) - p(e_2 \otimes e_1) + s(e_1 \otimes e_3) - s(e_3 \otimes e_1) + u(e_2 \otimes e_3) - u(e_3 \otimes e_2)$ and $p, s, u \in k$.

Theorem 13.2.3. Let L be a Lie algebra with a basis $\{e_1, e_2, e_3\}$ such that

$$[e_1 e_2] = 0, [e_1, e_3] = e_1 + \beta e_2, [e_2, e_3] = \delta e_2,$$

where $\delta, \beta \in k$. Let $p, s, u \in k$ and $r = p(e_1 \otimes e_2) - p(e_2 \otimes e_1) + s(e_1 \otimes e_3) - s(e_3 \otimes e_1) + u(e_2 \otimes e_3) - u(e_3 \otimes e_2)$.

(I) $(L, [\cdot, \cdot], \Delta_r, r)$ is a coboundary Lie bialgebra iff

$$(\delta + 1)((\delta - 1)u + \beta s)s = 0;$$

(II) If $\beta = 0$, then $(L, [\cdot, \cdot], \Delta_r, r)$ is a triangular Lie bialgebra iff

$$(1 - \delta)us = 0;$$

(III) If $\beta \neq 0$ and $\delta = 1$, then $(L, [\cdot, \cdot], \Delta_r, r)$ is a triangular Lie bialgebra iff

$$s = 0$$

iff $(L, [\cdot, \cdot], \Delta_r, r)$ is a coboundary Lie bialgebra.

Proof . (I) We get by computation

$$\begin{aligned}
& (1 \otimes \Delta)\Delta(e_1) \\
&= \{(e_1 \otimes e_1 \otimes e_2)(\beta\delta ss - \delta us) + (e_1 \otimes e_2 \otimes e_1)(-\beta\delta ss + \delta us) + (e_2 \otimes e_1 \otimes e_1)(0)\} \\
&+ \{(e_2 \otimes e_2 \otimes e_1)(-\beta us + uu + \beta^2 s^2 - \beta su) \\
&+ (e_2 \otimes e_1 \otimes e_2)(\beta us - uu + \beta us - \beta^2 s^2) + (e_1 \otimes e_2 \otimes e_2)(0)\} \\
& (1 \otimes \Delta)\Delta(e_2) \\
&= \{(e_1 \otimes e_1 \otimes e_2)(\delta^2 ss) + (e_1 \otimes e_2 \otimes e_1)(-\delta^2 s^2) + (e_2 \otimes e_1 \otimes e_1)(0)\} \\
&+ \{(e_2 \otimes e_2 \otimes e_1)(\delta\beta ss - \delta su) + (e_2 \otimes e_1 \otimes e_2)(\delta su - \delta\beta ss) + (e_1 \otimes e_2 \otimes e_2)(0)\} \\
& (1 \otimes \Delta)\Delta(e_3) \\
&= \{(e_1 \otimes e_2 \otimes e_3)(\delta su + \beta ss) + (e_2 \otimes e_3 \otimes e_1)(-\delta us - \beta ss) \\
&+ (e_3 \otimes e_1 \otimes e_2)(\delta\delta us + \beta\delta ss + \beta ss - su)\} \\
&+ \{(e_1 \otimes e_3 \otimes e_2)(-\delta su - \beta ss) + (e_3 \otimes e_2 \otimes e_1)(-\beta\delta ss - \delta\delta us - \beta ss + su) \\
&+ (e_2 \otimes e_1 \otimes e_3)(\delta us + \beta ss)\} \\
&+ \{(e_1 \otimes e_1 \otimes e_2)(-\delta sp - \delta\delta ps + sp + \delta sp) \\
&+ (e_1 \otimes e_2 \otimes e_1)(\delta ps + \delta\delta ps - \delta sp - sp) + (e_2 \otimes e_1 \otimes e_1)(0)\} \\
&+ \{(e_1 \otimes e_1 \otimes e_3)(ss) + (e_1 \otimes e_3 \otimes e_1)(-ss) + (e_3 \otimes e_1 \otimes e_1)(0)\} \\
&+ \{(e_2 \otimes e_2 \otimes e_1)(-\beta ps + pu - \delta\beta sp - \delta\delta up - \delta\beta ps + \delta pu - \beta sp - \delta up) \\
&+ (e_2 \otimes e_1 \otimes e_2)(\beta ps - pu + \delta\beta sp + \delta\delta up + \delta\beta ps - \delta pu + \beta sp + \delta up)) \\
&- (e_1 \otimes e_2 \otimes e_2)(0)\} \\
&+ \{(e_2 \otimes e_2 \otimes e_3)(\delta\delta uu + \delta\beta us + \beta\delta us + \beta\beta ss) \\
&+ (e_2 \otimes e_3 \otimes e_2)(-\delta\delta uu - \delta\beta us - \beta\delta us - \beta\beta ss) + (e_3 \otimes e_2 \otimes e_2)(0)\}.
\end{aligned}$$

Consequently,

$$(1 + \xi + \xi^2)(1 \otimes \Delta)\Delta(e_1) = 0,$$

$$(1 + \xi + \xi^2)(1 \otimes \Delta)\Delta(e_2) = 0$$

and

$$(1 + \xi + \xi^2)(1 \otimes \Delta)\Delta(e_3) = 0$$

iff

$$\delta^2 us + \delta\beta s^2 + \beta s^2 - us = 0.$$

This implies that $(L, [\], \Delta_r, r)$ is a coboundary Lie bialgebra iff

$$\delta^2 us + \delta\beta s^2 + \beta s^2 - us = 0 \quad .$$

(II) If $(L, [\], \Delta_r, r)$ is a triangular Lie bialgebra, then $(\delta^2 - 1)us = 0$ by part (I). If $(\delta + 1) \neq 0$, then $(1 - \delta)us = 0$. If $\delta + 1 = 0$, then $(1 - \delta)us = 0$ by Proposition 13.1.6 (II). Conversely, if $(1 - \delta)us = 0$, then we have that $(L, [\], \Delta_r, r)$ is a coboundary Lie bialgebra by part (I). Since

r is skew symmetric, we have that r is a solution of CYBE by Proposition 13.1.6 (II) (IV). Thus $(L, [\cdot, \cdot], \Delta_r, r)$ is a triangular Lie bialgebra.

(III) Similarly, we can show that part (III) holds by part (I) and Proposition 13.1.6 (III). $\square$

Theorem 13.2.4. *If L is a Lie algebra with $\dim L = 2$ and $r \in L \otimes L$, then $(L, [\cdot, \cdot], \Delta_r, r)$ is a triangular Lie bialgebra iff $(L, [\cdot, \cdot], \Delta_r, r)$ is a coboundary Lie bialgebra iff r is skew symmetric.*

Proof. It is an immediate consequence of the main result of [84]. $\square$

13.3 Strongly symmetric elements

In this section, we show that all strongly symmetric elements are solutions of constant classical Yang-Baxter equation in Lie algebra, or of quantum Yang-Baxter equation in algebra. Throughout this section k is an arbitrary field.

Definition 13.3.1. *Let $\{e_i \mid i \in \Omega\}$ be a basis of vector space V and $r = \sum_{i,j \in \Omega} k_{ij}(e_i \otimes e_j) \in V \otimes V$, where $k_{ij} \in k$.*

If

$$k_{ij} = k_{ji} \quad \text{and} \quad k_{ij}k_{lm} = k_{il}k_{jm}$$

for $i, j, l, m \in \Omega$, then r is said to be strongly symmetric to the basis $\{e_i \mid i \in \Omega\}$.

Obviously, symmetry does not depend on the particular choice of basis of V . We need to know if strong symmetry depends on the particular choice of basis of V .

Lemma 13.3.2. *Let $\{e_i \mid i \in \Omega\}$ be a basis of vector space V and $r = \sum_{i,j \in \Omega} k_{ij}(e_i \otimes e_j) \in V \otimes V$, where $k_{ij} \in k$. Then*

(I) the strong symmetry does not depend on the particular choice of basis of V ;

(II) r is strongly symmetric iff

$$k_{ij}k_{lm} = k_{il}k_{jm}$$

for any $i, j, l, m \in \Omega$;

(III) r is strongly symmetric iff

$$k_{ij}k_{lm} = k_{st}k_{uv}$$

where $\{s, t, u, v\} = \{i, j, l, m\}$ as sets for any $i, j, l, m \in \Omega$;

(IV) r is strongly symmetric iff

$$r = 0 \quad \text{or}$$

$$r = \sum_{i,j \in \Omega} k_{i_0 i} k_{i_0 j} k_{i_0 i_0}^{-1} (e_i \otimes e_j) \in V \otimes V, \quad \text{where } k_{i_0 i_0} \neq 0.$$

Proof. (I) Let $\{e'_i \mid i \in \Omega'\}$ be another basis of V . It is sufficient to show that r is strongly symmetric to the basis $\{e'_i \mid i \in \Omega'\}$. Obviously, there exists $q_{ij} \in k$ such that $e_i = \sum_s e'_s q_{si}$ for $i \in \Omega$. By computation, we have that

$$r = \sum_{s,t \in \Omega} \left(\sum_{i,j \in \Omega'} k_{ij} q_{si} q_{tj} \right) (e'_s \otimes e'_t).$$

If set $k'_{st} = \sum_{i,j} k_{ij} q_{si} q_{tj}$, then for $l, m, u, v \in \Omega'$ we have that

$$\begin{aligned} k'_{lm} k'_{uv} &= \sum_{i,j,s,t} k_{ij} k_{st} q_{li} q_{mj} q_{us} q_{vt} \\ &= \sum_{i,j,s,t} k_{is} k_{jt} q_{li} q_{mj} q_{us} q_{vt} \quad (\text{by } k_{ij} k_{st} = k_{is} k_{jt}) \\ &= k'_{lu} k'_{mv} \end{aligned}$$

and

$$k'_{lm} = k'_{ml},$$

which implies that r is strongly symmetric to the basis $\{e'_i \mid i \in \Omega'\}$.

(II) The necessity is obvious. Now we show the sufficiency. For any $i, j \in \Omega$, if $k_{ii} \neq 0$, then $k_{ij} = k_{ji}$ since $k_{ij} k_{ii} = k_{ii} k_{ji}$. If $k_{ii} = 0$, then $k_{ij} = k_{ji} = 0$ since $k_{ij}^2 = k_{ii} k_{jj} = 0 = k_{ji}^2$.

(III) The sufficiency follows from part (II). Now we show the necessity. For any $i, j, l, m \in \Omega$, we see that $k_{ij} k_{lm} = k_{ij} k_{ml} = k_{il} k_{jm} = k_{il} k_{mj} = k_{im} k_{lj}$.

Similarly, we can get that $k_{ij} k_{lm} = k_{st} k_{uv}$ when $s = j$ or $s = l$ or $s = m$.

(IV) If r is a case in part (IV), then r is strongly symmetric by straightforward verification. Conversely, if r is strongly symmetric and $r \neq 0$, then there exists $i_0 \in \Omega$ such that $k_{i_0 i_0} \neq 0$. Thus $k_{ij} = k_{i_0 i} k_{i_0 j} k_{i_0 i_0}^{-1}$. $\square$

If A is an algebra, then we can get a Lie algebra $L(A)$ by defining $[x, y] = xy - yx$ for any $x, y \in A$.

Theorem 13.3.3. *Let A be an algebra and $r \in A \otimes A$ be strongly symmetric. Then r is a solution of CYBE in $L(A)$; meantime, it is also a solution of QYBE in A .*

Proof. Let $\{e_i \mid i \in \Omega\}$ be a basis of A and the multiplication of A be defined as follows:

$$e_i e_j = \sum_m a_{ij}^m e_m$$

for any $i, j \in \Omega$, $a_{ij}^m \in k$.

We first show that r is a solution of CYBE in Lie algebra $L(A)$. By computation, for all $i, j, l \in \Omega$, we have that the coefficient of $e_i \otimes e_j \otimes e_l$ in $[r^{12}, r^{13}]$ is

$$\begin{aligned} & \sum_{st} a_{s,t}^i k_{sj} k_{tl} - a_{ts}^i k_{sj} k_{tl} \\ &= \sum_{s,t} a_{st}^i k_{sj} k_{tl} - \sum_{s,t} a_{st}^i k_{tj} k_{sl} \\ &= 0 \quad \text{by Lemma 13.3.2 (III).} \end{aligned}$$

Similarly, we have that the coefficients of $e_i \otimes e_j \otimes e_l$ in $[r^{12}, r^{23}]$ and $[r^{13}, r^{23}]$ are zero. Thus r is a solution of CYBE in $L(A)$.

Next we show that r is a solution of QYBE in algebra A . We denote r by R since the solution of QYBE is usually denoted by R .

By computation, for all $i, j, l \in \Omega$, we have that the coefficient of $e_i \otimes e_j \otimes e_l$ in $R^{12}R^{13}R^{23}$ is

$$\sum_{s,t,u,v,w,m} a_{st}^i a_{um}^j a_{vw}^l k_{su} k_{tv} k_{mw}$$

and the coefficient of $e_i \otimes e_j \otimes e_l$ in $R^{23}R^{13}R^{12}$ is

$$\begin{aligned} & \sum_{s,t,u,v,w,m} a_{tm}^i a_{sw}^j a_{uv}^l k_{su} k_{tv} k_{mw} \\ &= \sum_{s,t,u,v,w,m} a_{st}^i a_{um}^j a_{vw}^l k_{su} k_{tv} k_{mw} \quad \text{by Lemma 13.3.2 (III).} \end{aligned}$$

Thus r is a solution of QYBE in algebra A . $\square$

Corollary 13.3.4. *Let L be a Lie algebra and $r \in L \otimes L$ be strongly symmetric. Then r is a solution of CYBE.*

Proof: Let A denote the universal enveloping algebra of L . It is clear that r is also strongly symmetric in $A \otimes A$. Thus

$$[r^{12}, r^{13}] + [r^{12}, r^{23}] + [r^{13}, r^{23}] = 0$$

by Theorem 13.3.3. Consequently, r is a solution of CYBE in L . $\square$

13.4 The solutions of CYBE in the case of $\text{char } k = 2$

In this section, we find the general solution of CYBE for Lie algebra L over field k of characteristic 2 with $\dim L \leq 3$. Throughout this section, the characteristic of k is 2.

Proposition 13.4.1. *Let L be a Lie algebra with $\dim L = 2$. Then r is a solution of CYBE iff r is symmetric.*

Proof. The proof is not hard since the tensor r has only 4 coefficients. $\square$

If $r = x(e_1 \otimes e_1) + y(e_2 \otimes e_2) + z(e_3 \otimes e_3) + p(e_1 \otimes e_2) + p(e_2 \otimes e_1) + s(e_1 \otimes e_3) + s(e_3 \otimes e_1) + u(e_2 \otimes e_3) + u(e_3 \otimes e_2)$ with $\alpha yz + \beta xz + xy + \beta s^2 + \alpha u^2 + p^2 = 0$, then r is called α, β -symmetric to the basis $\{e_1, e_2, e_3\}$.

Proposition 13.4.2. *Let L be a Lie algebra with a basis $\{e_1, e_2, e_3\}$ such that $[e_1, e_2] = e_3$, $[e_2, e_3] = \alpha e_1$, $[e_3, e_1] = \beta e_2$, where $\alpha, \beta \in k$. Let $p, q, s, t, u, v, x, y, z \in k$.*

Then (I) if r is α, β -symmetric to basis $\{e_1, e_2, e_3\}$, then r is a solution of CYBE in L ;

(II) if $\alpha\beta \neq 0$, then r is a solution of the CYBE in L iff r is α, β -symmetric to the basis $\{e_1, e_2, e_3\}$;

(III) if $\alpha = \beta = 0$, then r is a solution of the CYBE in L iff

$$r = p(e_1 \otimes e_2) + p(e_2 \otimes e_1) + s(e_1 \otimes e_3) + t(e_3 \otimes e_1) + u(e_2 \otimes e_3) + v(e_3 \otimes e_2) + x(e_1 \otimes e_1) + y(e_2 \otimes e_2) + z(e_3 \otimes e_3)$$

$$\text{with } tu = vs, \quad xy = p^2, \quad (s+t)p + (u+v)x = 0 \quad (s+t)y + (u+v)p = 0.$$

Proof . Let $r = \sum_{i,j=1}^3 k_{ij}(e_i \otimes e_j) \in L \otimes L$ and $k_{ij} \in k$, with $i, j = 1, 2, 3$. It is clear that

$$\begin{aligned} [r^{12}, r^{13}] &= \sum_{i,j=1}^3 \sum_{s,t=1}^3 k_{ij} k_{st} [e_i, e_s] \otimes e_j \otimes e_t, \\ [r^{12}, r^{23}] &= \sum_{i,j=1}^3 \sum_{s,t=1}^3 k_{ij} k_{st} e_i \otimes [e_j, e_s] \otimes e_t, \\ [r^{13}, r^{23}] &= \sum_{i,j=1}^3 \sum_{s,t=1}^3 k_{ij} k_{st} e_i \otimes e_s \otimes [e_j, e_t]. \end{aligned}$$

By computation, for all $i, j, n = 1, 2, 3$, we have that the coefficient of $e_j \otimes e_i \otimes e_i$ in $[r^{12}, r^{13}]$ is zero and $e_i \otimes e_i \otimes e_j$ in $[r^{13}, r^{23}]$ is zero.

We now see the coefficient of $e_i \otimes e_j \otimes e_n$ in $[r^{12}, r^{13}] + [r^{12}, r^{23}] + [r^{13}, r^{23}]$.

- (1) $e_1 \otimes e_1 \otimes e_1(\alpha k_{12} k_{31} - \alpha k_{13} k_{21});$
- (2) $e_2 \otimes e_2 \otimes e_2(-\beta k_{21} k_{32} + \beta k_{23} k_{12});$
- (3) $e_3 \otimes e_3 \otimes e_3(k_{31} k_{23} - k_{32} k_{13});$
- (4) $e_1 \otimes e_2 \otimes e_3(\alpha k_{22} k_{33} - \alpha k_{32} k_{23} - \beta k_{11} k_{33} + \beta k_{13} k_{13} + k_{11} k_{22} - k_{12} k_{21});$
- (5) $e_2 \otimes e_3 \otimes e_1(\beta k_{33} k_{11} - \beta k_{13} k_{31} - k_{22} k_{11} + k_{21} k_{21} + \alpha k_{22} k_{33} - \alpha k_{23} k_{32});$
- (6) $e_3 \otimes e_1 \otimes e_2(k_{11} k_{22} - k_{21} k_{12} - \alpha k_{33} k_{22} + \alpha k_{32} k_{32} + \beta k_{33} k_{11} - \beta k_{31} k_{13});$
- (7) $e_1 \otimes e_3 \otimes e_2(-\alpha k_{33} k_{22} + \alpha k_{23} k_{32} + k_{11} k_{22} - k_{12} k_{12} - \beta k_{11} k_{33} + \beta k_{13} k_{31});$
- (8) $e_3 \otimes e_2 \otimes e_1(-k_{22} k_{11} + k_{12} k_{21} + \beta k_{33} k_{11} - \beta k_{31} k_{31} - \alpha k_{33} k_{22} + \alpha k_{32} k_{23});$
- (9) $e_2 \otimes e_1 \otimes e_3(-\beta k_{11} k_{33} + \beta k_{31} k_{13} + \alpha k_{22} k_{33} - \alpha k_{23} k_{23} - k_{22} k_{11} + k_{21} k_{12});$
- (10) $e_1 \otimes e_1 \otimes e_2(-\alpha k_{31} k_{22} + \alpha k_{21} k_{32} + \alpha k_{12} k_{32} - \alpha k_{13} k_{22});$
- (11) $e_2 \otimes e_1 \otimes e_1(-\alpha k_{23} k_{21} + \alpha k_{22} k_{31} + \alpha k_{22} k_{13} - \alpha k_{23} k_{12});$
- (12) $e_1 \otimes e_1 \otimes e_3(\alpha k_{21} k_{33} - \alpha k_{31} k_{23} + \alpha k_{12} k_{33} - \alpha k_{13} k_{23});$
- (13) $e_3 \otimes e_1 \otimes e_1(\alpha k_{32} k_{31} - \alpha k_{33} k_{21} + \alpha k_{32} k_{13} - \alpha k_{33} k_{12});$
- (14) $e_2 \otimes e_2 \otimes e_1(-\beta k_{12} k_{31} + \beta k_{32} k_{11} + \beta k_{23} k_{11} - \beta k_{21} k_{31});$
- (15) $e_1 \otimes e_2 \otimes e_2(-\beta k_{11} k_{32} + \beta k_{13} k_{12} - \beta k_{11} k_{23} + \beta k_{13} k_{21});$
- (16) $e_2 \otimes e_2 \otimes e_3(\beta k_{32} k_{13} - \beta k_{12} k_{33} - \beta k_{21} k_{33} + \beta k_{23} k_{13});$
- (17) $e_3 \otimes e_2 \otimes e_2(-\beta k_{31} k_{32} + \beta k_{33} k_{12} + \beta k_{33} k_{21} - \beta k_{31} k_{23});$
- (18) $e_3 \otimes e_3 \otimes e_1(k_{13} k_{21} - k_{23} k_{11} + k_{31} k_{21} - k_{32} k_{11});$
- (19) $e_3 \otimes e_3 \otimes e_2(-k_{23} k_{12} + k_{13} k_{22} - k_{32} k_{12} + k_{31} k_{22});$
- (20) $e_2 \otimes e_3 \otimes e_3(k_{21} k_{23} - k_{22} k_{13} + k_{21} k_{32} - k_{22} k_{31});$
- (21) $e_1 \otimes e_3 \otimes e_3(-k_{12} k_{13} + k_{11} k_{23} - k_{12} k_{31} + k_{11} k_{32});$
- (22) $e_1 \otimes e_3 \otimes e_1(\alpha k_{23} k_{31} - \alpha k_{33} k_{21} - k_{12} k_{11} + k_{11} k_{21} + \alpha k_{12} k_{33} - \alpha k_{13} k_{32});$
- (23) $e_1 \otimes e_2 \otimes e_1(-\alpha k_{32} k_{21} + \alpha k_{22} k_{31} - \beta k_{11} k_{31} + \beta k_{13} k_{11} - \alpha k_{13} k_{22} + \alpha k_{12} k_{23});$
- (24) $e_2 \otimes e_1 \otimes e_2(\beta k_{31} k_{12} - \beta k_{11} k_{32} + \alpha k_{22} k_{32} - \alpha k_{23} k_{22} - \beta k_{21} k_{13} + \beta k_{23} k_{11});$
- (25) $e_2 \otimes e_3 \otimes e_2(-\beta k_{13} k_{32} + \beta k_{33} k_{12} + k_{21} k_{22} - k_{22} k_{12} - \beta k_{21} k_{33} + \beta k_{23} k_{31});$
- (26) $e_3 \otimes e_2 \otimes e_3(k_{12} k_{23} - k_{22} k_{13} - \beta k_{31} k_{33} + \beta k_{33} k_{13} + k_{31} k_{22} - k_{32} k_{21});$
- (27) $e_3 \otimes e_1 \otimes e_3(-k_{21} k_{13} + k_{11} k_{23} + \alpha k_{32} k_{33} - \alpha k_{33} k_{23} + k_{31} k_{12} - k_{32} k_{11}).$

Let $k_{11} = x, k_{22} = y, k_{33} = z, k_{12} = p, k_{21} = q, k_{13} = s, k_{31} = t, k_{23} = u, k_{32} = v$.

It follows from (1)–(27) that

- (28) $\alpha pt = \alpha qs$;
- (29) $\beta qv = \beta pu$;
- (30) $tu = vs$;
- (31) $\alpha yz - \beta xz + xy - \alpha uv + \beta s^2 - pq = 0$;
- (32) $\beta zx - yx + \alpha yz - \beta st + q^2 - \alpha uv = 0$;
- (33) $xy - \alpha zy + \beta zx - pq + \alpha v^2 - \beta st = 0$;
- (34) $-\alpha zy + xy - \beta xz + \alpha uv - p^2 + \beta st = 0$;
- (35) $-xy + \beta xz - \alpha yz + pq - \beta t^2 + \alpha uv = 0$;
- (36) $-\beta xz + \alpha yz - yx + \beta st - \alpha u^2 + pq = 0$;
- (37) $\alpha(-ty + qv + pv - sy) = 0$;
- (38) $\alpha(-uq + yt + ys - up) = 0$;
- (39) $\alpha(qz - tu + pz - su) = 0$;
- (40) $\alpha(vt - zq + vs - zp) = 0$;
- (41) $\beta(-pt + vx + ux - qt) = 0$;
- (42) $\beta(-xv + sp - xu + sq) = 0$;
- (43) $\beta(vs - pz + us - qz) = 0$;
- (44) $\beta(-tv + zp + zq - tu) = 0$;
- (45) $sq - ux + tq - vx = 0$;
- (46) $-up + sy - vp + ty = 0$;
- (47) $qu - ys + qv - yt = 0$;
- (48) $-ps + xu - pt + xv = 0$;
- (49) $\alpha ut - \alpha zq - px + xq + \alpha pz - \alpha sv = 0$;
- (50) $-\alpha vq + \alpha yt - \beta xt + \beta sx - \alpha sy + \alpha pu = 0$;
- (51) $\beta tp - \beta xv + \alpha yv - \alpha uy - \beta qs + \beta ux = 0$;
- (52) $-\beta sv + \beta zp + qy - yp - \beta qz + \beta ut = 0$;
- (53) $pu - ys - \beta tz + \beta zs + ty - vq = 0$;
- (54) $-qs + xu + \alpha vz - \alpha zu + tp - vx = 0$.

It is clear that r is the solution of CYBE iff relations (28)–(54) hold.

- (55) $q^2 = p^2$ (by (34) + (32));
- (56) $\alpha u^2 = \alpha v^2$ (by (33) + (36));
- (57) $\beta t^2 = \beta s^2$ (by (31) + (35));

By computation, we have that if r is α, β -symmetric then relations (28)–(54) hold and so r is a solution of CYBE. That is, part (I) holds.

(II) By part (I), we have that the sufficiency of part (II) holds. For the necessity, if r is a solution of CYBE, then $p = q, s = t$ and $u = v$ by (55)–(57), since $\text{char } k = 2$. By relation (31), we have that

$$\alpha yz + \beta xz + xy + \beta s^2 + \alpha u^2 + p^2 = 0.$$

Thus r is α, β -symmetric.

(III) It is trivial. $\square$

In particular, Proposition 13.4.2 implies:

Example 13.4.3. *Let*

$$sl(2) := \{x \mid x \text{ is a } 2 \times 2 \text{ matrix with trace zero over } k\}$$

and

$$e_1 = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, e_2 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, e_3 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Thus L is a Lie algebra (defined by $[x, y] = xy - yx$) with a basis $\{e_1, e_2, e_3\}$. It is clear that

$$[e_1, e_2] = e_3, [e_2, e_3] = 0e_1, [e_3, e_1] = 0e_2.$$

Consequently, r is a solution of CYBE iff

$$r = p(e_1 \otimes e_2) + p(e_2 \otimes e_1) + s(e_1 \otimes e_3) + t(e_3 \otimes e_1) + u(e_2 \otimes e_3) + v(e_3 \otimes e_2) + x(e_1 \otimes e_1) + y(e_2 \otimes e_2) + z(e_3 \otimes e_3)$$

$$\text{with } tu = vs, \quad xy = p^2, \quad (s+t)p + (u+v)x = 0, \quad (s+t)y + (u+v)p = 0.$$

By the way, we have that α, β -symmetry does not depend on the particular choice of basis of L when $\alpha\beta \neq 0$.

Proposition 13.4.4. *Let L be a Lie algebra with a basis $\{e_1, e_2, e_3\}$ such that $[e_1, e_2] = 0, [e_1, e_3] = e_1 + \beta e_2, [e_2, e_3] = \delta e_2$, where $\beta, \delta \in k$. Let $p, q, s, t, u, v, x, y, z \in k$.*

(I) *If $\beta = 0, \delta \neq 0$, then r is a solution of CYBE in L iff*

$$r = p(e_1 \otimes e_2) + q(e_2 \otimes e_1) + s(e_1 \otimes e_3) + s(e_3 \otimes e_1) + u(e_2 \otimes e_3) + u(e_3 \otimes e_2) + x(e_1 \otimes e_1) + y(e_2 \otimes e_2) + z(e_3 \otimes e_3)$$

$$\text{where } (\delta + 1)zp = (\delta + 1)qz = (\delta + 1)us, (\delta + 1)uq = (\delta + 1)up \text{ and } (\delta + 1)ps = (\delta + 1)qs.$$

(II) *If $\beta \neq 0, \delta = 1$, then r is a solution of CYBE in L iff*

$$r = p(e_1 \otimes e_2) + q(e_2 \otimes e_1) + s(e_1 \otimes e_3) + s(e_3 \otimes e_1) + u(e_2 \otimes e_3) + u(e_3 \otimes e_2) + x(e_1 \otimes e_1) + y(e_2 \otimes e_2) + z(e_3 \otimes e_3)$$

$$\text{with } sp = sq, up = qu, zq = zp \text{ and } s^2 = xz.$$

(III) *If $\beta = \delta = 0$, then r is a solution of CYBE in L iff*

$$r = p(e_1 \otimes e_2) + q(e_2 \otimes e_1) + s(e_1 \otimes e_3) + s(e_3 \otimes e_1) + u(e_2 \otimes e_3) + v(e_3 \otimes e_2) + x(e_1 \otimes e_1) + y(e_2 \otimes e_2) + z(e_3 \otimes e_3)$$

$$\text{with } vs = pz, us = qz, qv = pu \text{ and } (u + v)x = (p + q)s.$$

Proof. Let $r = \sum_{i,j=1}^3 k_{ij}(e_i \otimes e_j) \in L \otimes L$ and $k_{ij} \in k$, with $i, j = 1, 2, 3$. By computation, for all $i, j, n = 1, 2, 3$, we have that the coefficient of $e_j \otimes e_i \otimes e_i$ in $[r^{12}, r^{13}]$ is zero and $e_i \otimes e_i \otimes e_j$ in $[r^{13}, r^{23}]$ is zero.

We now see the coefficient of $e_i \otimes e_j \otimes e_n$

$$\text{in } [r^{12}, r^{13}] + [r^{12}, r^{23}] + [r^{13}, r^{23}].$$

$$(1) \quad e_1 \otimes e_1 \otimes e_1(-k_{13}k_{11} + k_{11}k_{31});$$

$$(2) \quad e_2 \otimes e_2 \otimes e_2(-\beta k_{23}k_{12} + \beta k_{21}k_{32} - \delta k_{23}k_{22} + \delta k_{22}k_{32});$$

$$(3) \quad e_3 \otimes e_3 \otimes e_3(0);$$

$$(4) \quad e_1 \otimes e_2 \otimes e_3(-k_{32}k_{13} + k_{12}k_{33} - \beta k_{13}k_{13} + \beta k_{11}k_{33} - \delta k_{13}k_{23} + \delta k_{12}k_{33});$$

$$(5) \quad e_2 \otimes e_3 \otimes e_1(-\beta k_{33}k_{11} + \beta k_{13}k_{31} - \delta k_{33}k_{21} + \delta k_{23}k_{31} - k_{23}k_{31} + k_{21}k_{33});$$

- (6) $e_3 \otimes e_1 \otimes e_2(-k_{33}k_{12} + k_{31}k_{32} - \beta k_{33}k_{11} + \beta k_{31}k_{13} - \delta k_{33}k_{12} + \delta k_{32}k_{13});$
- (7) $e_1 \otimes e_3 \otimes e_2(-k_{33}k_{12} + k_{13}k_{32} - \beta k_{13}k_{32} + \beta k_{11}k_{33} - \delta k_{13}k_{32} + \delta k_{12}k_{33});$
- (8) $e_3 \otimes e_2 \otimes e_1(-\beta k_{33}k_{11} + \beta k_{31}k_{31} - \delta k_{33}k_{21} + \delta k_{32}k_{31} - k_{33}k_{21} + k_{31}k_{23});$
- (9) $e_2 \otimes e_1 \otimes e_3(-\beta k_{31}k_{13} + \beta k_{11}k_{33} - \delta k_{31}k_{23} + \delta k_{21}k_{33} - k_{23}k_{13} + k_{21}k_{33});$
- (10) $e_1 \otimes e_1 \otimes e_2(-k_{31}k_{12} + k_{11}k_{32} - k_{13}k_{12} + k_{11}k_{32});$
- (11) $e_2 \otimes e_1 \otimes e_1(-k_{23}k_{11} + k_{21}k_{31} - k_{23}k_{11} + k_{21}k_{13});$
- (12) $e_1 \otimes e_1 \otimes e_3(-k_{31}k_{13} + k_{11}k_{33} - k_{13}k_{13} + k_{11}k_{33});$
- (13) $e_3 \otimes e_1 \otimes e_1(-k_{33}k_{11} + k_{31}k_{31} - k_{33}k_{11} + k_{31}k_{13});$
- (14) $e_2 \otimes e_2 \otimes e_1(-\beta k_{32}k_{11} + \beta k_{12}k_{31} - \delta k_{32}k_{21} + \delta k_{22}k_{31} - \beta k_{23}k_{11} + \beta k_{21}k_{31} - \delta k_{23}k_{21} + \delta k_{22}k_{31});$
- (15) $e_1 \otimes e_2 \otimes e_2(-\beta k_{13}k_{12} + \beta k_{11}k_{32} - \delta k_{13}k_{22} + \delta k_{12}k_{32} - \beta k_{13}k_{21} + \beta k_{11}k_{23} - \delta k_{13}k_{22} + \delta k_{12}k_{23});$
- (16) $e_2 \otimes e_2 \otimes e_3(-\beta k_{32}k_{13} + \beta k_{12}k_{33} - \delta k_{32}k_{23} + \delta k_{22}k_{33} - \beta k_{23}k_{13} + \beta k_{21}k_{33} - \delta k_{23}k_{23} + \delta k_{22}k_{33});$
- (17) $e_3 \otimes e_2 \otimes e_2(-\beta k_{33}k_{12} + \beta k_{31}k_{32} - \delta k_{33}k_{22} + \delta k_{32}k_{32} - \beta k_{33}k_{21} + \beta k_{31}k_{23} - \delta k_{33}k_{22} + \delta k_{32}k_{23});$
- (18) $e_3 \otimes e_3 \otimes e_1(0);$
- (19) $e_1 \otimes e_3 \otimes e_3(0);$
- (20) $e_3 \otimes e_3 \otimes e_2(0);$
- (21) $e_2 \otimes e_3 \otimes e_3(0);$
- (22) $e_1 \otimes e_3 \otimes e_1(-k_{33}k_{11} + k_{13}k_{31} - k_{13}k_{31} + k_{11}k_{33}) = e_1 \otimes e_3 \otimes e_1(0);$
- (23) $e_1 \otimes e_2 \otimes e_1(-k_{32}k_{11} + k_{12}k_{31} - \beta k_{13}k_{11} + \beta k_{11}k_{31} - \delta k_{13}k_{21} + \delta k_{12}k_{31} - k_{13}k_{21} + k_{11}k_{23});$
- (24) $e_2 \otimes e_1 \otimes e_2(-\beta k_{31}k_{12} + \beta k_{11}k_{32} - \delta k_{31}k_{22} + \delta k_{21}k_{32} - k_{23}k_{12} + k_{21}k_{32} - \beta k_{23}k_{11} + \beta k_{21}k_{13} - \delta k_{23}k_{12} + \delta k_{22}k_{13});$

- (25) $e_2 \otimes e_3 \otimes e_2(-\beta k_{33}k_{12} + \beta k_{13}k_{32} - \delta k_{33}k_{22} + \delta k_{23}k_{32} - \beta k_{23}k_{31} + \beta k_{21}k_{33} + \delta k_{33}k_{22} - \delta k_{22}k_{33});$
- (26) $e_3 \otimes e_2 \otimes e_3(-\beta k_{33}k_{13} + \beta k_{31}k_{33} - \delta k_{33}k_{23} + \delta k_{32}k_{33});$
- (27) $e_3 \otimes e_1 \otimes e_3(-k_{33}k_{13} + k_{31}k_{33});$

Let $k_{11} = x, k_{22} = y, k_{33} = z, k_{12} = p, k_{21} = q, k_{13} = s, k_{31} = t, k_{23} = u, k_{32} = v$.

It follows from (1)–(27) that

- (28) $-sx + xt = 0;$
- (29) $-\beta up + \beta qv - \delta uy + \delta yv = 0;$
- (30) $-vs + pz - \beta s^2 + \beta xz - \delta su + \delta zp = 0;$
- (31) $-\beta xz + \beta st - \delta zq + \delta ut - ut + qz = 0;$
- (32) $-zp + tv - \beta zx + \beta ts - \delta zp + \delta vs = 0;$
- (33) $-zp + sv - \beta st + \beta xz - \delta sv + \delta zp = 0;$
- (34) $-\beta zx + \beta tt - \delta zq + \delta vt - zq + tu = 0;$
- (35) $-\beta st + \beta xz - \delta tu + \delta qz - us + qz = 0;$
- (36) $-tp + xv - sp + vx = 0;$
- (37) $-ux + qt - ux + qs = 0;$
- (38) $-st + xz - s^2 + xz = 0;$
- (39) $-zx + tt - zx + st = 0;$
- (40) $-\beta vx + \beta pt - \delta vq + \delta yt - \beta ux + \beta qt - \delta uq + \delta yt = 0;$
- (41) $-\beta sp + \beta xv - \delta sy + \delta pv - \beta sq + \beta xu - \delta sy + \delta pu = 0;$
- (42) $-\beta vs + \beta pz - \delta vu + \delta yz - \beta su + \beta zq - \delta u^2 + \delta yz = 0;$
- (43) $-\beta zp + \beta tv - \delta zy + \delta vv - \beta zq + \beta tu - \delta zy + \delta vu = 0;$

$$(44) -vx + pt - \beta sx + \beta xt - \delta sq + \delta pt - sq + xu = 0;$$

$$(45) -\beta pt + \beta vx - \delta ty + \delta qv - up + qv - \beta ux + \beta qs - \delta up + \delta ys = 0;$$

$$(46) -\beta zp + \beta sv - \beta ut + \beta qz = 0;$$

$$(47) -\beta zs + \beta tz - \delta zu + \delta vz = 0;$$

$$(48) -zs + tz = 0;$$

It is clear that r is a solution of CYBE iff (28)–(48) hold.

By computation we have that

$$(49) t = s \text{ (by (39) + (38))};$$

$$(50) \delta u = \delta v \text{ (by (42) + (43))}.$$

(I) Let $\beta = 0$ and $\delta \neq 0$. If r is the case in part (II), then r is a solution of CYBE by straightforward verification. Conversely, if r is a solution of CYBE, then $u = v$ by (50) and we have that

$$(1 + \delta)zp = (1 + \delta)us \quad \text{by (30)}$$

$$(1 + \delta)zq = (1 + \delta)us \quad \text{by (31)}$$

$$(1 + \delta)sp = (1 + \delta)qs \quad \text{by (44)}$$

$$(1 + \delta)qu = (1 + \delta)up \quad \text{by (45)}$$

Thus r is the case in part (II).

(II) Let $\beta \neq 0$ and $\delta = 1$. If r is the case in part (III), then r is a solution of CYBE by straightforward verification. Conversely, if r is a solution of CYBE then $u = v$ by (50) and we have that

$$up = qu \quad \text{by (29)}$$

$$s^2 = xz \quad \text{by (30)}$$

$$sp = qs \quad \text{by (40)}$$

$$zp = zq \quad \text{by (46)}$$

Thus r is the case in part (III).

(III) Let $\beta = \delta = 0$. It is clear that the system of equations (28)–(48) is equivalent to the following:

$$\left\{ \begin{array}{l} s = t \\ us = qz \\ vs = zp \\ up = qv \\ (p + q)s = (u + v)x \end{array} \right.$$

Thus we complete the proof. $\square$

13.5 Coboundary Lie bialgebras in the case of $\text{char } k = 2$

In this section, using the general solution, which are obtained in the preceding section, of CYBE in Lie algebra L with $\dim L \leq 3$, we give the sufficient and necessary conditions for

$(L, [\cdot, \cdot], \Delta_r, r)$ to be a coboundary (or triangular) Lie bialgebra over field k of characteristic 2. Throughout this section, the characteristic of field k is 2.

We now observe the connection between solutions of CYBE and triangular Lie bialgebra structures.

Lemma 13.5.1. *If $r \in \text{Im}(1 - \tau)$, then*

$$x \cdot C(r) = (1 + \xi + \xi^2)(1 \otimes \Delta)\Delta(x)$$

for any $x \in L$.

Proof: Let $r = \sum_i (a_i \otimes b_i - b_i \otimes a_i)$.

We see that

$$\begin{aligned} C(r) &= \sum_{i,k} ([a_k, a_i] \otimes b_k \otimes b_i - [a_k, b_i] \otimes b_k \otimes a_i - [b_k, a_i] \otimes a_k \otimes b_i \\ &\quad + [b_k, b_i] \otimes a_k \otimes a_i + a_k \otimes [b_k, a_i] \otimes b_i - a_k \otimes [b_k, b_i] \otimes a_i \\ &\quad - b_k \otimes [a_k, a_i] \otimes b_i - b_k \otimes [a_k, a_i] \otimes b_i + b_k \otimes [a_k, b_i] \otimes a_i \\ &\quad + a_k \otimes a_i \otimes [b_k, b_i] - a_k \otimes b_i \otimes [b_k, a_i] - b_k \otimes a_i \otimes [a_k, b_i] \\ &\quad + b_k \otimes b_i \otimes [a_k, a_i]) \end{aligned}$$

and

$$\begin{aligned} &(1 \otimes \Delta)\Delta(x) \\ &= \sum_{i,k} ([x, a_k] \otimes [b_k, a_i] \otimes b_i + [x, a_k] \otimes a_i \otimes [b_k, b_i] - [x, a_k] \otimes [b_k, b_i] \otimes a_i \\ &\quad - [x, a_k] \otimes b_i \otimes [b_k, a_i] - [x, b_k] \otimes [a_k, a_i] \otimes b_i - [x, b_k] \otimes a_i \otimes [a_k, b_i] \\ &\quad + [x, b_k] \otimes [a_k, b_i] \otimes a_i + [x, b_k] \otimes b_i \otimes [a_k, a_i] + a_k \otimes [[x, b_k], a_i] \otimes b_i \\ &\quad + a_k \otimes a_i \otimes [[x, b_k], b_i] - a_k \otimes [[x, b_k], b_i] \otimes a_i - a_k \otimes b_i \otimes [[x, b_k], a_i] \\ &\quad - b_k \otimes [[x, a_k], a_i] \otimes b_i - b_k \otimes a_i \otimes [[x, a_k], b_i] + b_k \otimes [[x, a_k], b_i] \otimes a_i \\ &\quad + b_k \otimes b_i \otimes [[x, a_k], a_i]). \end{aligned}$$

It is easy to check that the sum of the terms whose third factor in $(1 + \xi + \xi^2)(1 \otimes \Delta)\Delta(x)$ includes element x is equal to

$$\begin{aligned}
& \sum_{i,k} \{ [b_k, a_i] \otimes b_i \otimes [x, a_k] + a_i \otimes [b_k, b_i] \otimes [x, a_k] - [b_k, b_i] \otimes a_i \otimes [x, a_k] \\
& - b_i \otimes [b_k, a_i] \otimes [x, a_k] - [a_k, a_i] \otimes b_i \otimes [x, b_k] - a_i \otimes [a_k, b_i] \otimes [x, b_k] \\
& + [a_k, b_i] \otimes a_i \otimes [x, b_k] + b_i \otimes [a_k, a_i] \otimes [x, b_k] \\
& + (b_i \otimes a_k \otimes [[x, b_k], a_i] - b_k \otimes a_i \otimes [[x, a_k], b_i]) \\
& + (a_k \otimes a_i \otimes [[x, b_k], b_i] - a_i \otimes a_k \otimes [[x, b_k], b_i]) \\
& + (-a_k \otimes b_i \otimes [[x, b_k], a_i] + a_i \otimes b_k \otimes [[x, a_k], b_i]) \\
& + (-b_i \otimes b_k \otimes [[x, a_k], a_i] + b_k \otimes b_i \otimes [[x, a_k], a_i]) \} \\
& = (1 \otimes 1 \otimes L_x)C(r) \quad \text{by Jacobi identity,}
\end{aligned}$$

where L_x denotes the adjoint action $L_x(y) = [x, y]$ of L on L .

Consequently,

$$(1 + \xi + \xi^2)(1 \otimes \Delta)\Delta(x) = (L_x \otimes L_x \otimes L_x)C(r) = x \cdot C(r)$$

for any $x \in L$. $\square$

By Lemma 13.5.1 and [84, Proposition 2.11], we have

Theorem 13.5.2. *$(L, [\cdot, \cdot], \Delta_r, r)$ is a triangular Lie bialgebra iff r is a solution of CYBE in L and $r \in \text{Im}(1 - \tau)$.*

Consequently, we can easily get a triangular Lie bialgebra structure by means of a solution of CYBE.

Theorem 13.5.3. *Let L be a Lie algebra with a basis $\{e_1, e_2, e_3\}$ such that $[e_1, e_2] = e_3, [e_2, e_3] = \alpha e_1, [e_3, e_1] = \beta e_2$, where $\alpha, \beta \in k$, $\alpha\beta \neq 0$ or $\alpha = \beta = 0$. Then*

(I) *$(L, [\cdot, \cdot], \Delta_r, r)$ is a coboundary Lie bialgebra iff $r \in \text{Im}(1 - \tau)$;*

(II) *$(L, [\cdot, \cdot], \Delta_r, r)$ is a triangular Lie bialgebra iff $r \in \text{Im}(1 - \tau)$ and r is α, β -symmetric to the basis of $\{e_1, e_2, e_3\}$.*

Proof. (I) Obviously, $r \in \text{Im}(1 - \tau)$ when $(L, [\cdot, \cdot], \Delta_r, r)$ is a coboundary Lie bialgebra. Conversely, if $r \in \text{Im}(1 - \tau)$, let $r = p(e_1 \otimes e_2) - p(e_2 \otimes e_1) + s(e_1 \otimes e_3) - s(e_3 \otimes e_1) + u(e_2 \otimes e_3) - u(e_3 \otimes e_2)$. It is sufficient to show that

$$(1 + \xi + \xi^2)(1 \otimes \Delta)\Delta(e_i) = 0$$

for $i = 1, 2, 3$ by [84, Proposition 2.11]. First, by computation, we have that

$$\begin{aligned}
& (1 \otimes \Delta)\Delta(e_1) \\
& = \{(e_1 \otimes e_2 \otimes e_3)(\beta ps - \beta ps) + (e_3 \otimes e_1 \otimes e_2)(\beta ps) + (e_2 \otimes e_3 \otimes e_1)(-\beta sp)\} \\
& + \{(e_1 \otimes e_3 \otimes e_2)(-\beta ps + \beta ps) + (e_3 \otimes e_2 \otimes e_1)(-\beta ps) + (e_2 \otimes e_1 \otimes e_3)(\beta sp)\} \\
& + \{(e_1 \otimes e_1 \otimes e_2)(\alpha\beta su) + (e_1 \otimes e_2 \otimes e_1)(-\beta\alpha su) + (e_2 \otimes e_1 \otimes e_1)(0)\} \\
& + \{(e_1 \otimes e_1 \otimes e_3)(-\alpha pu) + (e_1 \otimes e_3 \otimes e_1)(\alpha pu) + (e_3 \otimes e_1 \otimes e_1)(0)\} \\
& + \{(e_2 \otimes e_1 \otimes e_2)(-\beta^2 s^2) + (e_1 \otimes e_2 \otimes e_2)(0) + (e_2 \otimes e_2 \otimes e_1)(\beta^2 s^2)\} \\
& + \{(e_3 \otimes e_3 \otimes e_1)(p^2) + (e_3 \otimes e_1 \otimes e_3)(-p^2) + (e_1 \otimes e_3 \otimes e_3)(0)\}.
\end{aligned}$$

Thus

$$(1 + \xi + \xi^2)(1 \otimes \Delta)\Delta(e_1) = 0.$$

Similarly, we have that

$$(1 + \xi + \xi^2)(1 \otimes \Delta)\Delta(e_2) = 0, \quad (1 + \xi + \xi^2)(1 \otimes \Delta)\Delta(e_3) = 0.$$

Thus $(L, [\], \Delta_r, r)$ is a coboundary Lie bialgebra.

(II) It follows from Theorem 13.5.2 and Proposition 13.4.2. $\square$

Example 13.5.4. Under Example 13.4.3, we have the following:

- (i) $(sl(2), [\], \Delta_r, r)$ is a coboundary Lie bialgebra iff $r \in \text{Im}(1 - \tau)$;
- (ii) $(sl(2), [\], \Delta_r, r)$ is a triangular Lie bialgebra iff r is 0,0- symmetric to the basis of $\{e_1, e_2, e_3\}$ iff $r = s(e_1 \otimes e_3) + s(e_3 \otimes e_1) + u(e_2 \otimes e_3) + u(e_3 \otimes e_2)$.

Theorem 13.5.5. Let L be a Lie algebra with a basis $\{e_1, e_2, e_3\}$ such that

$$[e_1, e_2] = 0, [e_1, e_3] = e_1 + \beta e_2, [e_2, e_3] = \delta e_2,$$

where $\delta, \beta \in k$ and $\delta = 1$ when $\beta \neq 0$. Let $p, s, u \in k$ and $r = p(e_1 \otimes e_2) - p(e_2 \otimes e_1) + s(e_1 \otimes e_3) - s(e_3 \otimes e_1) + u(e_2 \otimes e_3) - u(e_3 \otimes e_2)$. Then

(I) $(L, [\], \Delta_r, r)$ is a coboundary Lie bialgebra iff $r \in \text{Im}(1 - \tau)$ and

$$(\delta + 1)((\delta + 1)u + \beta s)s = 0;$$

(II) $(L, [\], \Delta_r, r)$ is a triangular Lie bialgebra iff $r \in \text{Im}(1 - \tau)$ and

$$\beta s + (1 + \delta)us = 0.$$

Proof . (I) We get by computation

$$\begin{aligned}
& (1 \otimes \Delta)\Delta(e_1) \\
&= \{(e_1 \otimes e_1 \otimes e_2)(\beta\delta ss - \delta us) + (e_1 \otimes e_2 \otimes e_1)(-\beta\delta ss + \delta us) + (e_2 \otimes e_1 \otimes e_1)(0)\} \\
&+ \{(e_2 \otimes e_2 \otimes e_1)(-\beta us + uu + \beta^2 s^2 - \beta su) \\
&+ (e_2 \otimes e_1 \otimes e_2)(\beta us - uu + \beta us - \beta^2 s^2) + (e_1 \otimes e_2 \otimes e_2)(0)\} \\
& (1 \otimes \Delta)\Delta(e_2) \\
&= \{(e_1 \otimes e_1 \otimes e_2)(\delta^2 ss) + (e_1 \otimes e_2 \otimes e_1)(-\delta^2 s^2) + (e_2 \otimes e_1 \otimes e_1)(0)\} \\
&+ \{(e_2 \otimes e_2 \otimes e_1)(\delta\beta ss - \delta su) + (e_2 \otimes e_1 \otimes e_2)(\delta su - \delta\beta ss) + (e_1 \otimes e_2 \otimes e_2)(0)\} \\
& (1 \otimes \Delta)\Delta(e_3) \\
&= \{(e_1 \otimes e_2 \otimes e_3)(\delta su + \beta ss) + (e_2 \otimes e_3 \otimes e_1)(-\delta us - \beta ss) \\
&+ (e_3 \otimes e_1 \otimes e_2)(\delta\delta us + \beta\delta ss + \beta ss - su)\} \\
&+ \{(e_1 \otimes e_3 \otimes e_2)(-\delta su - \beta ss) + (e_3 \otimes e_2 \otimes e_1)(-\beta\delta ss - \delta\delta us - \beta ss + su) \\
&+ (e_2 \otimes e_1 \otimes e_3)(\delta us + \beta ss)\} \\
&+ \{(e_1 \otimes e_1 \otimes e_2)(-\delta sp - \delta\delta ps + sp + \delta sp) \\
&+ (e_1 \otimes e_2 \otimes e_1)(\delta ps + \delta\delta ps - \delta sp - sp) + (e_2 \otimes e_1 \otimes e_1)(0)\} \\
&+ \{(e_1 \otimes e_1 \otimes e_3)(ss) + (e_1 \otimes e_3 \otimes e_1)(-ss) + (e_3 \otimes e_1 \otimes e_1)(0)\} \\
&+ \{(e_2 \otimes e_2 \otimes e_1)(-\beta ps + pu - \delta\beta sp - \delta\delta up - \delta\beta ps + \delta pu - \beta sp - \delta up) \\
&+ (e_2 \otimes e_1 \otimes e_2)(\beta ps - pu + \delta\beta sp + \delta\delta up + \delta\beta ps - \delta pu + \beta sp + \delta up)) \\
&- (e_1 \otimes e_2 \otimes e_2)(0)\} \\
&+ \{(e_2 \otimes e_2 \otimes e_3)(\delta\delta uu + \delta\beta us + \beta\delta us + \beta\beta ss) \\
&+ (e_2 \otimes e_3 \otimes e_2)(-\delta\delta uu - \delta\beta us - \beta\delta us - \beta\beta ss) + (e_3 \otimes e_2 \otimes e_2)(0)\}.
\end{aligned}$$

Consequently,

$$(1 + \xi + \xi^2)(1 \otimes \Delta)\Delta(e_1) = 0,$$

$$(1 + \xi + \xi^2)(1 \otimes \Delta)\Delta(e_2) = 0$$

and

$$(1 + \xi + \xi^2)(1 \otimes \Delta)\Delta(e_3) = 0$$

iff

$$\delta^2 us + \delta\beta s^2 + \beta s^2 - us = 0.$$

This implies that $(L, [\], \Delta_r, r)$ is a coboundary Lie bialgebra iff $r \in \text{Im}(1 - \tau)$ and

$$(\delta + 1)((\delta + 1)u + \beta s)s = 0 \quad .$$

(II) It follows from Theorem 13.5.2 and Proposition 13.4.4. $\square$

Theorem 13.5.6. *If L is a Lie algebra with $\dim L = 2$ and $r \in L \otimes L$, then $(L, [\], \Delta_r, r)$ is a triangular Lie bialgebra iff $(L, [\], \Delta_r, r)$ is a coboundary Lie bialgebra iff $r \in \text{Im}(1 - \tau)$.*

Proof. It is an immediate consequence of the main result of [84]. $\square$

13.6 The solutions of CYBE with char $k \neq 2$

In this section, we find the general solution of CYBE for Lie algebra L with $\dim L = 3$ and $\dim L' = 2$, where $\text{char } k \neq 2$. If k is not algebraically closed, let P be algebraic closure of k . we can construct a Lie algebra $L_P = P \otimes L$ over P , as in [59, Section 8].

By [59, P11–14], we have that

Lemma 13.6.1. *Let L be a vector space over k . Then L is a Lie algebra over field k with $\dim L = 3$ and $\dim L' = 2$ iff there is a basis e_1, e_2, e_3 in L such that $[e_1, e_2] = 0$, $[e_1, e_3] = \alpha e_1 + \beta e_2$, $[e_2, e_3] = \gamma e_1 + \delta e_2$, where $\alpha, \beta, \gamma, \delta \in k$, and $\alpha\delta - \beta\gamma \neq 0$.*

In following three sections, we only study the Lie algebra L in Lemma 13.6.1. Set $A = \begin{pmatrix} \alpha & \gamma \\ \beta & \delta \end{pmatrix}$. Thus A is similar to $\begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}$ or $\begin{pmatrix} \lambda_1 & 0 \\ 1 & \lambda_1 \end{pmatrix}$ in the algebraic closure P of k . Therefore, there is an invertible matrix D over P such that $AD = D \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}$, or $AD = D \begin{pmatrix} \lambda_1 & 0 \\ 1 & \lambda_2 \end{pmatrix}$. Let $Q = \begin{pmatrix} D & 0 \\ 0 & \frac{1}{\lambda_1} \end{pmatrix}$ and $(e'_1, e'_2, e'_3) = (e_1, e_2, e_3)Q$. By computation, we have that $[e'_1, e'_2] = 0$, $[e'_1, e'_3] = e'_1 + \beta' e'_2$, $[e'_2, e'_3] = \delta' e'_2$, where $\beta' = 0$ and $\delta' = \frac{\lambda_1}{\lambda_2}$ when A is similar to $\begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}$; $\beta' = \frac{1}{\lambda_1}$ and $\delta' = 1$ when A is similar to $\begin{pmatrix} \lambda_1 & 0 \\ 1 & \lambda_1 \end{pmatrix}$. Let $Q = (q_{ij})_{3 \times 3}$ and $Q^{-1} = (\bar{q}_{ij})_{3 \times 3}$. If $r = \sum_{i,j=1}^3 k_{ij}(e_i \otimes e_j) = \sum_{i,j=1}^3 k'_{ij}(e'_i \otimes e'_j)$, where $k_{ij} \in k$, $k'_{ij} \in P$ for $i, j = 1, 2, 3$, then

$$k'_{ij} = \sum_{m,n}^3 k_{mn} \bar{q}_{im} \bar{q}_{jn} \text{ and } k_{ij} = \sum_{m,n}^3 k'_{mn} q_{im} q_{jn}$$

for $i, j = 1, 2, 3$. Obviously, $k_{33} = k'_{33}$.

Lemma 13.6.2. (i) $k_{i3} = k_{3i}$, for $i = 1, 2, 3$ iff $k'_{i3} = k'_{3i}$ for $i = 1, 2, 3$;

(ii) $k_{i3} = -k_{3i}$ for $i = 1, 2, 3$ iff $k'_{i3} = -k'_{3i}$ for $i = 1, 2, 3$;

(iii) $k_{ij} = k_{ji}$ for $i, j = 1, 2, 3$ iff $k'_{ij} = k'_{ji}$ for $i, j = 1, 2, 3$;

(iv) $k_{ij} = -k_{ji}$ for $i, j = 1, 2, 3$ iff $k'_{ij} = -k'_{ji}$ for $i, j = 1, 2, 3$.

Proof (i) If $k'_{i3} = k'_{3i}$ for $i = 1, 2, 3$, we see that

$$\begin{aligned}
k_{i3} &= \sum_{m,n}^3 k'_{mn} q_{im} q_{3n} \\
&= \sum_m^3 k'_{m3} q_{im} q_{33} \quad (\text{since } q_{31} = q_{32} = 0) \\
&= \sum_m^3 k'_{3m} q_{im} q_{33} \quad (\text{by assumption}) \\
&= \sum_m^3 k'_{3m} q_{33} q_{im} \\
&= \sum_{m,n}^3 k'_{mn} q_{3n} q_{im} \\
&= k_{3i}.
\end{aligned}$$

Therefore, $k_{i3} = k_{3i}$ for $i = 1, 2, 3$. The others can be proved similarly. $\square$

Theorem 13.6.3. *Let L be a Lie algebra with a basis e_1, e_2, e_3 such that $[e_1, e_2] = 0$, $[e_1, e_3] = \alpha e_1 + \beta e_2$, $[e_2, e_3] = \gamma e_1 + \delta e_2$, where $\alpha, \beta, \gamma, \delta \in k$, and $\alpha\delta - \beta\gamma \neq 0$. Let $p, q, s, t, u, v, x, y, z \in k$. Then r is a solution of CYBE iff r is strongly symmetric, or $r = p(e_1 \otimes e_2) + q(e_2 \otimes e_1) + s(e_1 \otimes e_3) - s(e_3 \otimes e_1) + u(e_2 \otimes e_3) - u(e_3 \otimes e_2) + x(e_1 \otimes e_1) + y(e_2 \otimes e_2)$ with $s(2\alpha x + \gamma(p + q)) = u(2\delta y + \beta(q + p)) = u(2\alpha x + \gamma(q + p)) = s(2\delta y + \beta(q + p)) = (\alpha - \delta)us + \gamma u^2 - \beta s^2 = s(2\gamma y + 2\beta x + (\alpha + \delta)(q + p)) = u(2\gamma y + 2\beta x + (\alpha + \delta)(p + q)) = 0$.*

Proof Let $r = \sum_{i,j=1}^3 k_{ij}(e_i \otimes e_j) \in L \otimes L$, and $k_{ij} \in k$, with $i, j = 1, 2, 3$. By computation, for all $i, j, n = 1, 2, 3$, we have that the coefficient of $e_j \otimes e_i \otimes e_i$ in $[r^{12}, r^{23}]$ is zero and the coefficient of $e_i \otimes e_i \otimes e_j$ in $[r^{13}, r^{23}]$ is zero.

We can obtain the following equations by seeing the coefficient of $e_i \otimes e_j \otimes e_n$ in $[r^{12}, r^{13}] + [r^{12}, r^{23}] + [r^{13}, r^{23}]$, as in [131, Proposition 2.6]. To simplify notation, let $k_{11} = x, k_{22} = y, k_{33} = z, k_{12} = p, k_{21} = q, k_{13} = s, k_{31} = t, k_{23} = u, k_{32} = v$.

- (1) $-\alpha s x + \alpha x t - \gamma s q + \gamma p t = 0$;
- (2) $-\beta u p + \beta q v - \delta u y + \delta y v = 0$;
- (3) $-\alpha v s + \alpha p z - \gamma u v + \gamma y z - \beta s^2 + \beta x z - \delta s u + \delta z p = 0$;
- (4) $-\beta x z + \beta s t - \delta z q + \delta u t - \alpha u t + \alpha q z - \gamma u v + \gamma y z = 0$;
- (5) $-\alpha z p + \alpha t v - \gamma z y + \gamma u v - \beta z x + \beta t s - \delta z p + \delta v s = 0$;
- (6) $-\alpha z p + \alpha s v - \gamma z y + \gamma u v - \beta s t + \beta x z - \delta s v + \delta z p = 0$;
- (7) $-\beta z x + \beta t t - \delta z q + \delta v t - \alpha z q + \alpha t u - \gamma z y + \gamma u v = 0$;
- (8) $-\beta s t + \beta x z - \delta t u + \delta q z - \alpha u s + \alpha q z - \gamma u u + \gamma y z = 0$;
- (9) $-\alpha t p + \alpha x v - \gamma t y + \gamma q v - \alpha s p + \alpha v x - \gamma s y + \gamma p v = 0$;
- (10) $-\alpha u x + \alpha q t - \gamma u q + \gamma y t - \alpha u x + \alpha q s - \gamma u p + \gamma y s = 0$;
- (11) $-\alpha s t + \alpha x z - \gamma t u + \gamma q z - \alpha s^2 + \alpha x z - \gamma s u + \gamma p z = 0$;
- (12) $-\alpha z x + \alpha t t - \gamma z q + \gamma v t - \alpha z x + \alpha s t - \gamma z p + \gamma v s = 0$;

- (13) $-\beta vx + \beta pt - \delta vq + \delta yt - \beta ux + \beta qt - \delta uq + \delta yt = 0;$
- (14) $-\beta sp - \beta xv - \delta sy + \delta pv - \beta sq + \beta xu - \delta sy + \delta pu = 0;$
- (15) $-\beta vs + \beta pz - \delta vu + \delta yz - \beta su + \beta zq - \delta u^2 + \delta yz = 0;$
- (16) $-\beta zp + \beta tv - \delta zy + \delta vv - \beta zq + \beta tu - \delta zy + \delta vu = 0;$
- (17) $-\gamma zq + \gamma ut - \gamma sv + \gamma pz = 0;$
- (18) $-\alpha vx + \alpha pt - \gamma vq + \gamma yt - \beta sx + \beta xt - \delta sq + \delta pt - \alpha sq + \alpha xu - \gamma sy + \gamma pu = 0;$
- (19) $-\beta pt + \beta vx - \delta ty + \delta qv - \alpha up + \alpha qv - \gamma uy + \gamma yv - \beta ux + \beta qs - \delta up + \delta ys = 0;$
- (20) $-\beta zp + \beta sv - \delta zy + \delta uv - \beta ut + \beta qz - \delta uv + \delta yz = 0;$
- (21) $-\beta zs + \beta tz - \delta zu + \delta vz = 0;$
- (22) $-\alpha zs + \alpha tz - \gamma zu + \gamma vz = 0;$

It is clear that r is a solution of CYBE iff (1)-(22) hold.

By simple computation, we have the sufficiency. Now we show the necessity, If $k_{33} \neq 0$, then $k'_{33} \neq 0$ and so r is a strongly symmetric element in $L_P \otimes L_P$ by [131, The proof of Proposition 1.6]. Thus r is a strongly symmetric element in $L \otimes L$. If $k_{33} = 0$, then $k'_{33} = 0$. By [131, Proposition 1.6], we have that $k'_{i3} = -k'_{3i}$ for $i = 1, 2, 3$, which implies that $k_{i3} = -k_{3i}$ for $i = 1, 2, 3$ by Lemma 13.6.2.

It immediately follows from (1)-(22) that

- (23) $s(-2\alpha x - \gamma(q + p)) = 0;$
- (24) $u(2\delta y + \beta(q + p)) = 0;$
- (25) $\gamma u^2 - \beta s^2 + (\alpha - \delta)us = 0;$
- (26) $u(2\alpha x + \gamma(q + p)) = 0;$
- (27) $s(2\delta y + \beta(q + p)) = 0;$
- (28) $2\alpha ux - 2\gamma ys - 2\beta xs + (-s(\alpha + \delta) + \gamma u)(q + p) = 0;$
- (29) $-2\beta ux + 2\delta ys - 2\gamma uy + (-u(\alpha + \delta) + \beta s)(q + p) = 0;$

By (26), (27), (28) and (29), we have that

- (30) $s(2\gamma y + 2\beta x + (\alpha + \delta)(q + p)) = 0;$
- (31) $u(2\gamma y + 2\beta x + (\alpha + \delta)(p + q)) = 0. \quad \square$

Example 13.6.4. Let L be a Lie algebra over real field R with $\dim L = 3$ and $\dim L' = 2$. If there is complex characteristic root $\lambda_1 = a + bi$ of A and the root is not real, then $r \in L \otimes L$ is a solution of CYBE iff r is strongly symmetric, or $r = p(e_1 \otimes e_2) + q(e_2 \otimes e_1) + x(e_1 \otimes e_1) + y(e_2 \otimes e_2)$ for any $p, q, x, y \in R$.

Proof. There are two different characteristic roots: $\lambda_1 = a + bi$ and $\lambda_2 = a - bi$, where $a, b \in R$. Thus A must be similar to $\begin{pmatrix} a & b \\ -b & a \end{pmatrix}$. By Theorem 13.6.3, we can complete the proof. $\square$

13.7 The solutions of CYBE with char $k = 2$

In this section, we find the general solution of CYBE for Lie algebra L with $\dim L = 3$ and $\dim L' = 2$, where char $k = 2$.

Theorem 13.7.1. *Let L be a Lie algebra with a basis e_1, e_2, e_3 such that $[e_1, e_2] = 0$, $[e_1, e_3] = \alpha e_1 + \beta e_2$, $[e_2, e_3] = \gamma e_1 + \delta e_2$, where $\alpha, \beta, \gamma, \delta \in k$, and $\alpha\delta - \beta\gamma \neq 0$. Let $p, q, s, t, u, v, x, y, z \in k$ and $A = \begin{pmatrix} \alpha & \gamma \\ \beta & \delta \end{pmatrix}$.*

(I) *If two characteristic roots of A are equal and A is similar to a diagonal matrix in the algebraic closure P of k , then r is a solution of CYBE in L for any $r \in L \otimes L$;*

(II) *If the condition in Part (I) does not hold, then r is a solution of CYBE in L iff $r = p(e_1 \otimes e_1) + p(e_2 \otimes e_1) + s(e_1 \otimes e_3) + s(e_3 \otimes e_1) + u(e_2 \otimes e_3) + u(e_3 \otimes e_2) + x(e_1 \otimes e_1) + y(e_2 \otimes e_2) + z(e_3 \otimes e_3)$ with $\alpha us + \alpha pz + \gamma u^2 + \gamma yz + \beta s^2 + \beta xz + \delta su + \delta zp = 0$ and $z \neq 0$; or $r = p(e_1 \otimes e_2) + q(e_2 \otimes e_1) + s(e_1 \otimes e_3) + s(e_3 \otimes e_1) + u(e_2 \otimes e_3) + u(e_3 \otimes e_2) + x(e_1 \otimes e_1) + y(e_2 \otimes e_2)$ with $s\gamma(p+q) = u\beta(p+q) = u\gamma(p+q) = s\beta(p+q) = (\alpha+\delta)us + \gamma u^2 + \beta s^2 = s(\alpha+\delta)(p+q) = u(\alpha+\delta)(p+q) = 0$.*

Proof We only show the necessity since the sufficiency can easily be shown. By the proof of Theorem 13.6.3, there exists an invertible matrix Q such that $(e'_1, e'_2, e'_3) = (e_1, e_2, e_3)Q$ and $[e'_1, e'_2] = 0$, $[e'_1, e'_3] = e'_1 + \beta' e'_2$, $[e'_2, e'_3] = \delta' e'_2$. We use the notations before Lemma 13.6.2. By [11, Proposition 2.4], $k'_{i3} = k'_{3i}$ for $i = 1, 2, 3$, which implies that $k_{3i} = k_{i3}$ for $i = 1, 2, 3$ by Lemma 13.6.2.

(I) If A is similar to $\begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_1 \end{pmatrix}$, then $\beta' = 0$ and $\delta' = 1$. By [11, Proposition 2.4], we have Part (I).

(II) Let A be not similar to $\begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_1 \end{pmatrix}$.

(a). If $z \neq 0$, then $k'_{33} \neq 0$. Thus $k'_{12} = k'_{21}$ by [136, Proposition 2.4], which implies $k_{12} = k_{21}$. It is straightforward to check that relation (1)-(22) in the proof of Theorem 13.6.3 hold iff $\alpha us + \alpha pz + \gamma u^2 + \gamma yz + \beta s^2 + \beta xz + \delta su + \delta zp = 0$.

(b). If $z = 0$, then we can obtain that r is the second case in Part (II) by using the method similar to the proof of Theorem 13.6.3. $\square$

13.8 Coboundary Lie bialgebras

In this section, using the general solution, which are obtained in the section above, of CYBE in Lie algebra L with $\dim L = 3$ and $\dim L' = 2$, we give the sufficient and necessary conditions which $(L, [\cdot, \cdot], \Delta_r, r)$ is a coboundary (or triangular) Lie bialgebra.

Theorem 13.8.1. *Let L be a Lie algebra with a basis e_1, e_2, e_3 such that $[e_1, e_2] = 0$, $[e_1, e_3] = \alpha e_1 + \beta e_2$, $[e_2, e_3] = \gamma e_1 + \delta e_2$, where $\alpha, \beta, \gamma, \delta \in k$, and $\alpha\delta - \beta\gamma \neq 0$. Set $A = \begin{pmatrix} \alpha & \gamma \\ \beta & \delta \end{pmatrix}$. Let $p, u, s \in k$, $r \in \text{Im}(1 - \tau)$ and $r = p(e_1 \otimes e_2) - p(e_2 \otimes e_1) + s(e_1 \otimes e_3) - s(e_3 \otimes e_1) + u(e_2 \otimes e_3) - u(e_3 \otimes e_2)$. Then*

(I) *$(L, [\cdot, \cdot], \Delta_r, r)$ is a coboundary Lie bialgebra iff*

$$(s, u) \begin{pmatrix} \beta\delta + \alpha\beta & -\beta\gamma - \alpha^2 \\ \delta^2 + \gamma\beta & -\delta\gamma - \gamma\alpha \end{pmatrix} \begin{pmatrix} s \\ u \end{pmatrix} = 0;$$

(II) If two characteristic roots of A are equal and A is similar to a diagonal matrix in the algebraic closure P of k with $\text{char } k = 2$, then $(L, [\cdot, \cdot], \Delta_r, r)$ is a triangular Lie bialgebra for any $r \in \text{Im}(1 - \tau)$;

(III) If the condition in Part(II) does not hold, then $(L, [\cdot, \cdot], \Delta_r, r)$ is a triangular Lie bialgebra iff $-\beta s^2 + \gamma u^2 + (\alpha - \delta)us = 0$.

Proof We can complete the proof as in the proof of [131, Theorem 3.3]. $\square$

Example 13.8.2. Under Example 13.6.4, and $r \in \text{Im}(1 - \tau)$, we have the following :

(I) $(L, [\cdot, \cdot], \Delta_r, r)$ is a coboundary Lie bialgebra iff $r = p(e_1 \otimes e_2) - p(e_2 \otimes e_1) + s(e_1 \otimes e_3) - s(e_3 \otimes e_1) + u(e_2 \otimes e_3) - u(e_3 \otimes e_2)$ with $a(s^2 + u^2) = 0$;

(II) $(L, [\cdot, \cdot], \Delta_r, r)$ is a triangular Lie bialgebra iff $r = p(e_1 \otimes e_2) - p(e_2 \otimes e_1)$.

Chapter 14

Classification of PM Quiver Hopf Algebras

In noncommutative algebras quivers and associated path algebras are intensively studied because of a remarkable interaction between homology theory, algebraic geometry, and Lie theory. Ringel's approach to quantum groups via quivers suggested that there may be an interesting overlapping also between the theory of quivers and Hopf algebras. Some years ago Cibils and Rosso [36, 37] started to study quivers admitting a (graded) Hopf algebra structure and there are many papers (e.g. [21, 94]) follow these works. The representation theory of (graded) Hopf algebras has applications in many branches of physics and mathematics such as the construction of solutions to the quantum Yang-Baxter equation and topological invariants (see e.g. [74, 108]). Quivers have also been used in gauge theory and string theory (see e.g. [144, 105]).

This chapter can be viewed an extension of the analysis of Cibils and Rosso. Let G be a group and kG be the group algebra of G over a field k . It is well-known [37] that the kG -Hopf bimodule category ${}^{kG}_k\mathcal{M}_k^{kG}$ is equivalent to the direct product category $\prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$, where $\mathcal{K}(G)$ is the set of conjugate classes in G , $u : \mathcal{K}(G) \rightarrow G$ is a map such that $u(C) \in C$ for any $C \in \mathcal{K}(G)$, $Z_{u(C)} = \{g \in G \mid gu(C) = u(C)g\}$ and $\mathcal{M}_{kZ_{u(C)}}$ denotes the category of right $kZ_{u(C)}$ -modules (see [37, Proposition 3.3] [36, Theorem 4.1] and Theorem 14.1.2). Thus for any Hopf quiver (Q, G, r) , the kG -Hopf bimodule structures on the arrow comodule kQ_1^c can be derived from the right $kZ_{u(C)}$ -module structures on ${}^{u(C)}(kQ_1^c)^1$ for all $C \in \mathcal{K}(G)$. If the arrow comodule kQ_1^c admits a kG -Hopf bimodule structure, then there exist six graded Hopf algebras: co-path Hopf algebra kQ^c , one-type-co-path Hopf algebras $kG[kQ_1^c]$, one-type-path Hopf algebras $(kG)^*[kQ_1^a]$, semi-path Hopf algebra kQ^s , semi-co-path Hopf algebra kQ^{sc} and path Hopf algebra kQ^a . We call these Hopf algebras quiver Hopf algebras over G . If the corresponding $kZ_{u(C)}$ -modules ${}^{u(C)}(kQ_1^c)^1$ is pointed (i.e. it is zero or a direct sum of one dimensional $kZ_{u(C)}$ -modules) for all $C \in \mathcal{K}(G)$, then kQ_1^c is called a PM kG -Hopf bimodule. The six graded Hopf algebras derived from the PM kG -Hopf bimodule kQ_1^c are called PM quiver Hopf algebras. A Yetter-Drinfeld H -module V is *pointed* if $V = 0$ or V is a direct sum of one dimensional YD H -modules. If V is a pointed YD H -module, then the corresponding Nichols algebra $\mathcal{B}(V)$ is called a PM *Nichols algebra*. For example, when G is a finite abelian group of exponent m and k contains a primitive

m -th root of 1 (e.g. k is the complex field), all quiver Hopf algebras are PM (see Lemma 14.1.3), all YD kG -modules are pointed and all Nichols algebras of all YD kG -modules are PM (see Lemma 14.2.3).

The aim of this chapter is to provide parametrization of PM quiver Hopf algebras, multiple Taft algebras and PM Nichols algebras. In other words, this chapter provides a kind of classification of these Hopf algebras. This chapter is organized as follows. In Section 14.1, we examine the PM quiver Hopf algebras by means of ramification system with characters. In Section 14.2, we describe PM Nichols algebras and multiple Taft algebras by means of element system with characters. In Section 14.3, we show that the diagram of a quantum weakly commutative multiple Taft algebra is not only a Nichols algebra but also a quantum linear space in ${}^{kG}_{kG}\mathcal{YD}$; the diagram of a semi-path Hopf algebra of ESC is a quantum tensor algebra in ${}^{kG}_{kG}\mathcal{YD}$; the quantum enveloping algebra of a complex semisimple Lie algebra is a quotient of a semi-path Hopf algebra.

Throughout this chapter, we work over a fixed field k . All algebras, coalgebras, Hopf algebras, and so on, are defined over k ; $\dim$, $\otimes$ and Hom stand for $\dim_k$, $\otimes_k$ and Hom_k , respectively. Books [40, 87, 114] provide the necessary background for Hopf algebras and book [3] provides a nice description of the path algebra approach.

Let $\mathbb{Z}$, $\mathbb{Z}^+$ and $\mathbb{N}$ denote sets of all integers, all positive integers and all non-negative integers, respectively. For sets X and Y , we denote by $|X|$ the cardinal number of X and by X^Y or $X^{\mathbb{N}}$ the Cartesian product $\prod_{y \in Y} X_y$ with $X_y = X$ for any $y \in Y$. If X is finite, then $|X|$ is the number of elements in X . If $X = \bigoplus_{i \in I} X_{(i)}$ as vector spaces, then we denote by ι_i the natural injection from $X_{(i)}$ to X and by π_i the corresponding projection from X to $X_{(i)}$. We will use μ to denote the multiplication of an algebra, Δ to denote the comultiplication of a coalgebra, α^- , α^+ , δ^- and δ^+ to denote the left module, right module, left comodule and right comodule structure maps, respectively. The Sweedler's sigma notations for coalgebras and comodules are $\Delta(x) = \sum x_{(1)} \otimes x_{(2)}$, $\delta^-(x) = \sum x_{(-1)} \otimes x_{(0)}$, $\delta^+(x) = \sum x_{(0)} \otimes x_{(1)}$. Let G be a group. We denote by $Z(G)$ the center of G . Let $\widehat{G}$ denote the set of characters of all one-dimensional representations of G . It is clear that $\widehat{G} = \{\chi \mid \chi \text{ is a group homomorphism from } G \text{ to the multiplicative group of all non-zero elements in } k\}$.

A quiver $Q = (Q_0, Q_1, s, t)$ is an oriented graph, where Q_0 and Q_1 are the sets of vertices and arrows, respectively; s and t are two maps from Q_1 to Q_0 . For any arrow $a \in Q_1$, $s(a)$ and $t(a)$ are called its start vertex and end vertex, respectively, and a is called an arrow from $s(a)$ to $t(a)$. For any $n \geq 0$, an n -path or a path of length n in the quiver Q is an ordered sequence of arrows $p = a_n a_{n-1} \cdots a_1$ with $t(a_i) = s(a_{i+1})$ for all $1 \leq i \leq n-1$. Note that a 0-path is exactly a vertex and a 1-path is exactly an arrow. In this case, we define $s(p) = s(a_1)$, the start vertex of p , and $t(p) = t(a_n)$, the end vertex of p . For a 0-path x , we have $s(x) = t(x) = x$. Let Q_n be the set of n -paths. Let ${}^y Q_n^x$ denote the set of all n -paths from x to y , $x, y \in Q_0$. That is, ${}^y Q_n^x = \{p \in Q_n \mid s(p) = x, t(p) = y\}$.

A quiver Q is *finite* if Q_0 and Q_1 are finite sets. A quiver Q is *locally finite* if ${}^y Q_1^x$ is a finite set for any $x, y \in Q_0$.

Let G be a group. Let $\mathcal{K}(G)$ denote the set of conjugate classes in G . A formal sum $r =$

$\sum_{C \in \mathcal{K}(G)} r_C C$ of conjugate classes of G with cardinal number coefficients is called a *ramification* (or *ramification data*) of G , i.e. for any $C \in \mathcal{K}(G)$, r_C is a cardinal number. In particular, a formal sum $r = \sum_{C \in \mathcal{K}(G)} r_C C$ of conjugate classes of G with non-negative integer coefficients is a ramification of G .

For any ramification r and a $C \in \mathcal{K}(G)$, since r_C is a cardinal number, we can choose a set $I_C(r)$ such that its cardinal number is r_C without lost generality. Let $\mathcal{K}_r(G) := \{C \in \mathcal{K}(G) \mid r_C \neq 0\} = \{C \in \mathcal{K}(G) \mid I_C(r) \neq \emptyset\}$. If there exists a ramification r of G such that the cardinal number of ${}^y Q_1^x$ is equal to r_C for any $x, y \in G$ with $x^{-1}y \in C \in \mathcal{K}(G)$, then Q is called a *Hopf quiver with respect to the ramification data r* . In this case, there is a bijection from $I_C(r)$ to ${}^y Q_1^x$, and hence we write ${}^y Q_1^x = \{a_{y,x}^{(i)} \mid i \in I_C(r)\}$ for any $x, y \in G$ with $x^{-1}y \in C \in \mathcal{K}(G)$. Denote by (Q, G, r) the Hopf quiver of G with respect to r .

The coset decomposition of $Z_{u(C)}$ in G is

$$G = \bigcup_{\theta \in \Theta_C} Z_{u(C)} g_\theta, \quad (14.1)$$

where Θ_C is an index set. It is easy to check that $|\Theta_C| = |C|$. We always assume that the representative element of the coset $Z_{u(C)}$ is the identity 1 of G . We claim that $\theta = \eta$ if $g_\theta^{-1}u(C)g_\theta = g_\eta^{-1}u(C)g_\eta$. In fact, by the equation $g_\theta^{-1}u(C)g_\theta = g_\eta^{-1}u(C)g_\eta$ one gets $g_\eta g_\theta^{-1}u(C) = u(C)g_\eta g_\theta^{-1}$. Thus $g_\eta g_\theta^{-1} \in Z_{u(C)}$, and hence $\theta = \eta$. For any $x, y \in G$ with $x^{-1}y \in C \in \mathcal{K}(G)$, there exists a unique $\theta \in \Theta_C$ such that

$$x^{-1}y = g_\theta^{-1}u(C)g_\theta. \quad (14.2)$$

Without specification, we will always assume that x, y, θ and C satisfy the above relation (14.2). Note that θ is only determined by $x^{-1}y$. For any $h \in G$ and $\theta \in \Theta_C$, there exist unique $h' \in Z_{u(C)}$ and $\theta' \in \Theta_C$ such that $g_\theta h = h' g_{\theta'}$. Let $\zeta_\theta(h) = h'$. Then we have

$$g_\theta h = \zeta_\theta(h) g_{\theta'}. \quad (14.3)$$

If $u(C)$ lies in the center $Z(G)$ of G , we have $\zeta_\theta = id_G$. In particular, if G is abelian, then $\zeta_\theta = id_G$ since $Z_{u(C)} = G$.

Let H be a Hopf algebra. A (left-left) *Yetter-Drinfeld module* V over H (simply, YD H -module) is simultaneously a left H -module and a left H -comodule satisfying the following compatibility condition:

$$\sum (h \cdot v)_{(-1)} \otimes (h \cdot v)_{(0)} = \sum h_{(1)} v_{(-1)} S(h_{(3)}) \otimes h_{(2)} \cdot v_{(0)}, \quad v \in V, h \in H. \quad (14.4)$$

We denote by ${}^H_H \mathcal{YD}$ the category of YD H -modules; the morphisms in this category preserve both the action and the coaction of H .

The structure of a Nichols algebra appeared first in the paper [91], and N. Andruskiewitsch and H. J. Schneider used it to classify finite-dimensional pointed Hopf algebras [4, 5, 7, 6]. Its definition can be found in [7, Definition 2.1].

If $\phi : A \rightarrow A'$ is an algebra homomorphism and (M, α^-) is a left A' -module, then M becomes a left A -module with the A -action given by $a \cdot x = \phi(a) \cdot x$ for any $a \in A, x \in M$, called a pullback

A -module through ϕ , written as ${}_{\phi}M$. Dually, if $\phi : C \rightarrow C'$ be a coalgebra homomorphism and (M, δ^-) is a left C -comodule, then M is a left C' -comodule with the C' -comodule structure given by $\delta'^- := (\phi \otimes \text{id})\delta^-$, called a push-out C' -comodule through ϕ , written as ${}^{\phi}M$.

Let A be an algebra and M be an A -bimodule. Then the tensor algebra $T_A(M)$ of M over A is a graded algebra with $T_A(M)_{(0)} = A$, $T_A(M)_{(1)} = M$ and $T_A(M)_{(n)} = \otimes_A^n M$ for $n > 1$. That is, $T_A(M) = A \oplus (\bigoplus_{n>0} \otimes_A^n M)$. Let D be another algebra. If h is an algebra map from A to D and f is an A -bimodule map from M to ${}^h D_h$, then by the universal property of $T_A(M)$ (see [91, Proposition 1.4.1]) there is a unique algebra map $T_A(h, f) : T_A(M) \rightarrow D$ such that $T_A(h, f)\iota_0 = h$ and $T_A(h, f)\iota_1 = f$. One can easily see that $T_A(h, f) = h + \sum_{n>0} \mu^{n-1} T_n(f)$, where $T_n(f)$ is the map from $\otimes_A^n M$ to $\otimes_A^n D$ given by $T_n(f)(x_1 \otimes x_2 \otimes \cdots \otimes x_n) = f(x_1) \otimes f(x_2) \otimes \cdots \otimes f(x_n)$, i.e., $T_n(f) = f \otimes_A f \otimes_A \cdots \otimes_A f$. Note that μ can be viewed as a map from $D \otimes_A D$ to D .

Dually, let C be a coalgebra and let M be a C -bicomodule. Then the cotensor coalgebra $T_C^c(M)$ of M over C is a graded coalgebra with $T_C^c(M)_{(0)} = C$, $T_C^c(M)_{(1)} = M$ and $T_C^c(M)_n = \square_C^n M$ for $n > 1$. That is, $T_C^c(M) = C \oplus (\bigoplus_{n>0} \square_C^n M)$. Let D be another coalgebra. If h is a coalgebra map from D to C and f is a C -bicomodule map from ${}^h D^h$ to M such that $f(\text{corad}(D)) = 0$, then by the universal property of $T_C^c(M)$ (see [91, Proposition 1.4.2]) there is a unique coalgebra map $T_C^c(h, f)$ from D to $T_C^c(M)$ such that $\pi_0 T_C^c(h, f) = h$ and $\pi_1 T_C^c(h, f) = f$. It is not difficult to see that $T_C^c(h, f) = h + \sum_{n>0} T_n^c(f) \Delta_{n-1}$, where $T_n^c(f)$ is the map from $\square_C^n D$ to $\square_C^n M$ induced by $T_n^c(f)(x_1 \otimes x_2 \otimes \cdots \otimes x_n) = f(x_1) \otimes f(x_2) \otimes \cdots \otimes f(x_n)$, i.e., $T_n^c(f) = f \otimes f \otimes \cdots \otimes f$.

14.1 PM quiver Hopf algebras

In this section we describe PM quiver Hopf algebras by parameters.

We first describe the category of Hopf bimodules by categories of modules.

Let G be a group and let (B, δ^-, δ^+) be a kG -bicomodule. Then the (x, y) -isotypic component of B is

$${}^y B^x = \{b \in B \mid \delta^-(b) = y \otimes b, \delta^+(b) = b \otimes x\},$$

where $x, y \in G$. Let M be another kG -bicomodule and $f : B \rightarrow M$ be a kG -bicomodule homomorphism. Then $f({}^y B^x) \subseteq {}^y M^x$ for any $x, y \in G$. Denote by ${}^y f^x$ the restriction map $f|_{{}^y B^x} : {}^y B^x \rightarrow {}^y M^x$, $x, y \in G$.

Theorem 14.1.1. (See [37, Proposition 3.3] and [36, Theorem 4.1]) *The category ${}^{kG}_{kG} \mathcal{M}_{kG}^{kG}$ of kG -Hopf bimodules is equivalent to the Cartesian product category $\prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}^{kG}$ of categories $\mathcal{M}_{kZ_{u(C)}}^{kG}$ of right $kZ_{u(C)}$ -modules for all $C \in \mathcal{K}(G)$.*

For later use, we give the mutually inverse functors between the two categories here. The functors W from ${}^{kG}_{kG} \mathcal{M}_{kG}^{kG}$ to $\prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}^{kG}$ is defined by

$$W(B) = \{{}^{u(C)} B^1\}_{C \in \mathcal{K}(G)}, \quad W(f) = \{{}^{u(C)} f^1\}_{C \in \mathcal{K}(G)}$$

for any object B and morphism f in ${}^k_G\mathcal{M}_{{}^k_G}$, where the right $kZ_{u(C)}$ -module action on ${}^{u(C)}B^1$ is given by

$$b \triangleleft h = h^{-1} \cdot b \cdot h, \quad b \in {}^{u(C)}B^1, \quad h \in Z_{u(C)}. \quad (14.5)$$

The functor V from $\prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$ to ${}^k_G\mathcal{M}_{{}^k_G}$ is defined as follows:

For $M = \{M(C)\}_{C \in \mathcal{K}(G)} \in \prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$, $V(M)$ is given by

$$\begin{aligned} {}^yV(M)^x &= x \otimes M(C) \otimes_{kZ_{u(C)}} g_\theta, \\ h \cdot (x \otimes m \otimes_{kZ_{u(C)}} g_\theta) &= hx \otimes m \otimes_{kZ_{u(C)}} g_\theta, \\ (x \otimes m \otimes_{kZ_{u(C)}} g_\theta) \cdot h &= xh \otimes m \otimes_{kZ_{u(C)}} g_\theta h = xh \otimes (m \triangleleft \zeta_\theta(h)) \otimes_{kZ_{u(C)}} g_{\theta'}, \end{aligned} \quad (14.6)$$

where $h, x, y \in G$ with $x^{-1}y \in C$ and the relation (14.2) and (14.3), $m \in M(C)$. For any morphism $f = \{f_C\}_{C \in \mathcal{K}(G)} : \{M(C)\}_{C \in \mathcal{K}(G)} \rightarrow \{N(C)\}_{C \in \mathcal{K}(G)}$, $V(f)(x \otimes m \otimes_{kZ_{u(C)}} g_\theta) = x \otimes f_C(m) \otimes_{kZ_{u(C)}} g_\theta$ for any $m \in M_C$, $x, y \in G$ with $x^{-1}y = g_\theta^{-1}u(C)g_\theta$. That is, ${}^yV(f)^x = \text{id} \otimes f_C \otimes \text{id}$.

An A -module M is called *pointed* if $M = 0$ or M is a direct sum of one dimensional A -modules.

Definition 14.1.2. A kG -Hopf bimodule N is called a kG -Hopf bimodule with pointed module structure (or a PM kG -Hopf bimodule in short) if there exists an object $\prod_{C \in \mathcal{K}(G)} M(C)$ in $\prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$ such that $M(C)$ is a right pointed $kZ_{u(C)}$ -module for any $C \in \mathcal{K}(G)$ and $N \cong V(\{M(C)\}_{C \in \mathcal{K}(G)}) := \bigoplus_{y=g_\theta^{-1}u(C)g_\theta, x, y \in G} x \otimes M(C) \otimes_{kZ_{u(C)}} g_\theta$ as kG -Hopf bimodules. Here $V(\{M(C)\}_{C \in \mathcal{K}(G)}) := \bigoplus_{y=g_\theta^{-1}u(C)g_\theta, x, y \in G} x \otimes M(C) \otimes_{kZ_{u(C)}} g_\theta$ is defined in the proof of Theorem 14.1.1.

Lemma 14.1.3 – 14.1.7, Theorem 14.1.9 and Lemma 14.2.7 are well-known.

Lemma 14.1.3. Assume that G is a finite commutative group of exponent m . If k contains a primitive m -th root of 1, then (i) every kG -module is a pointed module; (ii) every kG -Hopf bimodule is PM.

Lemma 14.1.4. Let $H = \bigoplus_{i \geq 0} H_{(i)}$ be a graded Hopf algebra. Set $B := H_{(0)}$ and $M := H_{(1)}$. Then M is a B -Hopf bimodule with the B -actions and B -coactions given by

$$\alpha^- = \pi_1 \mu(\iota_0 \otimes \iota_1), \quad \alpha^+ = \pi_1 \mu(\iota_1 \otimes \iota_0), \quad \delta^- = (\pi_0 \otimes \pi_1) \Delta \iota_1, \quad \delta^+ = (\pi_1 \otimes \pi_0) \Delta \iota_1.$$

Lemma 14.1.5. (i) Let B be an algebra and M be a B -bimodule. Then the tensor algebra $T_B(M)$ of M over B admits a graded Hopf algebra structure if and only if B admits a Hopf algebra structure and M admits a B -Hopf bimodule structure.

(ii) Let B be a coalgebra and M be a B -bicomodule. Then the cotensor coalgebra $T_B^c(M)$ of M over B admits a graded Hopf algebra structure if and only if B admits a Hopf algebra structure and M admits a B -Hopf bimodule structure.

Let B be a bialgebra (Hopf algebra) and M be a B -Hopf bimodule. Then $T_B^c(M)$ is a graded bialgebra (Hopf algebra) by Lemma 14.1.5. Let $B[M]$ denote the subalgebra of $T_B^c(M)$ generated by B and M . Then $B[M]$ is a bialgebra (Hopf algebra) of type one by [91, section 2.2, p.1533]. $B[M]$ is a graded subspace of $T_B^c(M)$.

Lemma 14.1.6. *Let B and B' be two Hopf algebras. Let M and M' be B -Hopf bimodule and B' -Hopf bimodule, respectively. Assume that $\phi : B \rightarrow B'$ be a Hopf algebra map. If ψ is simultaneously a B -bimodule and B' -bicomodule map from ${}^\phi M^\phi$ to ${}^\phi M'_\phi$, then (i) $T_B(\iota_0\phi, \iota_1\psi) := \iota_0\phi + \sum_{n>0} \mu^{n-1}T_n(\iota_1\psi)$ is a graded Hopf algebra map from $T_B(M)$ to $T_{B'}(M')$. (ii) $T_{B'}^c(\phi\pi_0, \psi\pi_1) := \phi\pi_0 + \sum_{n>0} T_n^c(\psi\pi_1)\Delta_{n-1}$ is a graded Hopf algebra map from $T_B^c(M)$ to $T_{B'}^c(M')$.*

Lemma 14.1.7. *Let B and B' be two Hopf algebras. Let M and M' be B -Hopf bimodule and B' -Hopf bimodule, respectively. Then the following statements are equivalent: (i) There exists a Hopf algebra isomorphism $\phi : B \rightarrow B'$ such that $M \cong {}^\phi M'_\phi$ as B -Hopf bimodules. (ii) $T_B(M)$ and $T_{B'}(M')$ are isomorphic as graded Hopf algebras. (iii) $T_B^c(M)$ and $T_{B'}^c(M')$ are isomorphic as graded Hopf algebras. (iv) $B[M]$ and $B'[M']$ are isomorphic as graded Hopf algebras.*

Let $Q = (G, Q_1, s, t)$ be a quiver of a group G . Then kQ_1 becomes a kG -bicomodule under the natural comodule structures:

$$\delta^-(a) = t(a) \otimes a, \quad \delta^+(a) = a \otimes s(a), \quad a \in Q_1, \quad (14.7)$$

called an *arrow comodule*, written as kQ_1^c . In this case, the path coalgebra kQ^c is exactly isomorphic to the cotensor coalgebra $T_{kG}^c(kQ_1^c)$ over kG in a natural way (see [27] and [36]). We will regard $kQ^c = T_{kG}^c(kQ_1^c)$ in the following. Moreover, when G is finite, kQ_1 becomes a $(kG)^*$ -bimodule with the module structures defined by

$$p \cdot a := \langle p, t(a) \rangle a, \quad a \cdot p := \langle p, s(a) \rangle a, \quad p \in (kG)^*, a \in Q_1, \quad (14.8)$$

written as kQ_1^a , called an *arrow module*. Therefore, we have a tensor algebra $T_{(kG)^*}(kQ_1^a)$. Note that the tensor algebra $T_{(kG)^*}(kQ_1^a)$ of kQ_1^a over $(kG)^*$ is exactly isomorphic to the path algebra kQ^a . We will regard $kQ^a = T_{(kG)^*}(kQ_1^a)$ in the following.

Assume that Q is a finite quiver on finite group G . Let $\xi_{kQ_1^a}$ denote the linear map from kQ_1^a to $(kQ_1^c)^*$ by sending a to a^* for any $a \in Q_1$ and $\xi_{kQ_1^c}$ denote the linear map from kQ_1^c to $(kQ_1^a)^*$ by sending a to a^* for any $a \in Q_1$. Here $\{a^* \mid a^* \in (kQ_1)^*\}$ is the dual basis of $\{a \mid a \in Q_1\}$.

Lemma 14.1.8. *Assume that Q is a finite Hopf quiver on finite group G . Then*

(i) *If $(M, \alpha^-, \alpha^+, \delta^-, \delta^+)$ is a finite dimensional B -Hopf bimodule and B is a finite dimensional Hopf algebra, then $(M^*, \delta^{*-}, \delta^{*+}, \alpha^{*-}, \alpha^{*+})$ is a B^* -Hopf bimodule.*

(ii) *If $(kQ_1^c, \alpha^-, \alpha^+, \delta^-, \delta^+)$ is a kG -Hopf bimodule, then there exist unique left $(kG)^*$ -comodule operation $\delta_{kQ_1^a}^-$ and right $(kG)^*$ -comodule operation $\delta_{kQ_1^a}^+$ such that $(kQ_1^a, \alpha_{kQ_1^a}^-, \alpha_{kQ_1^a}^+, \delta_{kQ_1^a}^-, \delta_{kQ_1^a}^+)$ becomes a $(kG)^*$ -Hopf bimodule and $\xi_{kQ_1^a}$ becomes a $(kG)^*$ -Hopf bimodule isomorphism from $(kQ_1^a, \alpha_{kQ_1^a}^-, \alpha_{kQ_1^a}^+, \delta_{kQ_1^a}^-, \delta_{kQ_1^a}^+)$ to $((kQ_1^c)^*, \delta^{*-}, \delta^{*+}, \alpha^{*-}, \alpha^{*+})$.*

(iii) *If $(kQ_1^a, \alpha^-, \alpha^+, \delta^-, \delta^+)$ is a $(kG)^*$ -Hopf bimodule, then there exist unique left kG -module operation $\alpha_{kQ_1^c}^-$ and right kG -module operation $\alpha_{kQ_1^c}^+$ such that $(kQ_1^c, \alpha_{kQ_1^c}^-, \alpha_{kQ_1^c}^+, \delta_{kQ_1^c}^-, \delta_{kQ_1^c}^+)$ become a kG -Hopf bimodule and $\xi_{kQ_1^c}$ becomes a kG -Hopf bimodule isomorphism from $(kQ_1^c, \alpha_{kQ_1^c}^-, \alpha_{kQ_1^c}^+, \delta_{kQ_1^c}^-, \delta_{kQ_1^c}^+)$ to $((kQ_1^a)^*, \delta^{*-}, \delta^{*+}, \alpha^{*-}, \alpha^{*+})$.*

(iv) *$\xi_{kQ_1^a}$ is a $(kG)^*$ -Hopf bimodule isomorphism from $(kQ_1^a, \alpha_{kQ_1^a}^-, \alpha_{kQ_1^a}^+, \delta_{kQ_1^a}^-, \delta_{kQ_1^a}^+)$ to $((kQ_1^c)^*, \delta_{kQ_1^c}^{*-}, \delta_{kQ_1^c}^{*+}, \alpha_{kQ_1^c}^{*-}, \alpha_{kQ_1^c}^{*+})$ if and only if $\xi_{kQ_1^c}$ becomes a kG -Hopf bimodule isomorphism from $(kQ_1^c, \alpha_{kQ_1^c}^-, \alpha_{kQ_1^c}^+, \delta_{kQ_1^c}^-, \delta_{kQ_1^c}^+)$ to $((kQ_1^a)^*, \delta_{kQ_1^a}^{*-}, \delta_{kQ_1^a}^{*+}, \alpha_{kQ_1^a}^{*-}, \alpha_{kQ_1^a}^{*+})$.*

Proof. It is easy to check (i)–(iii). Now we show (iv). Let $A := kQ_1^a$, $B := kQ_1^c$. Let σ_B denote the canonical linear isomorphism from B to B^{**} by sending b to b^{**} for any $b \in B$, where $\langle b^{**}, f \rangle = \langle f, b \rangle$ for any $f \in B^*$. If $A \xrightarrow{\xi_A} B^*$ as $(kG)^*$ -Hopf bimodules, then $B \xrightarrow{\sigma_B} B^{**} \xrightarrow{(\xi_A)^*} A^*$ as kG -Hopf bimodules. It is easy to check $\xi_B = (\xi_A)^* \sigma_B$. Therefore ξ_B is a kG -Hopf bimodule isomorphism. Conversely, if $B \xrightarrow{\xi_B} A^*$ as kG -Hopf bimodules, we can similarly show that $A \cong B^*$ as $(kG)^*$ -Hopf bimodules. $\square$

Theorem 14.1.9. (see [36, Theorem 3.3] and [37, Theorem 3.1]) *Let Q be a quiver over group G . Then the following two statements are equivalent:*

(i) Q is a Hopf quiver.

(ii) Arrow comodule kQ_1^c admits a kG -Hopf bimodule structure.

Furthermore, if Q is finite, then the above are equivalent to the following:

(iii) Arrow module kQ_1^a admits a $(kG)^*$ -Hopf bimodule structure.

Assume that Q is a Hopf quiver. It follows from Theorem 14.1.9 that there exist a left kG -module structure α^- and a right kG -module structure α^+ on arrow comodule $(kQ_1^c, \delta^-, \delta^+)$ such that $(kQ_1^c, \alpha^-, \alpha^+, \delta^-, \delta^+)$ becomes a kG -Hopf bimodule, called a kG -Hopf bimodule with arrow comodule, written $(kQ_1^c, \alpha^-, \alpha^+)$ in short. We obtain three graded Hopf algebras $T_{kG}(kQ_1^c)$, $T_{kG}^c(kQ_1^c)$ and $kG[kQ_1^c]$, called semi-path Hopf algebra, co-path Hopf algebra and one-type-co-path Hopf algebra, written $kQ^s(\alpha^-, \alpha^+)$, $kQ^c(\alpha^-, \alpha^+)$ and $kG[kQ_1^c, \alpha^-, \alpha^+]$, respectively. Dually, when Q is finite, it follows from Theorem 14.1.9 that there exist a left $(kG)^*$ -comodule structure δ^- and a right $(kG)^*$ -comodule structure δ^+ on arrow module $(kQ_1^a, \alpha^-, \alpha^+)$ such that $(kQ_1^a, \alpha^-, \alpha^+, \delta^-, \delta^+)$ becomes a $(kG)^*$ -Hopf bimodule, called a $(kG)^*$ -Hopf bimodule with arrow module, written $(kQ_1^a, \delta^-, \delta^+)$ in short. We obtain three graded Hopf algebras $T_{(kG)^*}(kQ_1^a)$, $T_{(kG)^*}^c(kQ_1^a)$ and $(kG)^*[kQ_1^a]$, called path Hopf algebra, semi-co-path Hopf algebra and one-type-path Hopf algebra, written $kQ^a(\delta^-, \delta^+)$, $kQ^{sc}(\delta^-, \delta^+)$ and $(kG)^*[kQ_1^a, \alpha_1, \alpha^+]$, respectively. We call the six graded Hopf algebras the quiver Hopf algebras (over G). We usually omit the (co)module operations when we write these quiver Hopf algebras.

If $\xi_{kQ_1^a}$ or $\xi_{kQ_1^c}$ is a Hopf bimodule isomorphism, then, by Lemma 14.1.7 and Lemma 14.1.8, $T_{(kG)^*}(\iota_0, \iota_1 \xi_{kQ_1^a})$ and $T_{kG}^c(\pi_0, \xi_{kQ_1^c} \pi_1)$ are graded Hopf algebra isomorphisms from $T_{(kG)^*}(kQ_1^a)$ to $T_{(kG)^*}((kQ_1^c)^*)$ and from $T_{kG}^c(kQ_1^c)$ to $T_{kG}^c((kQ_1^a)^*)$, respectively; $T_{(kG)^*}^c(\pi_0, \xi_{kQ_1^a} \pi_1)$ and $T_{kG}(\iota_0, \iota_1 \xi_{kQ_1^c})$ are graded Hopf algebra isomorphisms from $T_{(kG)^*}^c(kQ_1^a)$ to $T_{(kG)^*}^c((kQ_1^c)^*)$ and from $T_{kG}(kQ_1^c)$ to $T_{kG}((kQ_1^a)^*)$, respectively. In this case, (kQ_1^a, kQ_1^c) , (kQ^a, kQ^c) and (kQ^s, kQ^{sc}) are said to be arrow dual pairings.

If $(kQ_1^c, \alpha^-, \alpha^+, \delta^-, \delta^+)$ is a PM kG -Hopf bimodule, and (kQ_1^c, kQ_1^a) is an arrow pairing, then $(kQ_1^a, \alpha^{*-}, \alpha^{*+})$ is called a PM $(kG)^*$ -Hopf bimodule and six quiver Hopf algebras induced by kQ_1^c and kQ_1^a are called PM quiver Hopf algebras.

Now we are going to describe the structure of all PM kG -Hopf bimodules and the corresponding graded Hopf algebras.

Definition 14.1.10. $(G, r, \vec{\chi}, u)$ is called a ramification system with characters (or RSC in short), if r is a ramification of G , u is a map from $\mathcal{K}(G)$ to G with $u(C) \in C$ for any

$C \in \mathcal{K}(G)$, and $\vec{\chi} = \{\chi_C^{(i)}\}_{i \in I_C(r), C \in \mathcal{K}_r(G)} \in \prod_{C \in \mathcal{K}_r(G)} (\widehat{Z_{u(C)}})^{r_C}$ with $\chi_C^{(i)} \in \widehat{Z_{u(C)}}$ for any $i \in I_C(r), C \in \mathcal{K}_r(G)$.

$\text{RSC}(G, r, \vec{\chi}, u)$ and $\text{RSC}(G', r', \vec{\chi}', u')$ are said to be isomorphic if the following conditions are satisfied:

- There exists a group isomorphism $\phi : G \rightarrow G'$.
- For any $C \in \mathcal{K}(G)$, there exists an element $h_C \in G$ such that $\phi(h_C^{-1}u(C)h_C) = u'(\phi(C))$.
- For any $C \in \mathcal{K}_r(G)$, there exists a bijective map $\phi_C : I_C(r) \rightarrow I_{\phi(C)}(r')$ such that $\chi'_{\phi(C)}^{(\phi_C(i))}(\phi(h_C^{-1}hh_C)) = \chi_C^{(i)}(h)$ for all $h \in Z_{u(C)}$ and $i \in I_C(r)$.

Remark. Assume that $G = G'$, $r = r'$ and $u(C) = u'(C)$ for any $C \in \mathcal{K}_r(G)$. If there is a permutation ϕ_C on $I_C(r)$ for any $C \in \mathcal{K}_r(G)$ such that $\chi'_{\phi(C)}^{(\phi_C(i))} = \chi_C^{(i)}$ for all $i \in I_C(r)$, then obviously $\text{RSC}(G, r, \vec{\chi}, u) \cong \text{RSC}(G, r, \vec{\chi}', u)$.

Proposition 14.1.11. *If N is a PM kG -Hopf bimodule, then there exist a Hopf quiver (Q, G, r) , an $\text{RSC}(G, r, \vec{\chi}, u)$ and a kG -Hopf bimodule $(kQ_1^c, \alpha^-, \alpha^+)$ with*

$$\alpha^-(h \otimes a_{y,x}^{(i)}) := h \cdot a_{y,x}^{(i)} = a_{hy,hx}^{(i)}, \quad \alpha^+(a_{y,x}^{(i)} \otimes h) := a_{y,x}^{(i)} \cdot h = \chi_C^{(i)}(\zeta_\theta(h))a_{yh,xh}^{(i)}$$

where $x, y, h \in G$ with $x^{-1}y = g_\theta^{-1}u(C)g_\theta$, ζ_θ is given by (14.3), $C \in \mathcal{K}_r(G)$ and $i \in I_C(r)$, such that $N \cong (kQ_1^c, \alpha^-, \alpha^+)$ as kG -Hopf bimodules.

Proof. Since N is a PM kG -Hopf bimodule, there exists an object $\prod_{C \in \mathcal{K}(G)} M(C)$ in $\prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$ such that $M(C)$ is a pointed $kZ_{u(C)}$ -module for any $C \in \mathcal{K}(G)$ and $N \cong V(\{M(C)\}_{C \in \mathcal{K}(G)}) = \bigoplus_{y=g_\theta^{-1}u(C)g_\theta, x,y \in G} x \otimes M(C) \otimes_{kZ_{u(C)}} g_\theta$ as kG -Hopf bimodules. Let $r = \sum_{C \in \mathcal{K}(G)} r_C C$ with $r_C = \dim M(C)$ for any $C \in \mathcal{K}(G)$. Notice that $\dim M(C)$ denotes the cardinal number of a basis of $M(C)$ when $M(C)$ is infinite dimensional. Since $(M(C), \alpha_C)$ is a pointed $kZ_{u(C)}$ -module, there exist a k -basis $\{x_C^{(i)} \mid i \in I_C(r)\}$ in $M(C)$ and a family of characters $\{\chi_C^{(i)} \in \widehat{Z_{u(C)}} \mid i \in I_C(r)\}$ such that $\alpha_C(x_C^{(i)} \otimes h) = x_C^{(i)} \triangleleft h = \chi_C^{(i)}(h)x_C^{(i)}$ for any $i \in I_C(r)$ and $h \in Z_{u(C)}$.

We have to show that $(kQ_1^c, \alpha^-, \alpha^+)$ is isomorphic to $\bigoplus_{y=g_\theta^{-1}u(C)g_\theta, x,y \in G} x \otimes M(C) \otimes_{kZ_{u(C)}} g_\theta$ as kG -Hopf bimodules. Observe that there is a canonical kG -bicomodule isomorphism $\varphi : kQ_1 \rightarrow \bigoplus_{y=g_\theta^{-1}u(C)g_\theta, x,y \in G} x \otimes M(C) \otimes_{kZ_{u(C)}} g_\theta$ given by

$$\varphi(a_{y,x}^{(i)}) = x \otimes x_C^{(i)} \otimes_{kZ_{u(C)}} g_\theta \tag{14.9}$$

where $x, y \in G$ with $x^{-1}y = g_\theta^{-1}u(C)g_\theta$, $C \in \mathcal{K}_r(G)$ and $i \in I_C(r)$. Now we have

$$\begin{aligned} \varphi(\alpha^-(h \otimes a_{y,x}^{(i)})) &= \varphi(a_{hy,hx}^{(i)}) = hx \otimes x_C^{(i)} \otimes_{kZ_{u(C)}} g_\theta \\ &= h \cdot (x \otimes x_C^{(i)} \otimes_{kZ_{u(C)}} g_\theta) \quad (\text{see (14.6)}) \\ &= h \cdot \varphi(a_{y,x}^{(i)}) \end{aligned}$$

and

$$\begin{aligned}
\varphi(\alpha^+(a_{y,x}^{(i)} \otimes h)) &= \chi_C^{(i)}(\zeta_\theta(h))\varphi(a_{yh,xh}^{(i)}) = \chi_C^{(i)}(\zeta_\theta(h))xh \otimes x_C^{(i)} \otimes_{kZ_{u(C)}} g_{\theta'} \\
&\quad (\text{since } hg_{\theta'}^{-1} = g_\theta^{-1}\zeta_\theta(h) \text{ and } yh = xhg_{\theta'}^{-1}u(C)g_{\theta'}) \\
&= (x \otimes x_C^{(i)} \otimes_{kZ_{u(C)}} g_\theta) \cdot h \quad (\text{see (14.6)}) \\
&= \varphi(a_{y,x}^{(i)}) \cdot h,
\end{aligned}$$

where $x, y, h \in G$ with $x^{-1}y = g_\theta^{-1}u(C)g_\theta$, ζ_θ is given by (14.3), $C \in \mathcal{K}_r(G)$ and $i \in I_C(r)$. Consequently, φ is a kG -Hopf bimodule isomorphism. $\square$

Let $(kQ_1^c, G, r, \vec{\chi}, u)$ denote the kG -Hopf bimodule $(kQ_1^c, \alpha^-, \alpha^+)$ given in Lemma 14.1.11. Furthermore, if (kQ_1^c, kQ_1^a) is an arrow dual pairing, then we denote the $(kG)^*$ -Hopf bimodule kQ_1^a by $(kQ_1^a, G, r, \vec{\chi}, u)$. We obtain six quiver Hopf algebras $kQ^c(G, r, \vec{\chi}, u)$, $kQ^s(G, r, \vec{\chi}, u)$, $kG[kQ_1^c, G, r, \vec{\chi}, u]$, $kQ^a(G, r, \vec{\chi}, u)$, $kQ^{sc}(G, r, \vec{\chi}, u)$, $(kG)^*[kQ_1^a, G, r, \vec{\chi}, u]$, called the quiver Hopf algebras determined by $\text{RSC}(G, r, \vec{\chi}, u)$.

From Proposition 14.1.11, it seems that the right kG -action on $(kQ_1^c, G, r, \vec{\chi}, u)$ depends on the choice of the set $\{g_\theta \mid \theta \in \Theta_C\}$ of coset representatives of $Z_{u(C)}$ in G (see, Eq.(14.1)). The following lemma shows that $(kQ_1^c, G, r, \vec{\chi}, u)$ is, in fact, independent of the choice of the coset representative set $\{g_\theta \mid \theta \in \Theta_C\}$, up to kG -Hopf bimodule isomorphisms. For a while, we write $(kQ_1^c, G, r, \vec{\chi}, u) = (kQ_1^c, G, r, \vec{\chi}, u, \{g_\theta\})$ given before. Now let $\{h_\theta \in G \mid \theta \in \Theta_C\}$ be another coset representative set of $Z_{u(C)}$ in G for any $C \in \mathcal{K}(G)$. That is,

$$G = \bigcup_{\theta \in \Theta_C} Z_{u(C)}h_\theta. \quad (14.10)$$

Lemma 14.1.12. *With the above notations, $(kQ_1^c, G, r, \vec{\chi}, u, \{g_\theta\})$ and $(kQ_1^c, G, r, \vec{\chi}, u, \{h_\theta\})$ are isomorphic kG -Hopf bimodules.*

Proof. We may assume $Z_{u(C)}h_\theta = Z_{u(C)}g_\theta$ for any $C \in \mathcal{K}(G)$ and $\theta \in \Theta_C$. Then $g_\theta h_\theta^{-1} \in Z_{u(C)}$. Now let $x, y, h \in G$ with $x^{-1}y = g_\theta^{-1}u(C)g_\theta$. Then $x^{-1}y = h_\theta^{-1}(g_\theta h_\theta^{-1})^{-1}u(C)(g_\theta h_\theta^{-1})h_\theta = h_\theta^{-1}u(C)h_\theta$ and $h_\theta h = (h_\theta g_\theta^{-1})g_\theta h = (h_\theta g_\theta^{-1})\zeta_\theta(h)g_{\theta'} = (h_\theta g_\theta^{-1})\zeta_\theta(h)(g_{\theta'}h_{\theta'}^{-1})h_{\theta'}$, where $g_\theta h = \zeta_\theta(h)g_{\theta'}$. Hence from Proposition 14.1.11 we know that the right kG -action on $(kQ_1^c, G, r, \vec{\chi}, u, \{h_\theta\})$ is given by

$$\begin{aligned}
a_{y,x}^{(i)} \cdot h &= \chi_C^{(i)}((h_\theta g_\theta^{-1})\zeta_\theta(h)(g_{\theta'}h_{\theta'}^{-1}))a_{yh,xh}^{(i)} \\
&= \chi_C^{(i)}((g_\theta h_\theta^{-1})^{-1})\chi_C^{(i)}(\zeta_\theta(h))\chi_C^{(i)}(g_{\theta'}h_{\theta'}^{-1})a_{yh,xh}^{(i)}, \quad i \in I_C(r).
\end{aligned}$$

However, we also have

$$(xh)^{-1}(yh) = h^{-1}(x^{-1}y)h = h^{-1}g_\theta^{-1}u(C)g_\theta h = g_{\theta'}^{-1}u(C)g_{\theta'}.$$

It follows that the k -linear isomorphism $f : kQ_1 \rightarrow kQ_1$ given by

$$f(a_{y,x}^{(i)}) = \chi_C^{(i)}(g_\theta h_\theta^{-1})a_{y,x}^{(i)}$$

for any $x, y \in G$ with $x^{-1}y = g_\theta^{-1}u(C)g_\theta$, $C \in \mathcal{K}_r(G)$ and $i \in I_C(r)$, is a kG -Hopf bimodule isomorphism from $(kQ_1^c, G, r, \vec{\chi}, u, \{g_\theta\})$ to $(kQ_1^c, G, r, \vec{\chi}, u, \{h_\theta\})$. $\square$

Now we state one of our main results, which classifies the PM (co-)path Hopf algebras, PM semi-(co-)path Hopf algebras and PM one-type-path Hopf algebras.

Theorem 14.1.13. *Let $(G, r, \vec{\chi}, u)$ and $(G', r', \vec{\chi}', u')$ are two RSC's. Then the following statements are equivalent:*

- (i) $\text{RSC}(G, r, \vec{\chi}, u) \cong \text{RSC}(G', r', \vec{\chi}', u')$.
- (ii) *There exists a Hopf algebra isomorphism $\phi : kG \rightarrow kG'$ such that $(kQ_1^c, G, r, \vec{\chi}, u) \cong_{\phi}^{\phi^{-1}} ((kQ_1'^c, G', r', \vec{\chi}', u'))_{\phi}^{\phi^{-1}}$ as kG -Hopf bimodules.*
- (iii) $kQ^c(G, r, \vec{\chi}, u) \cong kQ'^c(G, r, \vec{\chi}, u)$. (iv) $kQ^s(G, r, \vec{\chi}, u) \cong kQ'^s(G', r', \vec{\chi}', u')$.
- (v) $kG[kQ_1^c, G, r, \vec{\chi}, u] \cong kG'[kQ_1'^c, G', r', \vec{\chi}', u']$.

Furthermore, if Q is finite, then the above are equivalent to the following:

- (vi) $kQ^a(G, r, \vec{\chi}, u) \cong kQ'^a(G', r', \vec{\chi}', u')$. (vii) $kQ^{sc}(G, r, \vec{\chi}, u) \cong kQ'^{sc}(G', r', \vec{\chi}', u')$.
- (viii) $(kG)^*[kQ_1^a, G, r, \vec{\chi}, u] \cong (kG')^*[kQ_1'^a, G', r', \vec{\chi}', u']$. Notice that the isomorphisms above are ones of graded Hopf algebras but (i) (ii).

Proof. By Lemma 14.1.7 and Lemma 14.1.8, we only have to prove (i) $\Leftrightarrow$ (ii).

(i) $\Rightarrow$ (ii). Assume that $\text{RSC}(G, r, \vec{\chi}, u) \cong \text{RSC}(G', r', \vec{\chi}', u')$. Then there exist a group isomorphism $\phi : G \rightarrow G'$, an element $h_C \in G$ such that $\phi(h_C^{-1}u(C)h_C) = u'(\phi(C))$ for any $C \in \mathcal{K}(G)$ and a bijective map $\phi_C : I_C(r) \rightarrow I_{\phi(C)}(r')$ such that $\chi'_{\phi(C)}(\phi(h_C^{-1}hh_C)) = \chi_C^{(i)}(h)$ for any $h \in Z_{u(C)}$, $C \in \mathcal{K}_r(G)$ and $i \in I_C(r)$. Then $\phi(h_C^{-1}Z_{u(C)}h_C) = Z_{u'(\phi(C))}$ and $\phi : kG \rightarrow kG'$ is a Hopf algebra isomorphism. Now let $G = \bigcup_{\theta \in \Theta_C} Z_{u(C)}g_{\theta}$ be given as in (14.2) for any $C \in \mathcal{K}(G)$, and assume that the kG -Hopf bimodule $(kQ_1^c, G, r, \vec{\chi}, u)$ is defined by using these coset decompositions. Then

$$G' = \bigcup_{\theta \in \Theta_C} Z_{u'(\phi(C))}(\phi(h_C^{-1}g_{\theta}h_C)) \quad (14.11)$$

is a coset decomposition of $Z_{u'(\phi(C))}$ in G' for any $\phi(C) \in \mathcal{K}(G')$. By Lemma 14.1.12, we may assume that the structure of the kG' -Hopf bimodule $(kQ_1'^c, G', r', \vec{\chi}', u')$ is obtained by using these coset decompositions (14.11). Define a k -linear isomorphism $\psi : (kQ_1^c, G, r, \vec{\chi}, u) \rightarrow (kQ_1'^c, G', r', \vec{\chi}', u')$ by

$$\psi(a_{y,x}^{(i)}) = \chi_C^{(i)}(\zeta_{\theta}(h_C^{-1}))a_{\phi(y),\phi(x)}^{(\phi_C(i))}$$

for any $x, y \in G$ with $x^{-1}y = g_{\theta}^{-1}u(C)g_{\theta}$, and $i \in I_C(r)$, where $C \in \mathcal{K}_r(G)$ and $g_{\theta}h_C^{-1} = \zeta_{\theta}(h_C^{-1})g_{\eta}$ with $\zeta_{\theta}(h_C^{-1}) \in Z_{u(C)}$ and $\theta, \eta \in \Theta_C$. It is easy to see that ψ is a kG -bicomodule homomorphism from $(kQ_1^c, G, r, \vec{\chi}, u)$ to $_{\phi}^{\phi^{-1}}(kQ_1'^c, G', r', \vec{\chi}', u')_{\phi}^{\phi^{-1}}$. Since $(hx)^{-1}(hy) = x^{-1}y$ for any $x, y, h \in G$, it follows from Proposition 14.1.11 that ψ is also a left kG -module homomorphism from $(kQ_1^c, G, r, \vec{\chi}, u)$ to $_{\phi}^{\phi^{-1}}(kQ_1'^c, G', r', \vec{\chi}', u')_{\phi}^{\phi^{-1}}$.

Now let $x, y, h \in G$ with $x^{-1}y = g_{\theta}^{-1}u(C)g_{\theta}$, $C \in \mathcal{K}_r(G)$ and $\theta \in \Theta_C$. Assume that $g_{\theta}h_C^{-1} = \zeta_{\theta}(h_C^{-1})g_{\eta}$, $g_{\theta}h = \zeta_{\theta}(h)g_{\eta'}$, $g_{\eta}(h_C h h_C^{-1}) = \zeta_{\eta}(h_C h h_C^{-1})g_{\eta'}$ and $g_{\theta'}h_C^{-1} = \zeta_{\theta'}(h_C^{-1})g_{\theta''}$ with $\zeta_{\theta}(h_C^{-1}), \zeta_{\theta}(h), \zeta_{\eta}(h_C h h_C^{-1}), \zeta_{\theta'}(h_C^{-1}) \in Z_{u(C)}$ and $\eta, \theta', \eta', \theta'' \in \Theta_C$. Then we have

$$g_{\theta}h h_C^{-1} = \zeta_{\theta}(h)g_{\theta'}h_C^{-1} = \zeta_{\theta}(h)\zeta_{\theta'}(h_C^{-1})g_{\theta''}$$

and

$$g_{\theta}h h_C^{-1} = (g_{\theta}h_C^{-1})(h_C h h_C^{-1}) = \zeta_{\theta}(h_C^{-1})g_{\eta}(h_C h h_C^{-1}) = \zeta_{\theta}(h_C^{-1})\zeta_{\eta}(h_C h h_C^{-1})g_{\eta'}.$$

It follows that $\theta'' = \eta'$ and

$$\zeta_\theta(h)\zeta_{\theta'}(h_C^{-1}) = \zeta_\theta(h_C^{-1})\zeta_\eta(h_C h h_C^{-1}). \quad (14.12)$$

By Proposition 14.1.11, $a_{y,x}^{(i)} \cdot h = \chi_C^{(i)}(\zeta_\theta(h))a_{yh,xh}^{(i)}$ for any $i \in I_C(r)$. Moreover, we have $(xh)^{-1}(yh) = h^{-1}g_\theta^{-1}u(C)g_\theta h = g_{\theta'}^{-1}u(C)g_{\theta'}$. This implies that

$$\psi(a_{y,x}^{(i)} \cdot h) = \chi_C^{(i)}(\zeta_\theta(h))\psi(a_{yh,xh}^{(i)}) = \chi_C^{(i)}(\zeta_\theta(h))\chi_C^{(i)}(\zeta_{\theta'}(h_C^{-1}))a_{\phi(yh),\phi(xh)}^{(\phi_C(i))}.$$

On the other hand, we have $g_\theta = g_\theta h_C^{-1}h_C = \zeta_\theta(h_C^{-1})g_\eta h_C$, and hence

$$\begin{aligned} \phi(x)^{-1}\phi(y) &= \phi(x^{-1}y) = \phi(g_\theta^{-1}u(C)g_\theta) = \phi(h_C^{-1}g_\eta^{-1}u(C)g_\eta h_C) \\ &= \phi(h_C^{-1}g_\eta^{-1}h_C)\phi(h_C^{-1}u(C)h_C)\phi(h_C^{-1}g_\eta h_C) \\ &= \phi(h_C^{-1}g_\eta h_C)^{-1}u'(\phi(C))\phi(h_C^{-1}g_\eta h_C). \end{aligned}$$

We also have

$$\begin{aligned} \phi(h_C^{-1}g_\eta h_C)\phi(h) &= \phi(h_C^{-1}g_\eta h_C h h_C^{-1}h_C) \\ &= \phi(h_C^{-1}\zeta_\eta(h_C h h_C^{-1})g_{\eta'} h_C) \\ &= \phi(h_C^{-1}\zeta_\eta(h_C h h_C^{-1})h_C)\phi(h_C^{-1}g_{\eta'} h_C). \end{aligned}$$

Thus by Proposition 14.1.11 one gets

$$\begin{aligned} a_{\phi(y),\phi(x)}^{(\phi_C(i))} \cdot \phi(h) &= \chi_{\phi(C)}^{(\phi_C(i))}(\phi(h_C^{-1}\zeta_\eta(h_C h h_C^{-1})h_C))a_{\phi(yh),\phi(xh)}^{(\phi_C(i))} \\ &= \chi_C^{(i)}(\zeta_\eta(h_C h h_C^{-1}))a_{\phi(yh),\phi(xh)}^{(\phi_C(i))}. \end{aligned}$$

Now it follows from Eq.(14.12) that

$$\begin{aligned} \psi(a_{y,x}^{(i)} \cdot \phi(h)) &= \chi_C^{(i)}(\zeta_\theta(h_C^{-1}))a_{\phi(y),\phi(x)}^{(\phi_C(i))} \cdot \phi(h) \\ &= \chi_C^{(i)}(\zeta_\theta(h_C^{-1}))\chi_C^{(i)}(\zeta_\eta(h_C h h_C^{-1}))a_{\phi(yh),\phi(xh)}^{(\phi_C(i))} \\ &= \psi(a_{y,x}^{(i)} \cdot h). \end{aligned}$$

This shows that ϕ is a right kG -module homomorphism, and hence a kG -Hopf bimodule isomorphism from $(kQ_1^c, G, r, \vec{\chi}, u)$ to $\phi^{-1}(kQ_1'^c, G', r', \vec{\chi}', u')\phi^{-1}$.

(ii) $\Rightarrow$ (i). Assume that there exist a Hopf algebra isomorphism $\phi : kG \rightarrow kG'$ and a kG -Hopf bimodule isomorphism $\psi : (kQ_1^c, G, r, \vec{\chi}, u) \rightarrow \phi^{-1}(kQ_1'^c, G', r', \vec{\chi}', u')\phi^{-1}$. Then $\phi : G \rightarrow G'$ is a group isomorphism. Let $C \in \mathcal{K}(G)$. Then $\phi(u(C)), u'(\phi(C)) \in \phi(C) \in \mathcal{K}(G')$, and hence $u'(\phi(C)) = \phi(h_C)^{-1}\phi(u(C))\phi(h_C) = \phi(h_C^{-1}u(C)h_C)$ for some $h_C \in G$. Since ψ is a kG' -bicomodule isomorphism from $\phi(kQ_1^c, G, r, \vec{\chi}, u)\phi$ to $(kQ_1'^c, G', r', \vec{\chi}', u')$ and $\phi(h_C^{-1}u(C)h_C) = u'(\phi(C))$, by restriction one gets a k -linear isomorphism

$$\psi_C : h_C^{-1}u(C)h_C(kQ_1)^1 \rightarrow u'(\phi(C))(kQ_1')^1, \quad x \mapsto \psi(x).$$

We also have a k -linear isomorphism

$$f_C : u(C)(kQ_1)^1 \rightarrow h_C^{-1}u(C)h_C(kQ_1)^1, \quad x \mapsto h_C^{-1} \cdot x \cdot h_C.$$

Since $\phi(h_C^{-1}u(C)h_C) = u'(\phi(C))$ and $h_C^{-1}Z_{u(C)}h_C = Z_{h_C^{-1}u(C)h_C}$, one gets $\phi(h_C^{-1}Z_{u(C)}h_C) = Z_{u'(\phi(C))}$. Hence ϕ and h_C induce an algebra isomorphism

$$\sigma_C : kZ_{u(C)} \rightarrow kZ_{u'(\phi(C))}, \quad h \mapsto \phi(h_C^{-1}hh_C).$$

Using the hypothesis that ψ is a kG -bimodules homomorphism from $(kQ_1^c, G, r, \vec{\chi}, u)$ to ${}_{\phi}(kQ_1'^c, G', r', \vec{\chi}', u')_{\phi}$, one can easily check that the composition $\psi_C f_C$ is a right $kZ_{u(C)}$ -module isomorphism from ${}^{u(C)}(kQ_1)^1$ to $({}^{u'(\phi(C))}(kQ_1')^1)_{\sigma_C}$. Since both ${}^{u(C)}(kQ_1)^1$ and $({}^{u'(\phi(C))}(kQ_1')^1)_{\sigma_C}$ are pointed right $kZ_{u(C)}$ -modules, they are semisimple $kZ_{u(C)}$ -modules for any $C \in \mathcal{K}_r(G)$. Moreover, $ka_{u(C),1}^{(i)}$ and $ka_{u'(\phi(C)),1}^{(j)}$ are simple submodules of ${}^{u(C)}(kQ_1)^1$ and $({}^{u'(\phi(C))}(kQ_1')^1)_{\sigma_C}$, respectively, for any $i \in I_C(r)$ and $j \in I_{\phi(C)}(r')$, where $C \in \mathcal{K}_r(G)$. Thus for any $C \in \mathcal{K}_r(G)$, there exists a bijective map $\phi_C : I_C(r) \rightarrow I_{\phi(C)}(r')$ such that $ka_{u(C),1}^{(i)}$ and $(ka_{u'(\phi(C)),1}^{(\phi_C(i))})_{\sigma_C}$ are isomorphic right $kZ_{u(C)}$ -modules for any $i \in I_C(r)$, which implies $\chi_C^{(i)}(h) = \chi'_{\phi(C)}^{(\phi_C(i))}(\phi(h_C^{-1}hh_C))$ for any $h \in Z_{u(C)}$ and $i \in I_C(r)$. It follows that $\text{RSC}(G, r, \vec{\chi}, u) \cong \text{RSC}(G', r', \vec{\chi}', u')$. $\square$

Up to now we have classified the PM quiver Hopf algebras by means of RSC's. In other words, ramification systems with characters uniquely determine their corresponding PM quiver Hopf algebras up to graded Hopf algebra isomorphisms.

Example 14.1.14. Assume that k is a field with $\text{char}(k) \neq 2$. Let $G = \{1, g\} \cong \mathbf{Z}_2$ be the cyclic group of order 2 with the generator g . Let r be a ramification data of G with $r_1 = m$ and $r_{\{g\}} = 0$ and (Q, G, r) be the corresponding Hopf quiver, where m is a positive integer. Then ${}^1(Q_1)^1 = \{a_{1,1}^{(i)} \mid i = 1, 2, \dots, m\}$, ${}^g(Q_1)^g = \{a_{g,g}^{(i)} \mid i = 1, 2, \dots, m\}$, ${}^1(Q_1)^g$ and ${}^g(Q_1)^1$ are two empty sets. For simplification, we write $x_i = a_{1,1}^{(i)}$ and $y_i = a_{g,g}^{(i)}$ for any $1 \leq i \leq m$. Clearly, $Z_{u(\{1\})} = G$ and $\hat{G} = \{\chi_+, \chi_-\}$, where $\chi_{\pm}(g) = \pm 1$. For any $0 \leq n \leq m$, put $\vec{\chi}_n \in (\hat{G})^m$ with $\chi_{n\{1\}}^{(i)} = \begin{cases} \chi_-, & \text{if } i > n; \\ \chi_+, & \text{otherwise.} \end{cases}$ Then $\{\text{RSC}(G, r, \vec{\chi}_n, u_n) \mid n = 0, 1, 2, \dots, m\}$ are all non-isomorphic RSC's. Thus by Theorem 14.1.13 we know that the path coalgebra kQ^c exactly admits $m+1$ distinct PM co-path Hopf algebra structures $kQ^c(G, r, \vec{\chi}_n, u_n)$, $0 \leq n \leq m$, up to graded Hopf algebra isomorphism. Now let $0 \leq n \leq m$. Then by Proposition 14.1.11, the kG -actions on $(kQ_1^c, G, r, \vec{\chi}_n, u_n)$ are given by $g \cdot x_i = y_i$, $g \cdot y_i = x_i$, $1 \leq i \leq m$; $x_i \cdot g = \begin{cases} -y_i, & \text{if } i > n, \\ y_i, & \text{otherwise,} \end{cases}$ $y_i \cdot g = \begin{cases} -x_i, & \text{if } i > n, \\ x_i, & \text{otherwise.} \end{cases}$ Thus by [36, p.245 or Theorem 3.8], the products of these arrows x_i, y_j in $kQ^c(G, r, \vec{\chi}_n, u_n)$ can be described as follows. For any $i, j = 1, 2, \dots, m$, $x_i \cdot x_j = x_i x_j + x_j x_i$, $y_i \cdot x_j = y_i y_j + y_j y_i$, $x_i \cdot y_j = \begin{cases} -(y_i y_j + y_j y_i) & , \text{ if } i > n, \\ y_i y_j + y_j y_i & , \text{ otherwise,} \end{cases}$ $y_i \cdot y_j = \begin{cases} -(x_i x_j + x_j x_i) & , \text{ if } i > n, \\ x_i x_j + x_j x_i & , \text{ otherwise,} \end{cases}$ where $x \cdot y$ denotes the product of x and y in $kQ^c(G, r, \vec{\chi}_n, u_n)$ for any $x, y \in kQ^c(G, r, \vec{\chi}_n, u_n)$, $x_i x_j$ and $y_i y_j$ denote the 2-paths in the quiver Q as usual for any $1 \leq i, j \leq m$.

14.2 Multiple Taft algebras

In this section we discuss the PM quiver Hopf algebras determined by the RSC's with $\cup \mathcal{K}_r(G) \subseteq Z(G)$. We give the classification of PM Nichols algebras and multiple Taft algebras by means of element system with characters when G is finite abelian group and k is the complex field.

Let r be a ramification data of G and (Q, G, r) be the corresponding Hopf quiver. If C contains only one element of G for any $C \in \mathcal{K}_r(G)$, then $C = \{g\}$ for some $g \in Z(G)$, the center of

G . In this case, we say that the ramification r is *central*, and that $\text{RSC}(G, r, \overrightarrow{\chi}, u)$ a *central ramification system with characters*, or a CRSC in short. If $\text{RSC}(G, r, \overrightarrow{\chi}, u)$ is CRSC, then the PM co-path Hopf algebra $kQ^c(G, r, \overrightarrow{\chi}, u)$ is called a *multiple crown algebra* and $kG[kQ_1^c, G, r, \overrightarrow{\chi}, u]$ is called a *multiple Taft algebra*.

Definition 14.2.1. $(G, \overrightarrow{g}, \overrightarrow{\chi}, J)$ is called an *element system with characters* (simply, ESC) if G is a group, J is a set, $\overrightarrow{g} = \{g_i\}_{i \in J} \in Z(G)^J$ and $\overrightarrow{\chi} = \{\chi_i\}_{i \in J} \in \widehat{G}^J$ with $g_i \in Z(G)$ and $\chi_i \in \widehat{G}$. $\text{ESC}(G, \overrightarrow{g}, \overrightarrow{\chi}, J)$ and $\text{ESC}(G', \overrightarrow{g}', \overrightarrow{\chi}', J')$ are said to be isomorphic if there exist a group isomorphism $\phi : G \rightarrow G'$ and a bijective map $\sigma : J \rightarrow J'$ such that $\phi(g_i) = g'_{\sigma(i)}$ and $\chi'_{\sigma(i)}\phi = \chi_i$ for any $i \in J$.

$\text{ESC}(G, \overrightarrow{g}, \overrightarrow{\chi}, J)$ can be written as $\text{ESC}(G, g_i, \chi_i; i \in J)$ for convenience. Throughout this chapter, let $q_{ji} := \chi_i(g_j)$, $q_i = q_{ii}$ and N_i be the order of q_i ($N_i = \infty$ when q_i is not a root of unit, or $q_i = 1$) for $i, j \in J$.

Let H be a Hopf algebra with a bijective antipode S . A YD H -module V is *pointed* if $V = 0$ or V is a direct sum of one dimensional YD H -modules. If V is a pointed YD H -module, then the corresponding Nichols algebra $\mathcal{B}(V)$ is called a PM *Nichols algebra*.

Lemma 14.2.2. Let (V, α^-, δ^-) be a YD kG -module. Then (V, α^-, δ^-) is a pointed YD kG -module if and only if (V, α^-) is a pointed (left) kG -module with $\delta^-(V) \subseteq kZ(G) \otimes V$.

Proof. We may assume $V \neq 0$. Assume that V is a pointed YD kG -module. Then V is obviously a pointed kG -module. Now let U be a one dimensional YD kG -submodule of V and let $0 \neq x \in U$. Then $\delta^-(x) = g \otimes x$ for some $g \in G$. Let $h \in G$. Then $h \cdot x = \beta x$ for some $0 \neq \beta \in k$. By Eq.(14.4), one gets that $\beta g \otimes x = \delta^-(\beta x) = \delta^-(h \cdot x) = hgh^{-1} \otimes h \cdot x = \beta hgh^{-1} \otimes x$. This implies that $hgh^{-1} = g$ for any $h \in G$, and hence $g \in Z(G)$. It follows that $\delta^-(V) \subseteq kZ(G) \otimes V$. Conversely, assume that V is a pointed kG -module and $\delta^-(V) \subseteq kZ(G) \otimes V$. For any $g \in G$, let

$$V_g = \{v \in V \mid \delta^-(v) = g \otimes v\}.$$

Then V_g is a kG -subcomodule of V and any subspace of V_g is a kG -subcomodule for all $g \in G$. By $\delta^-(V) \subseteq kZ(G) \otimes V$, we know that $V = \bigoplus_{g \in Z(G)} V_g$. Now let $g \in Z(G)$. Then it follows from Eq.(14.4) that V_g is a kG -submodule of V . Since V is a pointed kG -module, any submodule of V is pointed, and hence V_g is a pointed kG -module. Thus if $V_g \neq 0$ then V_g is a direct sum of some one dimensional kG -submodules of V_g . However, any one dimensional kG -submodule of V_g is also a kG -subcomodule, and hence a YD kG -submodule. Thus V is a pointed YD kG -module. $\square$

Lemma 14.2.3. Assume that G is a finite abelian group of exponent m . If k contains a primitive m -th root of 1, then every YD kG -module is pointed and Nichols algebra of every YD kG -module is PM.

Proof. It follows from Lemma 14.1.3 and Lemma 14.2.2. $\square$

Let $(G, g_i, \chi_i; i \in J)$ be an ESC. Let V be a k -vector space with $\dim(V) = |J|$. Let $\{x_i \mid i \in J\}$ be a basis of V over k . Define a left kG -action and a left kG -coaction on V by

$$g \cdot x_i = \chi_i(g)x_i, \quad \delta^-(x_i) = g_i \otimes x_i, \quad i \in J, \quad g \in G.$$

Then it is easy to see that V is a pointed YD kG -module and kx_i is a one dimensional YD kG -submodule of V for any $i \in J$. Denote by $V(G, g_i, \chi_i; i \in J)$ the pointed YD kG -module V . Note that $V(G, g_i, \chi_i; i \in J) = 0$ if J is empty.

Proposition 14.2.4. *V is pointed YD kG -module if and only if V is isomorphic to $V(G, g_i, \chi_i; i \in J)$ for some ESC $(G, g_i, \chi_i; i \in J)$.*

Proof. If $V \cong V(G, g_i, \chi_i; i \in J)$ for some ESC $(G, g_i, \chi_i; i \in J)$ of G , then V is obviously a pointed YD kG -module. Conversely, assume that V is a nonzero pointed YD kG -module. By Lemma 14.2.2, $V = \bigoplus_{g \in Z(G)} V_g$ and $V_g = \{v \in V \mid \delta^-(v) = g \otimes v\}$ is a pointed YD kG -submodule of V for any $g \in Z(G)$. Let $g \in Z(G)$ with $V_g \neq 0$. Then V_g is a nonzero pointed kG -module. Hence there is a k -basis $\{x_i \mid i \in J_g\}$ such that kx_i is a kG -submodule of V_g for any $i \in J_g$. It follows that there is a character $\chi_i \in \hat{G}$ for any $i \in J_g$ such that $h \cdot x_i = \chi_i(h)x_i$ for all $h \in G$. For any $i \in J_g$, put $g_i = g$. We may assume that these index sets J_g are disjoint, that is, $J_g \cap J_h = \emptyset$ for any $g \neq h$ in $Z(G)$ with $V_g \neq 0$ and $V_h \neq 0$. Now let J be the union of all the J_g with $g \in Z(G)$ and $V_g \neq 0$. Then one can see that $(G, g_i, \chi_i; i \in J)$ is an ESC of G , and that V is isomorphic to $V(G, g_i, \chi_i; i \in J)$ as a YD kG -module. $\square$

Now we give the relation between RSC and ESC. Assume that $(G, g_i, \chi_i; i \in J)$ is an ESC of G . We define a binary relation $\sim$ on J by

$$i \sim j \Leftrightarrow g_i = g_j,$$

where $i, j \in J$. Clearly, this is an equivalence relation. Denote by $J/\sim$ the quotient set of J modulo $\sim$. For any $i \in J$, let $[i]$ denote the equivalence class containing i . That is, $[i] := \{j \in J \mid j \sim i\}$. Choose a subset $\bar{J} \subseteq J$ such that the assignment $i \mapsto [i]$ is a bijective map from $\bar{J}$ to $J/\sim$. That is, $\bar{J}$ is a set of representative elements of the equivalence class. Then $J = \bigcup_{i \in \bar{J}} [i]$ is a disjoint union. Let $r_{\{g_i\}} = |[i]|$ for any $i \in \bar{J}$. Then $r = \sum_{i \in \bar{J}} r_{\{g_i\}} \{g_i\}$ is a central ramification of G with $I_{\{g_i\}}(r) = [i]$ for $i \in \bar{J}$. Moreover, $\mathcal{K}_r(G) = \{\{g_i\} \mid i \in \bar{J}\}$. Put $\chi_{\{g_i\}}^{(j)} := \chi_j$ for any $i \in \bar{J}$ and $j \in [i]$. We obtain an CRSC, written $\text{CRSC}(G, r(g_i, \chi_i; i \in J), \vec{\chi}(g_i, \chi_i; i \in J), u)$. Let (Q, G, r) be the corresponding Hopf quiver with $r = r(g_i, \chi_i; i \in J)$ and denote by $(kQ_1^c, g_i, \chi_i; i \in J)$ the kG -Hopf bimodule $(kQ_1^c, G, r, \vec{\chi}, u)$. Denote by $kQ^c(G, g_i, \chi_i; i \in J)$ and $kG[kQ_1^c, G, g_i, \chi_i; i \in J]$ the corresponding multiple crown algebra $kQ^c(G, r, \vec{\chi}, u)$ and multiple Taft algebra $kG[kQ_1^c, G, r, \vec{\chi}, u]$, respectively. We also denote by $kQ^a(G, g_i, \chi_i; i \in J)$, $kQ^{sc}(G, g_i, \chi_i; i \in J)$, $kQ^s(G, g_i, \chi_i; i \in J)$ and $(kG)^*[kQ_1^a, G, g_i, \chi_i; i \in J]$ the corresponding path Hopf algebra $kQ^a(G, r, \vec{\chi}, u)$, semi-co-path Hopf algebra $kQ^{sc}(G, r, \vec{\chi}, u)$, semi-path Hopf algebra $kQ^s(G, r, \vec{\chi}, u)$ and one-type- path Hopf algebra $(kG)^*[kQ_1^a, G, r, \vec{\chi}, u]$, respectively.

Conversely, assume that $(G, r, \vec{\chi}, u)$ is a CRSC. We may assume $I_{\{g\}}(r) \cap I_{\{h\}}(r) = \emptyset$ for any $\{g\} \neq \{h\}$ in $\mathcal{K}_r(G)$. Let $J := \bigcup_{\{g\} \in \mathcal{K}_r(G)} I_{\{g\}}(r)$. For any $i \in J$, put $g_i := g$ and $\chi_i := \chi_{\{g\}}^{(i)}$ if $i \in I_{\{g\}}(r)$ with $\{g\} \in \mathcal{K}_r(G)$. We obtain an ESC, written $\text{ESC}(G, \vec{g}(r, \vec{\chi}, u), \vec{\chi}(r, \vec{\chi}, u), J)$.

From now on, assume $I_{\{g\}}(r) \cap I_{\{h\}}(r) = \emptyset$ for any $\{g\}, \{h\} \in \mathcal{K}_r(G)$ with $\{g\} \neq \{h\}$. Note that, in two cases above, for any $i, j \in I_{\{g\}}(r)$, we have

$$g_i = g_j = g, \quad a_{g_i,1}^{(j)} = a_{g_j,1}^{(j)}, \quad \chi_{\{g_i\}}^{(j)} = \chi_{\{g_j\}}^{(j)} = \chi_j. \quad (14.13)$$

Throughout this chapter, let $E_j := a_{g_j,1}^{(j)}$ for any $j \in J$.

Proposition 14.2.5. $\text{CRSC}(G, r, \vec{\chi}, u) \cong \text{CRSC}(G', r', \vec{\chi}', u')$ if and only if $\text{ESC}(G, \vec{g}(r, \vec{\chi}, u), \vec{\chi}(r, \vec{\chi}, u), J) \cong \text{ESC}(G', \vec{g}'(r', \vec{\chi}', u'), \vec{\chi}'(r', \vec{\chi}', u'), J')$.

Proof. We use notations above. If $\text{CRSC}(G, r, \vec{\chi}, u) \cong \text{CRSC}(G', r', \vec{\chi}', u')$, then there exists a group isomorphism $\phi : G \rightarrow G'$; for any $C \in \mathcal{K}_r(G)$, there exists a bijective map $\phi_C : I_C(r) \rightarrow I_{\phi(C)}(r')$ such that $\chi'_{\phi(C)}(\phi_C(i))\phi = \chi_C^{(i)}$. Let σ be the bijection from J to J' such that $\sigma(i) = \phi_C(i)$ for any $C \in \mathcal{K}_r(G)$, $i \in I_C(r)$. It is clear that $\phi(g_j) = g'_{\sigma(j)}$ and $\chi'_{\sigma(j)}\phi = \chi_j$ for any $j \in J$. Thus $\text{ESC}(G, \vec{g}(r, \vec{\chi}, u), \vec{\chi}(r, \vec{\chi}, u), J) \cong \text{ESC}(G', \vec{g}'(r', \vec{\chi}', u'), \vec{\chi}'(r', \vec{\chi}', u'), J')$.

Conversely, if $\text{ESC}(G, \vec{g}(r, \vec{\chi}, u), \vec{\chi}(r, \vec{\chi}, u), J) \cong \text{ESC}(G', \vec{g}'(r', \vec{\chi}', u'), \vec{\chi}'(r', \vec{\chi}', u'), J')$, then there exist a group isomorphism $\phi : G \rightarrow G'$, a bijective map $\sigma : J \rightarrow J'$ such that $\phi(g_j) = g'_{\sigma(j)}$ and $\chi'_{\sigma(j)}\phi = \chi_j$ for any $j \in J$. For any $C = \{g_i\} \in \mathcal{K}_r(G)$, $j \in I_C(r)$, we define $\phi_C = \sigma|_{I_C(r)}$ and have

$$\chi'_{\phi(C)}(\phi_C(j))\phi = \chi'_{g'_{\sigma(j)}}(\phi_C(j))\phi = \chi'_{\sigma(j)}\phi = \chi_j = \chi_C^{(j)}. \quad \square$$

If $(G, g_i, \chi_i; i \in J)$ is an ESC, then $(kQ_1^e, G, g_i, \chi_i; i \in J)$ is a kG -Hopf bimodule with module operations α^- and α^+ . Define a new left kG -action on kQ_1 by

$$g \triangleright x := g \cdot x \cdot g^{-1}, \quad g \in G, x \in kQ_1,$$

where $g \cdot x = \alpha^-(g \otimes x)$ and $x \cdot g = \alpha^+(x \otimes g)$ for any $g \in G$ and $x \in kQ_1$. With this left kG -action and the original left (arrow) kG -coaction δ^- , kQ_1 is a YD kG -module. Let $Q_1^1 := \{a \in Q_1 \mid s(a) = 1\}$. It is clear that kQ_1^1 is a YD kG -submodule of kQ_1 , denoted by $(kQ_1^1, \text{ad}(G, g_i, \chi_i; i \in J))$.

Lemma 14.2.6. $(kQ_1^1, \text{ad}(G, g_i, \chi_i; i \in J))$ and $V(G, g_i, \chi_i^{-1}; i \in J)$ are isomorphic YD kG -modules.

Proof. By definition, $V(G, g_i, \chi_i^{-1}; i \in J)$ has a k -basis $\{x_i \mid i \in J\}$ such that $\delta^-(x_i) = g_i \otimes x_i$ and $g \cdot x_i = \chi_i^{-1}(g)x_i$ for all $i \in J$ and $g \in G$. By Proposition 14.1.11, for any $j \in J$, we have that $g \triangleright a_{g_j,1}^{(j)} = \chi_j(g^{-1})a_{g_j,1}^{(j)}$ and $\delta^-(a_{g_j,1}^{(j)}) = g_j \otimes a_{g_j,1}^{(j)}$. It follows that there is a YD kG -module isomorphism from $(kQ_1^1, \text{ad}(G, g_i, \chi_i; i \in J))$ to $V(G, g_i, \chi_i^{-1}; i \in J)$ given by $a_{g_j,1}^{(j)} \mapsto x_j$ for any $j \in J$. $\square$

Lemma 14.2.7. Let B and B' be two Hopf algebras with bijective antipodes. Let V be a YD B -module. Assume that there is a Hopf algebra isomorphism $\phi : B' \rightarrow B$. Then $\phi^{-1}\mathcal{B}(V) \cong \mathcal{B}(\phi^{-1}V)$ as graded braided Hopf algebras in $B'_{\phi}\mathcal{YD}$.

Proof. We first show that ${}_{\phi}^{\phi^{-1}}\mathcal{B}(V)$ is a Nichols algebra of ${}_{\phi}^{\phi^{-1}}V$ by following steps. Let $R := {}_{\phi}^{\phi^{-1}}\mathcal{B}(V)$. (i) Obviously, $R_0 = k$, $R_1 = P(R)$ and R is generated by R_1 as algebras. Since $C(x \otimes y) = C'(x' \otimes y')$ for any $x', y' \in R$, $x, y \in \mathcal{B}(V)$ with $x' = x, y' = y$, where C and C' denote the braidings in ${}_{B'}^B\mathcal{YD}$ and ${}_{B'}^{B'}\mathcal{YD}$, respectively, we have that R is a graded braided Hopf algebra in ${}_{B'}^B\mathcal{YD}$. (ii) $R_1 = {}_{\phi}^{\phi^{-1}}\mathcal{B}(V)_1 \cong {}_{\phi}^{\phi^{-1}}V$ as YD B' -modules, since $\mathcal{B}(V)_1 \cong V$ as YD B -modules.

Consequently, ${}_{\phi}^{\phi^{-1}}\mathcal{B}(V)$ is a Nichols algebra of ${}_{\phi}^{\phi^{-1}}V$. By [7, Proposition 2.2 (iv)], $R \cong \mathcal{B}({}_{\phi}^{\phi^{-1}}V)$ as graded braided Hopf algebra in ${}_{B'}^B\mathcal{YD}$. $\square$

Theorem 14.2.8. Assume that $(G, g_i, \chi_i; i \in J)$ and $(G', g'_i, \chi'_i; i \in J')$ are two ESC's. Then the following statements are equivalent:

- (i) $\text{ESC}(G, g_i, \chi_i; i \in J) \cong \text{ESC}(G', g'_i, \chi'_i; i \in J')$.
- (ii) $\text{CRSC}(G, r(G, g_i, \chi_i; i \in J), \vec{\chi}(G, g_i, \chi_i; i \in J), u) \cong \text{CRSC}(G', r'(G', g'_i, \chi'_i; i \in J'), \vec{\chi}'(G', g'_i, \chi'_i; i \in J'), u')$.
- (iii) There is a Hopf algebra isomorphism $\phi : kG \rightarrow kG'$ such that $V(G, g_i, \chi_i; i \in J) \cong {}_{\phi}^{\phi^{-1}}V'(G', g'_i, \chi'_i; i \in J')$ as YD kG -modules.
- (iv) There is a Hopf algebra isomorphism $\phi : kG \rightarrow kG'$ such that $\mathcal{B}(V(G, g_i, \chi_i; i \in J)) \cong {}_{\phi}^{\phi^{-1}}\mathcal{B}(V'(G', g'_i, \chi'_i; i \in J'))$ as graded braided Hopf algebras in ${}_{kG}^{kG'}\mathcal{YD}$.
- (v) There is a Hopf algebra isomorphism $\phi : kG \rightarrow kG'$ such that $(kQ_1^1, \text{ad}(G, g_i, \chi_i; i \in J)) \cong {}_{\phi}^{\phi^{-1}}(kQ_1'^1, \text{ad}(G', g'_i, \chi'_i; i \in J'))$ as YD kG -modules.
- (vi) $kG[kQ_1^c, G, g_i, \chi_i; i \in J] \cong kG'[kQ_1'^c, G', g'_i, \chi'_i; i \in J']$.

Proof. We use the notations before Proposition 14.2.5.

(i) $\Rightarrow$ (ii). There exist a group isomorphism $\phi : G \rightarrow G'$ and a bijective map $\sigma : J \rightarrow J'$ such that $\phi(g_j) = g'_{\sigma(j)}$ and $\chi'_{\sigma(j)}\phi = \chi_j$ for any $j \in J$. For any $C = \{g_i\} \in \mathcal{K}_r(G)$ and $j \in I_C(r)$, we have $g_i = g_j$ and

$$\begin{aligned} \chi_C^{(j)} = \chi_{\{g_i\}}^{(j)} &= \chi_j = \chi'_{\sigma(j)}\phi \\ &= \chi'_{\{g'_{\sigma(i)}\}}(\sigma(j))\phi = \chi'_{\{\phi(g_i)\}}(\phi_{g_i}(j))\phi = \chi'_{\phi(C)}(\phi_C(j))\phi. \end{aligned}$$

(ii) $\Rightarrow$ (i). There is a group isomorphism $\phi : G \rightarrow G'$ and a bijection $\phi_C : I_C(r) \rightarrow I_{\phi(C)}(r')$ such that $\chi'_{\phi(C)}(\phi_C(j))\phi = \chi_C^{(j)}$ for any $C = \{g_i\} \in \mathcal{K}_r(G)$, $j \in I_C(r)$. Define a map $\sigma : J \rightarrow J'$ such that $\sigma|_{I_C(r)} = \phi_C$ for any $C \in \mathcal{K}_r(G)$. Thus σ is bijective. For any $C = \{g_i\} \in \mathcal{K}_r(G)$ and $j \in I_{\{g_i\}}(r)$, we have $\phi(g_j) = \phi(g_i) = g'_{\sigma(j)}$ and $\chi'_{\sigma(j)}\phi = \chi'_{\phi_{\{g_i\}}(j)}\phi = \chi'_{\phi(\{g_i\})}(\phi_{\{g_i\}}(j))\phi = \chi_{\{g_i\}}^{(j)} = \chi_j$. This shows that $\text{ESC}(G, \vec{g}, \vec{\chi}, J) \cong \text{ESC}(G', \vec{g}', \vec{\chi}', J')$.

(i) $\Leftrightarrow$ (iii). Let $V := V(G, g_i, \chi_i; i \in J)$ and $V' := V'(G', g'_i, \chi'_i; i \in J')$. By definition V has a k -basis $\{x_i \mid i \in J\}$ such that $g \cdot x_i = \chi_i(g)x_i$ and $\delta^-(x_i) = g_i \otimes x_i$ for any $i \in J$ and $g \in G$. Similarly, V' has a k -basis $\{y_j \mid j \in J'\}$ such that $h \cdot y_j = \chi'_j(h)y_j$ and $\delta^-(y_j) = g'_j \otimes y_j$ for all $j \in J'$ and $h \in G'$.

Assume $\text{ESC}(G, \vec{g}, \vec{\chi}, J) \cong \text{ESC}(G', \vec{g}', \vec{\chi}', J')$. Then there is a group isomorphism $\phi : G \rightarrow G'$ and a bijective map $\sigma : J \rightarrow J'$ such that $\phi(g_i) = g'_{\sigma(i)}$ and $\chi'_{\sigma(i)}\phi = \chi_i$ for any $i \in J$. Hence $\phi : kG \rightarrow kG'$ is a Hopf algebra isomorphism. Define a k -linear isomorphism $\psi : V \rightarrow V'$

by $\psi(x_i) = y_{\sigma(i)}$ for any $i \in J$. Then it is straightforward to check that ψ is a YD kG -module homomorphism from V to ${}_{\phi}^{\phi^{-1}}V'$.

Conversely, assume that $\phi : kG \rightarrow kG'$ is a Hopf algebra isomorphism and $\psi : V \rightarrow {}_{\phi}^{\phi^{-1}}V'$ is a YD kG -module isomorphism. Then $\phi : G \rightarrow G'$ is a group isomorphism. We use the notations in the proof of Lemma 14.2.2 and the notations above. Then $\psi(V_g) = V'_{\phi(g)}$ for any $g \in G$. Since $V = \bigoplus_{i \in \bar{J}} V_{g_i}$ and $V' = \bigoplus_{i \in \bar{J}'} V'_{g'_i}$, there is a bijection $\tau : \bar{J} \rightarrow \bar{J}'$ such that $\psi(V_{g_i}) = V'_{g'_{\tau(i)}}$ for any $i \in \bar{J}$. This shows that $\phi(g_i) = g'_{\tau(i)}$ and $V_{g_i} \cong {}_{\phi}(V'_{g'_{\tau(i)}})$ as left kG -modules for any $i \in \bar{J}$. However, V_{g_i} is a pointed kG -module and kx_j is its one dimensional submodule for any $j \in [i]$. Similarly, $V'_{g'_{\tau(i)}}$ is a pointed kG' -module and ky_j is its one dimensional submodule for any $j \in [\tau(i)]$. Hence there is a bijection $\phi_{\{g_i\}} : [i] \rightarrow [\tau(i)]$ for any $i \in \bar{J}$ such that kx_j and $\phi(ky_{\phi_{\{g_i\}}(j)})$ are isomorphic kG -modules for all $j \in [i]$. This implies that $\chi_j = \chi'_{\phi_{\{g_i\}}(j)}\phi$ for all $j \in [i]$ and $i \in \bar{J}$. Then the same argument as in the proof of (ii) $\Rightarrow$ (i) shows that $\text{ESC}(G, \vec{g}, \vec{\chi}, J) \cong \text{ESC}(G', \vec{g}', \vec{\chi}', J')$. (iii) $\Leftrightarrow$ (iv) It follows from Lemma 14.2.7. (iii) $\Leftrightarrow$ (v) It follows from Lemma 14.2.6. (ii) $\Leftrightarrow$ (vi) It follows from Theorem 14.1.13. $\square$

Up to now we have classified Nichols algebras and YD modules over finite abelian group and the complex field up to isomorphisms, which are under means of Theorem 14.2.8 (iv)(iii), respectively. In fact, we can explain these facts above by introducing some new concepts about isomorphisms. For convenience, if B is a Hopf algebra and M is a B -Hopf bimodule, then we say that (B, M) is a Hopf bimodules. For any two Hopf bimodules (B, M) and (B', M') , if ϕ is a Hopf algebra homomorphism from B to B' and ψ is simultaneously a B -bimodule homomorphism from M to ${}_{\phi}M'_{\phi}$ and a B' -bicomodule homomorphism from ${}^{\phi}M^{\phi}$ to M' , then (ϕ, ψ) is called a pull-push Hopf bimodule homomorphism. Similarly, we say that (B, M) and (B, X) are a YD module and a YD Hopf algebra if M is a YD B -module and X is a braided Hopf algebra in Yetter-Drinfeld category ${}^B_B\mathcal{YD}$, respectively. For any two YD modules (B, M) and (B', M') , if ϕ is a Hopf algebra homomorphism from B to B' , and ψ is simultaneously a left B -module homomorphism from M to ${}_{\phi}M'$ and a left B' -comodule homomorphism from ${}^{\phi}M$ to M' , then (ϕ, ψ) is called a pull-push YD module homomorphism. For any two YD Hopf algebra (B, X) and (B', X') , if ϕ is a Hopf algebra homomorphism from B to B' , ψ is simultaneously a left B -module homomorphism from X to ${}_{\phi}X'$ and a left B' -comodule homomorphism from ${}^{\phi}X$ to X' , meantime, ψ also is algebra and coalgebra homomorphism from X to X' , then (ϕ, ψ) is called a pull-push YD Hopf algebra homomorphism.

Consequently, we have classified Nichols algebras over finite abelian group and the complex field up to pull-push graded YD Hopf algebra isomorphisms and YD modules over finite abelian group and the complex field up to pull-push YD module isomorphisms, respectively. In other words, element systems with characters uniquely determine their corresponding Nichols algebras and YD modules up to their isomorphisms.

14.3 The relation between quiver Hopf algebras and quotients of free algebras

In this section we show that the diagram of a quantum weakly commutative multiple Taft algebra is not only a Nichols algebra but also a quantum linear space in ${}^k_G\mathcal{YD}$; the diagram of a semi-path Hopf algebra of ESC is a quantum tensor algebra in ${}^k_G\mathcal{YD}$; the quantum enveloping algebra of a complex semisimple Lie algebra is a quotient of a semi-path Hopf algebra.

14.3.1 The structure of multiple Taft algebras and semi-path Hopf algebras

Assume that $H = \bigoplus_{i \geq 0} H_{(i)}$ is a graded Hopf algebra with invertible antipode S . Let $B = H_{(0)}$, and let $\pi_0 : H \rightarrow H_{(0)} = B$ and $\iota_0 : B = H_{(0)} \rightarrow H$ denote the canonical projection and injection. Set $\omega := id_H * (\iota_0 \pi_0 S) : H \rightarrow H$. Then it is clear that (H, δ^+, α^+) is a right B -Hopf module with $\delta^+ := (id \otimes \pi_0)\Delta$ and $\alpha^+ := \mu(id \otimes \iota_0)$. Let $R := H^{coB} := \{h \in H \mid \delta^+(h) = h \otimes 1\}$, which is a graded subspace of H . Then it is known that $R = \text{Im}(\omega)$ and $\Delta(R) \subseteq H \otimes R$. Hence R is a left coideal subalgebra of H , and so R is a left H -comodule algebra. It is well known that R is a graded braided Hopf algebra in ${}^B_B\mathcal{YD}$ with the same multiplication, unit and counit as in H , the comultiplication $\Delta_R = (\omega \otimes id)\Delta$, where the left B -action α_R and left B -coaction δ_R on R are given by

$$\alpha_R(b \otimes x) = b \rightharpoonup_{ad} x = \sum b_{(1)} x S(b_{(2)}), \quad \delta_R^-(x) = \sum \pi_0(x_{(1)}) \otimes x_{(2)}, \quad b \in B, \quad x \in R \quad (14.14)$$

(see the proof of [99, Theorem 3]). R is called the diagram of H , written $diag(H)$. Note that diagram R of H is dependent on the gradation of H . By [99, Theorem 1], the biproduct of R and B is a Hopf algebra, written $R_{\alpha_R}^{\delta_R} \# B$, or $R \# B$ in short. The biproduct $R_{\alpha_R}^{\delta_R} \# B$ is also called the bosonization of R . Furthermore, we have the following well known result.

Theorem 14.3.1. (see [91, p.1530], [4] and [99]) *Under notations above, if $H = \bigoplus_{i \geq 0} H_{(i)}$ is a graded Hopf algebra, then R is a graded braided Hopf algebra in ${}^B_B\mathcal{YD}$ and $R \# B \cong H$ as graded Hopf algebras, where the isomorphism is $\alpha^+ := \mu_H(id_H \otimes \iota_0)$.*

Remark: If A be a Hopf algebra whose coradical A_0 is a Hopf subalgebra, then it is clear that $H := gr A$ is a graded Hopf algebra. The diagram of H with respect to gradation of $gr A$ is called the diagram of A in [5, Introduction].

Lemma 14.3.2. (i) *Assume that H and H' are two graded Hopf algebras with $B = H_{(0)}$ and $B' = H'_{(0)}$. Then $H \cong H'$ as graded Hopf algebras if and only if there exists a Hopf algebra isomorphism $\phi : B \rightarrow B'$ such that $diag(H) \cong \phi^{-1} diag(H')$ as YD B -modules and as graded braided Hopf algebras in ${}^B_B\mathcal{YD}$.*

(ii) *Let B and B' be two Hopf algebras. Let M and M' be B -Hopf bimodule and B' -Hopf bimodule, respectively. Then $B[M] \cong B'[M']$ as graded Hopf algebras if and only if there exists a Hopf algebra isomorphism $\phi : B \rightarrow B'$ such that $diag(B[M]) \cong \phi^{-1} diag(B'[M'])$ as YD B -modules and as graded braided Hopf algebras in ${}^B_B\mathcal{YD}$.*

Proof. (i) Assume that ξ is a graded Hopf algebra isomorphism from H to H' . Let $R := \text{diag}(H)$, $R' := \text{diag}(H')$, $\phi := \xi|_B$ and $\psi := \xi|_R$. It is easy to check that ψ is the map required.

Conversely, by Theorem 14.3.1, $R\#B \cong H$ and $R'\#B' \cong H'$ as graded Hopf algebras. Let ξ be a linear map from $R\#B$ to $R'\#B'$ by sending $r\#b$ to $\psi(r)\#\phi(b)$ for any $r \in R$, $b \in B$. Let ν be a linear map from $R'\#B'$ to $R\#B$ by sending $r'\#b'$ to $\psi^{-1}(r')\#\phi^{-1}(b')$ for any $r' \in R'$, $b' \in B'$. Obviously, ν is the inverse of ξ . Since ψ is graded, so is ξ .

Now we show that ξ is an algebra homomorphism. For any $r, r' \in R$, $b, b' \in B$, see

$$\begin{aligned} \xi\mu_{R\#B}((r\#b) \otimes (r'\#b')) &= \psi(r(b_{(1)} \cdot r'))\#\phi(b_{(2)}b') \\ &= \psi(r)(\phi(b_{(1)}) \cdot \psi(r'))\#\phi(b_{(2)})\phi(b) \\ &\quad (\text{since } \psi \text{ is a pullback module homomorphism} \\ &\quad \text{and an algebra homomorphism.}) \\ &= \mu_{R'\#B'}(\xi(r\#b) \otimes \xi(r'\#b')). \end{aligned}$$

Similarly, we can show that ξ is a coalgebra homomorphism.

(ii) It follows from (i). $\square$

Lemma 14.3.3. (i) $(kQ^c(G, r, \vec{\chi}, u))^{\text{co } kG} = \text{span}\{\beta \mid \beta \text{ is a path with } s(\beta) = 1\}$. (ii) $(kG[kQ_1^c, G, r, \vec{\chi}, u])^{\text{co } kG}$ is the subalgebra of $kG[kQ_1^c, G, r, \vec{\chi}, u]$ generated by Q_1^1 as algebras. (iii) $(kG[kQ_1^c, G, r, \vec{\chi}, u])^{\text{co } kG} \# kG \cong kG[kQ_1^c, G, r, \vec{\chi}, u]$ as graded Hopf algebra isomorphism. (iv) $(kQ^s(G, r, \vec{\chi}, u))^{\text{co } kG}$ is the subalgebra of $kG^s(G, r, \vec{\chi}, u)$ generated by Q_1^1 as algebras.

Proof. (i) For a path β , see that

$$\delta^+(\beta) = (id \otimes \pi_0)\Delta(\beta) = \beta \otimes s(\beta).$$

This implies $(kQ^c)^{\text{co } kG} = \text{span}\{\beta \mid \beta \text{ is a path with } s(\beta) = 1\}$.

(ii) Since every path generated by arrows in Q_1^1 is of start vertex 1, this path belongs to $(kG[kQ_1^c, G, r, \vec{\chi}, u])^{\text{co } kG}$. Let $R := (kG[kQ_1^c, G, r, \vec{\chi}, u])^{\text{co } kG}$ and $A :=$ the subalgebra of $kG[kQ_1^c, G, r, \vec{\chi}, u]$ generated by Q_1^1 as algebras. Obviously, $A \subseteq R$. It is clear that $\alpha^+(R\#kG) = \alpha^+(A\#kG) = kG[kQ_1^c, G, r, \vec{\chi}, u]$ and α^+ is injective. Thus $R\#kG = A\#kG$ and $R = A$.

(iii) It follows from Theorem 14.3.1.

(iv) We first show $(kQ^s)^{\text{co } kG} = \text{span}\{\beta \mid \beta = 1, \text{ or } \beta = \beta_n \otimes_{kG} \beta_{n-1} \otimes_{kG} \cdots \otimes_{kG} \beta_1 \text{ with } \prod_{i=1}^n s(\beta_i) = 1 \text{ and } \beta_i \in Q_1, i = 1, 2, \dots, n; n \in \mathbb{Z}^+\}$. Indeed, obviously right hand side $\subseteq$ the left hand side. For any $\beta = \beta_n \otimes_{kG} \beta_{n-1} \otimes_{kG} \cdots \otimes_{kG} \beta_1$ with $\beta_i \in Q_1$, called a monomial, define $s(\beta) = \prod_{i=1}^n s(\beta_i)$. For any $0 \neq u \in (kQ^s)^{\text{co } kG}$ with $u \notin kG$, there exist linearly independent monomials $u_1, u_2, \dots, u_n$ such that $u = \sum_{i=1}^n b_i u_i$ with $0 \neq b_i \in k$ for $i = 1, 2, \dots, n$. See $\delta^+(u) = \sum_{i=1}^n b_i u_i \otimes s(u_i) = u \otimes 1$. Consequently, $s(u_i) = 1$ for $i = 1, 2, \dots, n$. This implies that u belongs to the right hand side.

For any $\beta = \beta_n \otimes_{kG} \beta_{n-1} \otimes_{kG} \cdots \otimes_{kG} \beta_1$ with $\prod_{i=1}^n s(\beta_i) = 1$ and $\beta_i \in Q_1, i = 1, 2, \dots, n$, we show that β can be written as multiplication of arrows in Q_1^1 by induction. When $n = 1$, it

is clear. For $n > 1$, see $\beta = \beta_n \otimes_{kG} \beta_{n-1} \otimes_{kG} \cdots \otimes_{kG} \beta_2 \cdot s(\beta_1) \otimes_{kG} (s(\beta_1)^{-1} \cdot \beta_1)$. Thus β can be written as multiplication of arrows in Q_1^1 . Consequently, we complete the proof of (iv). $\square$

Recall that a braided algebra A in braided tensor category $(\mathcal{C}, C)$ with braiding C is said to be braided commutative or quantum commutative, if $ab = \mu C(a \otimes b)$ for any $a, b \in A$. An $\text{ESC}(G, g_i, \chi_i; i \in J)$ is said to be quantum commutative if

$$\chi_i(g_j)\chi_j(g_i) = 1$$

for any $i, j \in J$. An $\text{ESC}(G, g_i, \chi_i; i \in J)$ is said to be quantum weakly commutative if

$$\chi_i(g_j)\chi_j(g_i) = 1$$

for any $i, j \in J$ with $i \neq j$.

Lemma 14.3.4. (i) $\text{ESC}(G, g_i, \chi_i; i \in J)$ is quantum weakly commutative if and only if in $\text{diag}(kG[kQ_1^c, G, g_i, \chi_i; i \in J])$,

$$E_i \cdot E_j = \chi_j(g_i^{-1})E_j \cdot E_i \quad (14.15)$$

for any $i, j \in J$ with $i \neq j$.

(ii) $\text{diag}(kG[kQ_1^c, G, g_i, \chi_i; i \in J])$ is quantum commutative in ${}^{kG}_k\mathcal{YD}$ if and only if $\text{ESC}(G, g_i, \chi_i; i \in J)$ is quantum commutative.

Proof. (ii) By Theorem 14.3.1, $R := \text{diag}(kG[kQ_1^c, g_i, \chi_i; i \in J])$ is a braided Hopf algebra in ${}^{kG}_k\mathcal{YD}$. If $\text{diag}(kG[kQ_1^c, g_i, \chi_i; i \in J])$ is quantum commutative in ${}^{kG}_k\mathcal{YD}$, then

$$E_i \cdot E_j = \chi_j(g_i^{-1})E_j \cdot E_i \quad (14.16)$$

for any $i, j \in J$. By [36, Theorem 3.8] and Proposition 14.1.11, we have

$$\begin{aligned} E_i \cdot E_j &= a_{g_i g_j, g_i}^{(j)} a_{g_i, 1}^{(i)} + q_{ij} a_{g_i g_j, g_j}^{(i)} a_{g_j, 1}^{(j)}, \\ E_j \cdot E_i &= q_{ji} a_{g_i g_j, g_i}^{(j)} a_{g_i, 1}^{(i)} + a_{g_i g_j, g_j}^{(i)} a_{g_j, 1}^{(j)}. \end{aligned}$$

Thus $\chi_i(g_j^{-1}) = \chi_j(g_i)$.

Conversely, if $\chi_i(g_j^{-1}) = \chi_j(g_i)$ for any $i, j \in J$, see

$$\begin{aligned} E_i \cdot E_j &= a_{g_i, 1}^{(i)} \cdot a_{g_j, 1}^{(j)} \\ &= a_{g_i g_j, g_i}^{(j)} a_{g_i, 1}^{(i)} + q_{ij} a_{g_i g_j, g_j}^{(i)} a_{g_j, 1}^{(j)} \\ &= q_{ij} (q_{ji} a_{g_i g_j, g_i}^{(j)} a_{g_i, 1}^{(i)} + a_{g_i g_j, g_j}^{(i)} a_{g_j, 1}^{(j)}) \\ &= q_{ij} a_{g_j, 1}^{(j)} \cdot a_{g_i, 1}^{(i)} \\ &= q_{ij} E_j \cdot E_i. \end{aligned}$$

Since E'_i s generate R as algebras, R is quantum commutative.

(i) It is similar to the proof of (ii). $\square$

For any positive integers m and n , let

$$D_n^{n+m} = \{d = (d_{n+m}, d_{n+m-1}, \dots, d_1) \mid d_i = 0 \text{ or } 1, \sum_{i=1}^{n+m} d_i = n\}.$$

Let $d \in D_n^{n+m}$ and let $A = a_n a_{n-1} \cdots a_1 \in Q_n$ be an n -path. We define a sequence $dA = ((dA)_{n+m}, \dots, (dA)_1)$ by

$$(dA)_i = \begin{cases} t(a_{d(i)}) & , \text{ if } d_i = 0; \\ a_{d(i)} & , \text{ if } d_i = 1, \end{cases}$$

where $1 \leq i \leq n+m$ and $d(i) = \sum_{j=1}^i d_j$. Such a sequence dA is called an $n+m$ -thin splits of the n -path A . Note that if $d(i) = 0$ then we regard $t(a_{d(i)}) = s(a_1)$, since $s(a_{d(i)+1}) = s(a_1)$ in this case.

If $0 \neq q \in k$ and $0 \leq i \leq n < \text{ord}(q)$ (the order of q), we set $(0)_q! = 1$,

$$\binom{n}{i}_q = \frac{(n)_q!}{(i)_q!(n-i)_q!}, \quad \text{where } (n)_q! = \prod_{1 \leq i \leq n} (i)_q, \quad (n)_q = \frac{q^n - 1}{q - 1}.$$

In particular, $(n)_q = n$ when $q = 1$.

Lemma 14.3.5. *In $kQ^c(G, r, \vec{\chi}, u)$, assume $\{g\} \in \mathcal{K}_r(G)$ and $j \in I_{\{g\}}(r)$. Let $q := \chi_{\{g\}}^{(j)}(g)$. If $i_1, i_2, \dots, i_m$ be non-negative integers, then*

$$a_{g^{i_m+1}, g^{i_m}}^{(j)} \cdot a_{g^{i_{m-1}+1}, g^{i_{m-1}}}^{(j)} \cdots \cdots a_{g^{i_1+1}, g^{i_1}}^{(j)} = q^{\beta_m}(m)_q! P_{g^{\alpha_m}}^{(j)}(g, m)$$

where $\alpha_m = i_1 + i_2 + \cdots + i_m$, $P_h^{(j)}(g, m) = a_{g^m h, g^{m-1} h}^{(j)} a_{g^{m-1} h, g^{m-2} h}^{(j)} \cdots a_{g h, h}^{(j)}$, $\beta_1 = 0$ and $\beta_m = \sum_{j=1}^{m-1} (i_1 + i_2 + \cdots + i_j)$ if $m > 1$.

Proof. We prove the equality by induction on m . For $m = 1$, it is easy to see that the equality holds. Now suppose $m > 1$. We have

$$\begin{aligned} & a_{g^{i_m+1}, g^{i_m}}^{(j)} \cdot a_{g^{i_{m-1}+1}, g^{i_{m-1}}}^{(j)} \cdots \cdots a_{g^{i_1+1}, g^{i_1}}^{(j)} \\ &= a_{g^{i_m+1}, g^{i_m}}^{(j)} \cdot (a_{g^{i_{m-1}+1}, g^{i_{m-1}}}^{(j)} \cdots \cdots a_{g^{i_1+1}, g^{i_1}}^{(j)}) \\ &= q^{\beta_{m-1}}(m-1)_q! a_{g^{i_m+1}, g^{i_m}}^{(j)} \cdot P_{g^{\alpha_{m-1}}}^{(j)}(g, m-1) \quad (\text{by inductive assumption}) \\ &= q^{\beta_{m-1}}(m-1)_q! a_{g^{i_m+1}, g^{i_m}}^{(j)} \cdot (a_{g^{\alpha_{m-1}+m-1}, g^{\alpha_{m-1}+m-2}}^{(j)} \cdots a_{g^{\alpha_{m-1}+1}, g^{\alpha_{m-1}}}^{(j)}) \\ &= q^{\beta_{m-1}}(m-1)_q! \sum_{l=1}^m [(g^{i_m+1} \cdot a_{g^{\alpha_{m-1}+m-1}, g^{\alpha_{m-1}+m-2}}^{(j)}) \cdots (g^{i_m+1} \cdot a_{g^{\alpha_{m-1}+l}, g^{\alpha_{m-1}+l-1}}^{(j)}) \\ &\quad (a_{g^{i_m+1}, g^{i_m}}^{(j)} \cdot g^{\alpha_{m-1}+l-1}) (g^{i_m} \cdot a_{g^{\alpha_{m-1}+l-1}, g^{\alpha_{m-1}+l-2}}^{(j)}) \cdots (g^{i_m} \cdot a_{g^{\alpha_{m-1}+1}, g^{\alpha_{m-1}}}^{(j)})] \\ &\quad (\text{by [36, Theorem 3.8]}) \\ &= q^{\beta_{m-1}}(m-1)_q! \sum_{l=1}^m [a_{g^{\alpha_m+m}, g^{\alpha_m+m-1}}^{(j)} \cdots a_{g^{\alpha_m+l+1}, g^{\alpha_m+l}}^{(j)} \\ &\quad (\chi_{\{g\}}^{(j)}(g^{\alpha_{m-1}+l-1}) a_{g^{\alpha_m+l}, g^{\alpha_m+l-1}}^{(j)}) a_{g^{\alpha_m+l-1}, g^{\alpha_m+l-2}}^{(j)} \cdots a_{g^{\alpha_m+1}, g^{\alpha_m}}^{(j)}] \quad (\text{by Proposition 14.1.11}) \\ &= q^{\beta_{m-1}}(m-1)_q! \sum_{l=1}^m q^{\alpha_{m-1}+l-1} P_{g^{\alpha_m}}^{(j)}(g, m) \\ &= q^{\beta_{m-1}+\alpha_{m-1}}(m)_q! P_{g^{\alpha_m}}^{(j)}(g, m) \\ &= q^{\beta_m}(m)_q! P_{g^{\alpha_m}}^{(j)}(g, m). \quad \square \end{aligned}$$

Lemma 14.3.6. *(See [5, Lemma 3.3]) Let B be a Hopf algebra and R a braided Hopf algebra in ${}^B_B \mathcal{YD}$ with a linearly independent set $\{x_1, \dots, x_t\} \subseteq P(R)$. Assume that there exist $g_j \in G(B)$ (the set of all group-like elements in B) and $\chi_j \in \text{Alg}(B, k)$ such that*

$$\delta(x_j) = g_j \otimes x_j, \quad h \cdot x_j = \chi_j(h) x_j, \quad \text{for all } h \in B, j = 1, 2, \dots, t.$$

Then

$$\{x_1^{m_1}x_2^{m_2}\cdots x_t^{m_t} \mid 0 \leq m_j < N_j, 1 \leq j \leq t\}.$$

is linearly independent, where N_i is the order of $q_i := \chi_i(g_i)$ ($N_i = \infty$ when q_i is not a root of unit, or $q_i = 1$) for $1 \leq i \leq t$.

Proof. By the quantum binomial formula, if $1 \leq n_j < N_j$, then

$$\Delta(x_j^{n_j}) = \sum_{0 \leq i_j \leq n_j} \binom{n_j}{i_j}_{q_j} x_j^{i_j} \otimes x_j^{n_j - i_j}.$$

We use the following notation:

$$\mathbf{n} = (n_1, \dots, n_j, \dots, n_t), \quad x^{\mathbf{n}} = x_1^{n_1} \cdots x_j^{n_j} \cdots x_t^{n_t}, \quad |\mathbf{n}| = n_1 + \cdots + n_j + \cdots + n_t;$$

accordingly, $\mathbf{N} = (N_1, \dots, N_t)$, $\mathbf{1} = (1, \dots, 1)$. Also, we set

$$\mathbf{i} \leq \mathbf{n} \quad \text{if } i_j \leq n_j, j = 1, \dots, t.$$

Then, for $\mathbf{n} < \mathbf{N}$, we deduce from the quantum binomial formula that

$$\Delta(x^{\mathbf{n}}) = x^{\mathbf{n}} \otimes 1 + 1 \otimes x^{\mathbf{n}} + \sum_{0 \leq \mathbf{i} \leq \mathbf{n}, 0 \neq \mathbf{i} \neq \mathbf{n}} c_{\mathbf{n}, \mathbf{i}} x^{\mathbf{i}} \otimes x^{\mathbf{n} - \mathbf{i}}, \quad (14.17)$$

where $c_{\mathbf{n}, \mathbf{i}} \neq 0$ for all $\mathbf{i}$.

We shall prove by induction on r that the set

$$\{x^{\mathbf{n}} \mid |\mathbf{n}| \leq r, \mathbf{n} < \mathbf{N}\}$$

is linearly independent.

Let $r = 1$ and let $a_0 + \sum_{i=1}^t a_i x_i = 0$, with $a_j \in k$, $0 \leq j \leq t$. Applying ϵ , we see that $a_0 = 0$; by hypothesis we conclude that the other a_j 's are also 0.

Now let $r > 1$ and suppose that $z = \sum_{|\mathbf{n}| \leq r, \mathbf{n} < \mathbf{N}} a_{\mathbf{n}} x^{\mathbf{n}} = 0$. Applying ϵ , we see that $a_0 = 0$. Then

$$\begin{aligned} 0 &= \Delta(z) = z \otimes 1 + 1 \otimes z + \sum_{1 \leq |\mathbf{n}| \leq r, \mathbf{n} < \mathbf{N}} a_{\mathbf{n}} \sum_{0 \leq \mathbf{i} \leq \mathbf{n}, 0 \neq \mathbf{i} \neq \mathbf{n}} c_{\mathbf{n}, \mathbf{i}} x^{\mathbf{i}} \otimes x^{\mathbf{n} - \mathbf{i}} \\ &= \sum_{1 \leq |\mathbf{n}| \leq r, \mathbf{n} < \mathbf{N}} \sum_{0 \leq \mathbf{i} \leq \mathbf{n}, 0 \neq \mathbf{i} \neq \mathbf{n}} a_{\mathbf{n}} c_{\mathbf{n}, \mathbf{i}} x^{\mathbf{i}} \otimes x^{\mathbf{n} - \mathbf{i}}. \end{aligned}$$

Now, if $|\mathbf{n}| \leq r$, $0 \leq \mathbf{i} \leq \mathbf{n}$, and $0 \neq \mathbf{i} \neq \mathbf{n}$, then $|\mathbf{i}| < r$ and $|\mathbf{n} - \mathbf{i}| < r$. By inductive hypothesis, the elements $x^{\mathbf{i}} \otimes x^{\mathbf{n} - \mathbf{i}}$ are linearly independent. Hence $a_{\mathbf{n}} c_{\mathbf{n}, \mathbf{i}} = 0$ and $a_{\mathbf{n}} = 0$ for all $\mathbf{n}$, $|\mathbf{n}| \geq 1$. Thus $a_{\mathbf{n}} = 0$ for all $\mathbf{n}$. $\square$

Assume that $(G, g_i, \chi_i; j \in J)$ is an ESC. Let $\mathcal{T}(G, g_i, \chi_i; j \in J)$ be the free algebra generated by set $\{x_j \mid j \in J\}$. Let $\mathcal{S}(G, g_i, \chi_i; j \in J)$ be the algebra generated by set $\{x_j \mid j \in J\}$ with relations

$$x_i x_j = \chi_j(g_i) x_j x_i \quad \text{for any } i, j \in J \text{ with } i \neq j. \quad (14.18)$$

Let $\mathcal{R}(G, g_i, \chi_i; j \in J)$ be the algebra generated by set $\{x_j \mid j \in J\}$ with relations

$$x_l^{N_l} = 0, \quad x_i x_j = \chi_j(g_i) x_j x_i \quad \text{for any } i, j, l \in J \text{ with } N_l < \infty, i \neq j. \quad (14.19)$$

Define their coalgebra operations and kG -(co-)module operations as follows:

$$\Delta x_i = x_i \otimes 1 + 1 \otimes x_i, \quad \epsilon(x_i) = 0, \quad \delta^-(x_i) = g_i \otimes x_i, \quad h \cdot x_i = \chi_i(h) x_i. \quad (14.20)$$

$\mathcal{T}(G, g_i, \chi_i; j \in J)$ is called a quantum tensor algebra in ${}^{kG}_{kG}\mathcal{YD}$, $\mathcal{S}(G, g_i, \chi_i; j \in J)$ is called a quantum symmetric algebra in ${}^{kG}_{kG}\mathcal{YD}$ and $\mathcal{R}(G, g_i, \chi_i; j \in J)$ is called a quantum linear space in ${}^{kG}_{kG}\mathcal{YD}$. Note that when $\text{ESC}(G, g_i, \chi_i; i \in J)$ is quantum weakly commutative with finite J and finite N_j for any $j \in J$, the definition of quantum linear space is the same as in [5, Lemma 3.4]. Obviously, if N_i is infinite for all $i \in J$, then $\mathcal{S}(G, g_i, \chi_i; j \in J) = \mathcal{R}(G, g_i, \chi_i; j \in J)$.

Theorem 14.3.7. *Assume that $\text{ESC}(G, g_i, \chi_i; i \in J)$ is quantum weakly commutative. Let $\prec$ be a total order of J . Then*

(i) *The multiple Taft algebra $kG[kQ_1^c, G, g_i, \chi_i; i \in J]$ has a k -basis*

$$\{g \cdot E_{\nu_1}^{m_1} \cdot E_{\nu_2}^{m_2} \cdots E_{\nu_t}^{m_t} \mid 0 \leq m_j < N_j; \nu_j \prec \nu_{j+1}, \nu_j \in J, j = 1, 2, \dots, t; t \in \mathbb{Z}^+, g \in G\}.$$

Moreover, $kG[kQ_1^c, G, g_i, \chi_i; i \in J]$ is finite dimensional if and only if $|G|$, $|J|$ and N_j are finite for any $j \in J$. In this case, $\dim_k(kG[kQ_1^c, G, g_i, \chi_i; i \in J]) = |G|N_1 N_2 \cdots N_t$ with $J = \{1, 2, \dots, t\}$.

(ii) *diag($kG[kQ_1^c, G, g_i, \chi_i; i \in J]$) has a k -basis*

$$\{E_{\nu_1}^{m_1} \cdot E_{\nu_2}^{m_2} \cdots E_{\nu_t}^{m_t} \mid 0 \leq m_j < N_j; \nu_j \prec \nu_{j+1}, \nu_j \in J, j = 1, 2, \dots, t; t \in \mathbb{Z}^+\}. \quad (14.21)$$

(iii) *diag($kG[kQ_1^c, G, g_i, \chi_i; i \in J]$) is a Nichols algebra in ${}^{kG}_{kG}\mathcal{YD}$ and $\mathcal{R}(G, g_i, \chi_i^{-1}; j \in J) \cong \text{diag}(kG[kQ_1^c, G, g_i, \chi_i; i \in J])$ as graded braided Hopf algebras in ${}^{kG}_{kG}\mathcal{YD}$, by sending x_j to $a_{g_j, 1}^{(j)}$ for any $j \in J$.*

(iv) *$\mathcal{T}(G, g_i, \chi_i^{-1}; j \in J) \cong \text{diag}(kQ^s(g_i, \chi_i; i \in J))$ as graded braided Hopf algebras in ${}^{kG}_{kG}\mathcal{YD}$ algebras, by sending x_j to $a_{g_j, 1}^{(j)}$ for any $j \in J$.*

(v) *$kQ^s(G, g_i, \chi_i; i \in J) \cong \mathcal{T}(G, g_i, \chi_i^{-1}; j \in J) \# kG$ as graded Hopf algebras and $kQ^s(G, g_i, \chi_i; i \in J)$ has a k -basis*

$$\{g \cdot E_{\nu_1} \otimes_{kG} E_{\nu_2} \otimes_{kG} \cdots \otimes_{kG} E_{\nu_t} \mid \nu_j \in J, j = 1, 2, \dots, t; t \in \mathbb{Z}^+ \cup \{0\}, g \in G\},$$

where $g \cdot E_{\nu_1} \otimes_{kG} E_{\nu_2} \otimes_{kG} \cdots \otimes_{kG} E_{\nu_t} = g$ when $t = 0$.

Note that (iv) and (v) still hold without quantum weakly commutative condition.

Proof. (ii) Since $(N_j)_{q_j}! = 0$, it follows from Lemma 14.3.3 that $E_j^{N_j} = 0$ when $N_j < \infty$. By Lemma 14.3.4, $E_i \cdot E_j = \chi_j(g_i^{-1}) E_j \cdot E_i$ for any $i, j \in J$ with $i \neq j$. Considering Lemma 14.3.6, we complete the proof.

(iii) By Lemma 14.3.3 and Eq.(14.15), there exists an algebra homomorphism ψ from $\mathcal{R}(G, g_i, \chi_i^{-1}; j \in J)$ to $\text{diag}(kG[kQ_1^c, G, g_i, \chi_i; i \in J])$ by sending x_j to $a_{g_j, 1}^{(j)}$ for any $j \in J$. By (ii), ψ is bijective. It is clear that ψ is a graded braided Hopf algebra isomorphism.

Let $R := \mathcal{R}(G, g_i, \chi_i; i \in J)$. Obviously, $R_{(1)} \subseteq P(R)$. It is sufficient to show that any non-zero homogeneous element $z \in R$, whose degree $\deg(z)$ is not equal to 1, is not a primitive element. Obviously, z is not a primitive element when $\deg(z) = 0$. Now $\deg(z) > 1$. We can assume, without lost generality, that there exist $\nu_1, \nu_2, \dots, \nu_t \in J$ such that $z = \sum_{|\mathbf{i}|=n} k_{\mathbf{i}} x^{\mathbf{i}}$, where $k_{\mathbf{i}} \in k$, $x^{\mathbf{i}} = x_{\nu_1}^{i_1} x_{\nu_2}^{i_2} \cdots x_{\nu_t}^{i_t}$ with $i_1 + i_2 + \cdots + i_t = n$. It is clear

$$\Delta(z) = \sum_{|\mathbf{i}|=n} k_{\mathbf{i}} \Delta(x^{\mathbf{i}}) = z \otimes 1 + 1 \otimes z + \sum_{|\mathbf{i}|=n} \sum_{0 \leq \mathbf{j} \leq \mathbf{i}, 0 \neq \mathbf{j} \neq \mathbf{i}} k_{\mathbf{i}} c_{\mathbf{i}, \mathbf{j}} x^{\mathbf{j}} \otimes x^{\mathbf{i}-\mathbf{j}}. \quad (14.22)$$

If z is a primitive element, then $\sum_{|\mathbf{i}|=n} \sum_{0 \leq \mathbf{j} \leq \mathbf{i}, 0 \neq \mathbf{j} \neq \mathbf{i}} k_{\mathbf{i}} c_{\mathbf{i}, \mathbf{j}} x^{\mathbf{j}} \otimes x^{\mathbf{i}-\mathbf{j}} = 0$. Since $c_{\mathbf{i}, \mathbf{j}} \neq 0$, we have $k_{\mathbf{i}} = 0$ for any $\mathbf{i}$ with $|\mathbf{i}| = n$, hence $z = 0$. We get a contradiction. Thus z is not a primitive element. This show $R_{(1)} = P(R)$ and R is a Nichols algebra.

(iv) and (v). Let $A := \mathcal{T}(G, g_i, \chi_i^{-1}; j \in J)$ and $R = \text{diag}(kQ^s(G, g_i, \chi_i; i \in J))$. Let ψ be an algebra homomorphism from $\mathcal{T}(G, g_i, \chi_i^{-1}; j \in J)$ to $\text{diag}(kQ^s(G, g_i, \chi_i; i \in J))$ by sending x_j to $a_{g_j, 1}^{(j)}$.

It is clear that $\mathcal{T}(G, g_i, \chi_i^{-1}; j \in J)$ is a kG -module algebra. Define a linear map ν from $A \# kG$ to kQ^s by sending $x_j \# g$ to $a_{g_j, 1}^{(j)} \cdot g = \alpha^+(a_{g_j, 1}^{(j)} \otimes g)$ for any $g \in G, j \in J$. That is, ν is the composition of

$$A \# kG \xrightarrow{\psi \otimes id} R \# kG \xrightarrow{\alpha^+} kQ^s,$$

where $\alpha^+ = \mu_{kQ^s}(id \otimes \iota_0)$ (see Theorem 14.3.1). Define a linear map λ from kG to $A \# kG$ by sending g to $1 \# g$ for any $g \in G$ and another linear map γ from kQ_1^c to $A \# kG$ by sending $a_{g_i h, h}^{(i)}$ to $\chi_i^{-1}(h) x_i \# h$ for any $h \in G, i \in J$. It is clear that γ is a kG -bimodule homomorphism from kQ_1^c to ${}_{\lambda}(A \# kG)_{\lambda}$. Considering $kQ^s = T_{kG}(kQ_1^c)$ and universal property of tensor algebra over kG , we have that there exists an algebra homomorphism $\phi = T_{kG}(\lambda, \gamma)$ from kQ^s to $A \# kG$. Obviously, ϕ is the inverse of ν . Thus ϕ is bijective. It is easy to check that ϕ is graded Hopf algebra isomorphism. Obviously, $\{x_{\nu_1} x_{\nu_2} \cdots x_{\nu_t} \# g \mid \nu_j \in J, j = 1, 2, \dots, t; t \in \mathbb{Z}^+ \cup \{0\}, g \in G\}$ is a basis of $\mathcal{T}(G, g_i, \chi_i^{-1}; j \in J) \# kG$. See

$$\begin{aligned} & \nu(x_{\nu_1} x_{\nu_2} \cdots x_{\nu_t} \# g) \\ &= E_{\nu_1} \otimes_{kG} E_{\nu_2} \otimes_{kG} \cdots \otimes_{kG} E_{\nu_t} \cdot g \\ &= \chi_{\nu_1}(g) \chi_{\nu_2}(g) \cdots \chi_{\nu_t}(g) g \cdot E_{\nu_1} \otimes_{kG} E_{\nu_2} \otimes_{kG} \cdots \otimes_{kG} E_{\nu_t}. \end{aligned}$$

Thus $\{g \cdot E_{\nu_1} \otimes_{kG} E_{\nu_2} \otimes_{kG} \cdots \otimes_{kG} E_{\nu_t} \mid \nu_j \in J, j = 1, 2, \dots, t; t \in \mathbb{Z}^+ \cup \{0\}, g \in G\}$ is a basis of kQ^s . It is easy to check that ψ is graded braided Hopf algebra isomorphism.

(i) Considering (ii), Lemma 14.3.3 and Theorem 14.3.1, we complete the proof. $\square$

14.3.2 A characterization of multiple Taft algebras

In this subsection we characterize multiple Taft algebras by means of elements in themselves.

Definition 14.3.8. For a quantum weakly commutative $\text{ESC}(G, g_i, \chi_i; i \in J)$, let A be the Hopf algebra to satisfy the following conditions: (i) G is a subgroup of $G(A)$; (ii) there exists a linearly independent subset $\{X_i \mid i \in J\}$ of A such that A is generated by set $\{X_i \mid i \in J\} \cup G$

as algebras; (iii) X_j is $(1, g_j)$ -primitive, i.e., $\Delta(X_j) = X_j \otimes 1 + g_j \otimes X_j$, for any $j \in J$; (iv) $X_j g = \chi_j(g) g X_j$, for any $j \in J$, $g \in G$; (v) $X_j X_i = \chi_j(g_i) X_i X_j$, for $i, j \in J$ with $i \neq j$; (vi) $A_{(0)} \cap A_{(1)} = 0$, where $A_{(0)} := kG$ and $A_{(1)}$ is the vector space spanned by set $\{hX_i \mid i \in J; h \in G\}$. Furthermore, let $J(A)$ denote the ideal of A generated by the set

$$\{X_i^{N_i} \mid N_i < \infty, i \in J\}$$

and $H(G, g_i, \chi_i; i \in J)$ the quotient algebra $A/J(A)$.

Lemma 14.3.9. (See [5, Lemma 3.3]) Let H be a Hopf algebra with a linearly independent set $\{x_1, \dots, x_t\}$ and G a subgroup of $G(H)$. Assume that $g_i \in Z(G)$ and $\chi_i \in \hat{G}$ such that $\Delta(x_i) = x_i \otimes 1 + g_i \otimes x_i$, $x_i h = \chi_i(h) h x_i$, for $i = 1, 2, \dots, t$, $h \in G$. If the intersection of kG and $\text{span} \{h x_i \mid 1 \leq i \leq t, h \in G\}$ is zero, then

$$\{h x_1^{m_1} x_2^{m_2} \cdots x_t^{m_t} \mid 0 \leq m_j < N_j, 1 \leq j \leq t; h \in G\}.$$

is linearly independent, where N_i is the order of $q_i := \chi_i(g_i)$ ($N_i = \infty$ when q_i is not a root of unit, or $q_i = 1$) for $1 \leq i \leq t$.

Proof. (i) We first show set $\{h x_i \mid i = 1, 2, \dots, t; h \in G\}$ is linearly independent. Indeed, suppose that $z = \sum_{h \in G} \sum_{i=1}^t k_{i,h} h x_i = 0$ with $k_{i,h} \in k$ for $h \in G$, $1 \leq i \leq t$. Applying Δ , we get

$$\sum_{h \in G} \sum_{i=1}^t k_{i,h} (h g_i \otimes h x_i + h x_i \otimes h) = 0,$$

which implies that $\sum_{i=1}^t k_{i,h} h x_i = 0$ for any $h \in H$, hence $k_{i,h} = 0$ for any $h \in H$, $i = 1, 2, \dots, t$.

(ii) We use the notation in the proof of Lemma 14.3.6. We shall prove by induction on r that the set

$$\{h x^{\mathbf{n}} \mid |\mathbf{n}| \leq r, \mathbf{n} < \mathbf{N}; h \in G\}$$

is linearly independent.

By the quantum binomial formula, if $1 \leq n_j < N_j$, then

$$\Delta(x_j^{n_j}) = \sum_{0 \leq i_j \leq n_j} \binom{n_j}{i_j}_{q_j} g_j^{i_j} x_j^{n_j - i_j} \otimes x_j^{i_j}. \quad (14.23)$$

For $\mathbf{n} < \mathbf{N}$, we deduce from Eq. (14.23) or [40, Eq. (5.9)] that

$$\Delta(x^{\mathbf{n}}) = x^{\mathbf{n}} \otimes 1 + g^{\mathbf{n}} \otimes x^{\mathbf{n}} + \sum_{0 \leq \mathbf{i} \leq \mathbf{n}, 0 \neq \mathbf{i} \neq \mathbf{n}} c_{\mathbf{n}, \mathbf{i}} g^{\mathbf{i}} x^{\mathbf{n} - \mathbf{i}} \otimes x^{\mathbf{i}}, \quad (14.24)$$

where $g^{\mathbf{i}} = g_1^{i_1} g_2^{i_2} \cdots g_t^{i_t}$ and $0 \neq c_{\mathbf{n}, \mathbf{i}} \in k$ for all $\mathbf{i}$. Obviously, the claim holds when $r = 0$.

Let $r = 1$ and let $\sum_{h \in G} b_h h + \sum_{h \in G} \sum_{i=1}^t b_{h,i} h x_i = 0$, with $b_{h,j}, b_h \in k$, $0 \leq j \leq t$. It follows from hypothesis that $\sum_{h \in G} b_h h = 0$ and $\sum_{h \in G} \sum_{i=1}^t b_{h,i} h x_i = 0$, hence $b_h = 0$ and $b_{h,i} = 0$ by (i) for any $h \in G$, $1 \leq i \leq t$.

Now let $r > 1$ and suppose that $z = \sum_{h \in G} \sum_{|\mathbf{n}| \leq r, \mathbf{n} < \mathbf{N}} b_{h, \mathbf{n}} h x^{\mathbf{n}} = 0$. Then

$$\begin{aligned} 0 &= \Delta(z) = z \otimes 1 + 1 \otimes z + \sum_{h \in G} b_{h, \mathbf{0}} h \otimes h + \sum_{h \in G} \sum_{1 \leq |\mathbf{n}| \leq r, \mathbf{n} < \mathbf{N}} b_{h, \mathbf{n}} \sum_{0 \leq \mathbf{i} \leq \mathbf{n}, 0 \neq \mathbf{i} \neq \mathbf{n}} c_{\mathbf{n}, \mathbf{i}} h g^{\mathbf{i}} x^{\mathbf{n}-\mathbf{i}} \otimes x^{\mathbf{i}} \\ &= \sum_{h \in G} \sum_{1 \leq |\mathbf{n}| \leq r, \mathbf{n} < \mathbf{N}} \sum_{0 \leq \mathbf{i} \leq \mathbf{n}, 0 \neq \mathbf{i} \neq \mathbf{n}} b_{h, \mathbf{n}} c_{\mathbf{n}, \mathbf{i}} h g^{\mathbf{i}} x^{\mathbf{n}-\mathbf{i}} \otimes x^{\mathbf{i}}. \end{aligned}$$

Now, if $|\mathbf{n}| \leq r$, $0 \leq \mathbf{i} \leq \mathbf{n}$, and $0 \neq \mathbf{i} \neq \mathbf{n}$, then $|\mathbf{i}| < r$ and $|\mathbf{n} - \mathbf{i}| < r$. By inductive hypothesis, the set $\{h g^{\mathbf{i}} x^{\mathbf{n}-\mathbf{i}} \otimes h x^{\mathbf{i}} \mid \mathbf{0} \leq \mathbf{i} \leq \mathbf{n}, \mathbf{0} \neq \mathbf{i} \neq \mathbf{n}, |\mathbf{n}| < r, \mathbf{n} < \mathbf{N}, h \in G\} \cup \{h \otimes h \mid h \in G\}$ are linearly independent. Hence $b_{h, \mathbf{0}} = 0$ and $b_{h, \mathbf{n}} c_{\mathbf{n}, \mathbf{i}} = 0$, which implies $b_{h, \mathbf{n}} = 0$ for any $h \in G$, all $\mathbf{n}$. $\square$

Proposition 14.3.10. *If $\text{ESC}(G, g_i, \chi_i; i \in J)$ is quantum weakly commutative, then $H(G; g_i, \chi_i; i \in J)$ and multiple Taft algebra $kG[kQ_1^C, G, g_i, \chi_i; i \in J]$ are isomorphic as graded Hopf algebras.*

Proof. Let $H := H(G; g_i, \chi_i; i \in J)$. Denote by X_j the image of X_j under the canonical projection $A \rightarrow A/J(A)$. We say that $h X_{\nu_1}^{i_1} X_{\nu_2}^{i_2} \cdots X_{\nu_t}^{i_t}$ is a monomial with degree n if $h \in G$ and $i_1 + i_2 + \cdots + i_t = n$, where $\nu_1, \nu_2, \dots, \nu_t \in J$, $t \in \mathbb{Z}^+$. Let $H_{(0)} := kG$ and $H_{(n)}$ be the vector space spanned by all monomials with degree n . It follows from Lemma 14.3.9 that H is a graded Hopf algebra. By the method similar to the proof of Lemma 14.3.3(ii), we can obtain that $H^{co \ kG}$ is a subalgebra of H generated by X'_i 's. By Theorem 14.3.1, $H^{co \ kG}$ is a braided Hopf algebra in ${}^{kG} \mathcal{YD}$ and H is the bosonization of $H^{co \ kG}$. Considering Theorem 14.3.7 (iii), we only need to show that $\mathcal{R}(G, g_i, \chi_i^{-1}; j \in J) \cong H^{co \ kG}$ as graded braided Hopf algebras. Let $\prec$ be a total order of J . Since $(N_j)_{q_j}! = 0$, it follows from Lemma 14.3.3 that $X_j^{N_j} = 0$ when $N_j < \infty$. Considering Lemma 14.3.6, we have that

$$\{X_{\nu_1}^{m_1} X_{\nu_2}^{m_2} \cdots X_{\nu_t}^{m_t} \mid 0 \leq m_j < N_j; \nu_j \prec \nu_{j+1}, \nu_j \in J, j = 1, 2, \dots, t; t \in \mathbb{Z}^+\}$$

is a basis of $\text{diag}(H)$. Consequently, we have an algebra isomorphism ψ from $\mathcal{R}(G, g_i, \chi_i^{-1}; i \in J)$ to $H^{co \ kG}$ by sending x_i to X_i . It is easy to check that ψ is also a graded braided Hopf algebra isomorphism. $\square$

Remark: $H(G, g_i, \chi_i; i \in J)$ just is $H(C, n, c, c^*, 0, 0)$ in [40, Definition 5.6.8 and Definition 5.6.15] with $G = C$, $J = \{1, 2, \dots, t\}$, $1 < n_i = N_i < \infty$, $g_i = c_i, c_i^* = \chi_i$ for $i = 1, 2, \dots, t$.

14.3.3 The relation between semi-path Hopf algebras and quantum enveloping algebras

If $0 \neq q \in k$ and $0 \leq i \leq n < \text{ord}(q)$ (the order of q), we set

$$\begin{bmatrix} n \\ i \end{bmatrix}_q = \frac{[n]_q!}{[i]_q! [n-i]_q!}, \quad \text{where } [n]_q! = \prod_{1 \leq i \leq n} [i]_q, \quad [n]_q = \frac{q^n - q^{-n}}{q - q^{-1}}.$$

Let B be a Hopf algebra and R a braided Hopf algebra in ${}^B_B \mathcal{YD}$. For convenience, we denote $r \# 1$ by r and $1 \# b$ by b in biproduct $R \# B$ for any $r \in R$, $b \in B$.

Lemma 14.3.11. *Let B be a Hopf algebra and R a braided Hopf algebra in ${}^B_B\mathcal{YD}$ with $x_1, x_2 \in P(R)$ and $g_1, g_2 \in Z(G(B))$. Assume that there exist $\chi_1, \chi_2 \in \text{Alg}(B, k)$ with $\sqrt{\chi_i(g_j)} \in k$ such that*

$$\delta_R(x_j) = g_j \otimes x_j, \quad h \cdot x_j = \chi_j(h^{-1})x_j,$$

for all $h \in B, j = 1, 2$.

(i) *If r is a positive integer and*

$$\chi_2(g_1)\chi_1(g_2)\chi_1(g_1)^{r-1} = 1, \quad \chi_1(g_1)^{\frac{1}{2}(r-1)}\chi_2(g_1) = 1, \quad r-1 < \text{ord}(\chi_1(g_1)), \quad (14.25)$$

then

$$\sum_{m=0}^r (-1)^m \begin{bmatrix} r \\ m \end{bmatrix}_{\sqrt{\chi_1(g_1^{-1})}} x_1^{r-m} x_2 x_1^m \quad (14.26)$$

= $(ad_c x_1)^r x_2$ is a primitive element of R and a $(1, g_1^r g_2)$ -primitive element of biproduct $R \# B$, where $(ad_c x_1)x_2 = x_1 x_2 - \chi_2(g_1^{-1})x_2 x_1$.

(ii) *If $\sqrt{\chi_1(g_2)}\sqrt{\chi_2(g_1)} = 1$ and $x_i g_j = \chi_i(g_j)g_j x_i$ for $i, j = 1, 2$, then*

$$\sqrt{\chi_2(g_1)}x_1 x_2 - \sqrt{\chi_1(g_2)}x_2 x_1$$

is a primitive element of R and

$$\sqrt{\chi_2(g_1)}x_1 x_2 - \sqrt{\chi_1(g_2)}x_2 x_1 - \beta(g_1 g_2 - 1)$$

is a $(1, g_1 g_2)$ -primitive element of biproduct $R \# B$ for any $\beta \in k$.

Proof. (i) Let $z := \sum_{m=0}^r (-1)^m \begin{bmatrix} r \\ m \end{bmatrix}_{\sqrt{\chi_1(g_1^{-1})}} x_1^{r-m} x_2 x_1^m$. By [6, Lemma A.1], $(ad_c x_1)^r x_2$ is primitive. However,

$$\begin{aligned} & (ad_c x_1)^r x_2 \\ &= \sum_{m=0}^r (-1)^m \begin{pmatrix} r \\ m \end{pmatrix}_{\chi_1(g_1^{-1})} \chi_1(g_1^{-1})^{\frac{1}{2}m(m-1)} \chi_2(g_1^{-1})^m x_1^{r-m} x_2 x_1^m \quad (\text{by [6, (A.8)]}) \\ &= \sum_{m=0}^r (-1)^m \begin{bmatrix} r \\ m \end{bmatrix}_{\sqrt{\chi_1(g_1^{-1})}} \chi_1(g_1^{-1})^{\frac{1}{2}m(r-1)} \chi_2(g_1^{-1})^m x_1^{r-m} x_2 x_1^m \quad (\text{by [6, P40]}) \\ &= \sum_{m=0}^r (-1)^m \begin{bmatrix} r \\ m \end{bmatrix}_{\sqrt{\chi_1(g_1^{-1})}} x_1^{r-m} x_2 x_1^m \quad (\text{by assumption (14.25)}). \end{aligned}$$

Thus z is primitive in R , i.e.

$$\Delta_R(z) = z \otimes 1 + 1 \otimes z. \quad (14.27)$$

Since

$$\begin{aligned}
\delta_R(z) &= \sum_{m=0}^r (-1)^m \begin{bmatrix} r \\ m \end{bmatrix}_{\sqrt{\chi_1(g_1^{-1})}} \delta_R(x_1^{r-m} x_2 x_1^m) \\
&= \sum_{m=0}^r (-1)^m \begin{bmatrix} r \\ m \end{bmatrix}_{\sqrt{\chi_1(g_1^{-1})}} g_1^r g_2 \otimes x_1^{r-m} x_2 x_1^m \\
&= g_1^r g_2 \otimes z,
\end{aligned}$$

we have

$$\begin{aligned}
\Delta_{R\#B}(z) &= \Delta_{R\#B}(z\#1) \\
&= (id_R \otimes \mu_B \otimes id_R \otimes id_B)(id_R \otimes id_B \otimes C_{R,B} \otimes id_B) \\
&\quad (id_R \otimes \delta_R \otimes \Delta_B)(1\#z \otimes 1) + z \otimes 1 \quad (\text{by Theorem 14.3.1}) \\
&= g_1^r g_2 \otimes z + z \otimes 1,
\end{aligned}$$

where $C_{R,B}$ denotes the ordinary twist map from $R \otimes B$ to $B \otimes R$ by sending $r \otimes b$ to $b \otimes r$ for any $r \in R, b \in B$.

(ii) See

$$\begin{aligned}
&\Delta_{R\#B}(\sqrt{\chi_2(g_1)}x_1x_2 - \sqrt{\chi_1(g_2)}x_2x_1) \\
&= \sqrt{\chi_2(g_1)}(g_1g_2 \otimes x_1x_2 + x_1g_2 \otimes x_2 + g_1x_2 \otimes x_1 + x_1x_2 \otimes 1) \\
&\quad - \sqrt{\chi_1(g_2)}(g_1g_2 \otimes x_2x_1 + x_2g_1 \otimes x_1 + g_2x_1 \otimes x_2 + x_2x_1 \otimes 1) \\
&= g_1g_2 \otimes (\sqrt{\chi_2(g_1)}x_1x_2 - \sqrt{\chi_1(g_2)}x_2x_1) + (\sqrt{\chi_2(g_1)}x_1x_2 - \sqrt{\chi_1(g_2)}x_2x_1) \otimes 1 \\
&\quad (\text{by assumption of (ii)})
\end{aligned}$$

and

$$\begin{aligned}
&\Delta_{R\#B}(\sqrt{\chi_2(g_1)}x_1x_2 - \sqrt{\chi_1(g_2)}x_2x_1) - \Delta_{R\#B}(\beta(g_1g_2 - 1)) \\
&= g_1g_2 \otimes (\sqrt{\chi_2(g_1)}x_1x_2 - \sqrt{\chi_1(g_2)}x_2x_1 - \beta(g_1g_2 - 1)) \\
&\quad + (\sqrt{\chi_2(g_1)}x_1x_2 - \sqrt{\chi_1(g_2)}x_2x_1 - \beta(g_1g_2 - 1)) \otimes 1.
\end{aligned}$$

Thus $\sqrt{\chi_2(g_1)}x_1x_2 - \sqrt{\chi_1(g_2)}x_2x_1 - \beta(g_1g_2 - 1)$ is a $(1, g_1g_2)$ -primitive element of $R\#B$. See

$$\begin{aligned}
&\Delta_R(\sqrt{\chi_2(g_1)}x_1x_2 - \sqrt{\chi_1(g_2)}x_2x_1) \\
&= \sqrt{\chi_2(g_1)}(1 \otimes x_1x_2 + x_1 \otimes x_2 + \chi_2(g_1)^{-1}x_2 \otimes x_1 + x_1x_2 \otimes 1) \\
&\quad - \sqrt{\chi_1(g_2)}(1 \otimes x_2x_1 + x_2 \otimes x_1 + \chi_1(g_2)^{-1}x_1 \otimes x_2 + x_2x_1 \otimes 1) \\
&= 1 \otimes (\sqrt{\chi_2(g_1)}x_1x_2 - \sqrt{\chi_1(g_2)}x_2x_1) + (\sqrt{\chi_2(g_1)}x_1x_2 - \sqrt{\chi_1(g_2)}x_2x_1) \otimes 1 \\
&\quad (\text{by assumption of (ii)}).
\end{aligned}$$

Thus $\sqrt{\chi_2(g_1)}x_1x_2 - \sqrt{\chi_1(g_2)}x_2x_1$ is a primitive element of R . $\square$

For an $\text{ESC}(G, \chi_i, g_i; i \in J)$, we give the follows notations:

(FL1) N is a set and $J = \cup_{s \in N} (J_s \cup J'_s)$ is a disjoint union.

(FL2) There exists a bijection $\sigma : J^{(1)} \rightarrow J^{(2)}$ such that $\sigma|_{J_u}$ is a bijection from J_u to J'_u for $u \in N$, where $J^{(1)} := \cup_{u \in N} J_u$ and $J^{(2)} := \cup_{u \in N} J'_u$.

(FL3) There exists a $J^{(1)} \times J^{(1)}$ -matrix $A = (a_{ij})$ with $a_{ii} = 2$ and non-positive integer a_{ij} for any $i, j \in J^{(1)}$ and $i \neq j$. For any $u \in N$, there exists an integer $d_i^{(u)}$ such that $d_i^{(u)} a_{ij} = d_j^{(u)} a_{ji}$ for any $i, j \in J_u$.

(FL4) For any $u \in N$, there exists $0 \neq q_u \in k$ such that $\chi_i(g_j) = q_u^{-2d_i^{(u)} a_{ij}}$, $\chi_{\sigma(i)}(g_j) = \chi_i^{-1}(g_j)$ and $g_{\sigma(j)} = g_j$ for $i, j \in J_u$.

(FL5) There exists $\xi_i \in G$ such that $\chi_{\sigma(i)}(\xi_j) = \chi_i^{-1}(\xi_j)$, $\xi_{\sigma(i)} := \xi_i^{-1}$ and $g_i = g_{\sigma(i)} := \xi_i^2$ for any $i, j \in J_u$, $u \in N$; there exists a positive integer r_{ij} such that $r_{\sigma(i), \sigma(j)} = r_{ij}$ for any $i, j \in J^{(1)}$ with $i \neq j$.

(FL6) $\chi_j(g_i) \chi_i(g_j) \chi_i(g_i)^{r_{ij}-1} = 1$, $\chi_i(\xi_j) = \chi_j(\xi_i)$, $\chi_i(g_i)^{\frac{1}{2}(r_{ij}-1)} \chi_j(g_i) = 1$, for any $i, j \in J_u$, $i \neq j$, $u \in N$.

(FL7) G is a free commutative group generated by generator set $\{\xi_i \mid i \in J^{(1)}\}$.

An $\text{ESC}(G, g_i, \chi_i; i \in J)$ is said to be a local FL-matrix type (see [6, P.4]) if (FL1)–(FL4) hold. An $\text{ESC}(G, g_i, \chi_i; i \in J)$ is said to be a local FL-type if (FL1), (FL2), (FL5) and (FL6) hold. An $\text{ESC}(G, g_i, \chi_i; i \in J)$ is said to be a local FL-free type if (FL1), (FL2), (FL5), (FL6) and (FL7) hold. An $\text{ESC}(G, g_i, \chi_i; i \in J)$ is said to be a local FL-quantum group type if (FL1)–(FL4) and (FL7) hold. If N only contains one element and $J^{(1)} = \{1, 2, \dots, n\}$, then we delete ‘local’ in the terms above.

Let $\text{ESC}(G, \chi_i, g_i; i \in J)$ be a local FL-free type. Let I be the ideal of $kQ^s(G, g_i, \chi_i; i \in J)$ generated by the following elements:

$$\chi_j(\xi_i) E_i E_j - \chi_i^{-1}(\xi_j) E_j E_i - \delta_{\sigma(i), j} \frac{g_i^2 - 1}{\chi_i(\xi_i) - \chi_i(\xi_i)^{-1}}, \text{ for } i \in J_u, j \in J'_u, u \in N; \quad (14.28)$$

$$\sum_{m=0}^{r_{ij}} (-1)^m \begin{bmatrix} r_{ij} \\ m \end{bmatrix}_{\chi_i(\xi_i^{-1})} E_i^{r_{ij}-m} E_j E_i^m, \quad (14.29)$$

for any $i, j \in J_u$ or $i, j \in J'_u, i \neq j$ and $r_{ij} - 1 < \text{ord}(\chi_i(g_i)); u \in N$.

Let U be the algebra generated by set $\{K_i, X_i \mid i \in J\}$ with relations

$$\begin{aligned} X_i X_j - X_j X_i &= \delta_{\sigma(i), j} \frac{K_i^2 - K_i^{-2}}{\chi_i(\xi_i) - \chi_i(\xi_i)^{-1}}, \text{ for any } i \in J_u, j \in J'_u; u \in N; \\ K_i K_{\sigma(i)} &= K_{\sigma(i)} K_i = 1 \text{ for any } i \in J^{(1)}; \\ \chi_j(\xi_i) K_i X_j &= X_j K_i, \quad K_i K_j - K_j K_i = 0, \text{ for any } i, j \in J; \\ \sum_{m=0}^{r_{ij}} (-1)^m \begin{bmatrix} r_{ij} \\ m \end{bmatrix}_{\chi_i(\xi_i^{-1})} X_i^{r_{ij}-m} X_j X_i^m &= 0, \\ \text{for } i, j \in J_u \text{ or } i, j \in J'_u, i \neq j \text{ and } r_{ij} - 1 < \text{ord}(\chi_i(g_i)); u \in N. \end{aligned} \quad (14.30)$$

The comultiplication, counit and antipode of U are defined by

$$\begin{aligned} \Delta(X_j) &= X_j \otimes K_{\sigma(j)} + K_j \otimes X_j, & S(X_j) &= -\chi_j(\xi_j) X_j, & \epsilon(X_i) &= 0, \\ \Delta(K_i) &= K_i \otimes K_i, & S(K_j) &= K_{\sigma(j)}, & \epsilon(K_i) &= 1, \\ \Delta(X_{\sigma(j)}) &= X_{\sigma(j)} \otimes K_{\sigma(j)} + K_j \otimes X_{\sigma(j)}, & S(X_{\sigma(j)}) &= -\chi_{\sigma(j)}(\xi_j) X_{\sigma(j)} \end{aligned} \quad (14.31)$$

for any $j \in J^{(1)}, i \in J$.

In fact, $K_{\sigma(j)} = K_j^{-1}$ in U for any $j \in J^{(1)}$.

Theorem 14.3.12. *Under notation above, if $\text{ESC}(G, g_i, \chi_i; i \in J)$ is a local FL-free type, then $kQ^s(G, g_i, \chi_i; i \in J)/I \cong U$ as Hopf algebras.*

Proof. For any $i, j \in J_u, u \in N$, see $\chi_{\sigma(i)}(\xi_{\sigma(j)}) = \chi_i(\xi_j) = \chi_j(\xi_i) = \chi_{\sigma(j)}(\xi_{\sigma(i)})$, $\chi_{\sigma(i)}(\xi_j) = \chi_i^{-1}(\xi_j) = \chi_j^{-1}(\xi_i) = \chi_j(\xi_{\sigma(i)})$, $\chi_i(\xi_{\sigma(j)}) = \chi_i(\xi_j)^{-1} = \chi_j(\xi_i)^{-1} = \chi_{\sigma(j)}(\xi_i)$. Therefore,

$$\chi_i(\xi_j) = \chi_j(\xi_i) \quad (14.32)$$

For any $i, j \in J_u \cup J'_u, u \in N$. Obviously, for $i, j \in J'_u, i \neq j, u \in N$, (FL6) holds.

We show this theorem by following several steps.

(i) There is a algebra homomorphism Φ from kQ^s to U such that $\Phi(\xi_i) = K_i$, $\Phi(a_{hg_i, h}^{(i)}) = \Phi(h)K_iX_i$ and $\Phi(a_{hg_{\sigma(i)}, h}^{(\sigma(i))}) = \Phi(h)K_iX_{\sigma(i)}$ for all $h \in G$ and $i \in J^{(1)}$. Indeed, define algebra homomorphism $\phi : kG \rightarrow U$ given by $\phi(\xi_i) = K_i$ for $i \in J$ and a k -linear map $\psi : kQ_1^c \rightarrow U$ by $\psi(a_{gg_j, g}^{(j)}) = \phi(g)K_jX_j$ and $\psi(a_{gg_{\sigma(j)}, g}^{(\sigma(j))}) = \phi(g)K_jX_{\sigma(j)}$ for any $j \in J^{(1)}, g \in G$. For any $g, h \in G, j \in J^{(1)}$, see

$$\psi(h \cdot a_{gg_j, g}^{(j)}) = \psi(a_{hgg_j, hg}^{(j)}) = \phi(hg)K_jX_j = \phi(h)\phi(g)K_jX_j = \phi(h)\psi(a_{gg_j, g}^{(j)}),$$

and

$$\begin{aligned} \psi(a_{gg_j, g}^{(j)} \cdot h) &= \chi_j(h)\psi(a_{hgg_j, hg}^{(j)}) = \chi_j(h)\phi(hg)K_jX_j \\ &= \phi(g)K_jX_j\phi(h) \quad (\text{since } X_j\phi(h) = \phi(h)\chi_j(h)X_j) \\ &= \psi(a_{gg_j, g}^{(j)})\phi(h). \end{aligned}$$

Similarly, $\psi(h \cdot a_{gg_{\sigma(j)}, g}^{(\sigma(j))}) = \phi(h)\psi(a_{gg_{\sigma(j)}, g}^{(\sigma(j))})$ and $\psi(a_{gg_{\sigma(j)}, g}^{(\sigma(j))} \cdot h) = \psi(a_{gg_{\sigma(j)}, g}^{(\sigma(j))})\phi(h)$. This implies that ψ is a kG -bimodule map from $(kQ_1^c, g_i, \chi_i; i \in J)$ to ${}_{\phi}U_{\phi}$. Using the universal property of tensor algebra over kG , we complete the proof.

(ii) $\Phi(I) = 0$. For any $i \in J_u, j \in J'_u$, see that

$$\begin{aligned} &\Phi(\chi_j(\xi_i)E_iE_j - \chi_i(\xi_j)^{-1}E_jE_i - \delta_{\sigma(i), j} \frac{g_i^2 - 1}{\chi_i(\xi_i) - \chi_i(\xi_i)^{-1}}) \\ &= \chi_j(\xi_i)K_iX_iK_j^{-1}X_j - \chi_i(\xi_j)^{-1}K_j^{-1}X_jK_iX_i - \delta_{\sigma(i), j} \frac{K_i^4 - 1}{\chi_i(\xi_i) - \chi_i(\xi_i)^{-1}} \\ &= K_iK_j^{-1}X_iX_j - K_j^{-1}K_iX_jX_i - \delta_{\sigma(i), j} \frac{K_i^4 - 1}{\chi_i(\xi_i) - \chi_i(\xi_i)^{-1}} \\ &= K_iK_j^{-1}(X_iX_j - X_jX_i - \delta_{\sigma(i), j} \frac{K_i^3K_j - K_i^{-1}K_j}{\chi_i(\xi_i) - \chi_i(\xi_i)^{-1}}) \\ &= K_iK_j^{-1}(X_iX_j - X_jX_i - \delta_{\sigma(i), j} \frac{K_i^2 - K_i^{-2}}{\chi_i(\xi_i) - \chi_i(\xi_i)^{-1}}) = 0. \end{aligned}$$

For $i, j \in J_u, i \neq j$, see that

$$\begin{aligned} &\Phi(E_i^{r_{ij}-m}E_jE_i^m) \\ &= (K_iX_i)^{r_{ij}-m}K_jX_j(K_iX_i)^m \\ &= \chi_i(\xi_i)^{\frac{1}{2}((r_{ij}-m)(r_{ij}-m-1)+m(m-1)+2(r_{ij}-m)m)}\chi_i(\xi_j)^{r_{ij}-m}\chi_j(\xi_i)^mX_i^{r_{ij}-m}X_jX_i^m \\ &= \chi_i(\xi_i)^{\frac{1}{2}(r_{ij}-1)r_{ij}}\chi_i(\xi_j)^{r_{ij}}X_i^{r_{ij}-m}X_jX_i^m \quad (\text{by (FL6)}) \end{aligned}$$

and

$$\Phi\left(\sum_{m=0}^{r_{ij}}(-1)^m \begin{bmatrix} r_{ij} \\ m \end{bmatrix}_{\chi_i(\xi_i^{-1})} E_i^{r_{ij}-m} E_j E_i^m\right) = 0.$$

Similarly, the equation above holds for $i, j \in J'_u$, $i \neq j$, $u \in N$.

By (ii), there exists an algebra homomorphism $\bar{\Phi}$ from kQ^s/I to U such that $\bar{\Phi}(x+I) = \Phi(x)$ for any $x \in kQ^s$. For convenience, we will still use x to denote $x+I$.

(iii) It follows from the definition of U that there exists a unique algebra map $\Psi : U \rightarrow kQ^s/I$ such that $\Psi(K_i) = \xi_i$, $\Psi(X_i) = \xi_i^{-1}E_i$ and $\Psi(X_{\sigma(i)}) = \xi_i^{-1}E_{\sigma(i)}$ for all $i \in J^{(1)}$. It is easy to see that $\Psi\bar{\Phi} = id$ and $\bar{\Phi}\Psi = id$.

(iv) We show that I is a Hopf ideal of kQ^s . It follows from Theorem 14.3.7 that $R := \text{diag}(kQ^s)$ is a braided Hopf algebra with $\delta^-(E_i) = g_i \otimes E_i$, $h \triangleright E_i = \chi_i^{-1}(h)E_i$ for any $h \in G$, $i \in J$. By Theorem 14.3.1, $R \# B \cong kQ^s$ as Hopf algebras. It follows from Lemma 14.3.11 that I is a Hopf ideal of kQ^s .

Obviously, Ψ preserve the comultiplication and counit. Since kQ^s/I is a Hopf algebra, then Ψ is a Hopf algebra isomorphism. $\square$

Corollary 14.3.13. *The quantum enveloping algebra of a complex semisimple Lie algebra is isomorphic to a quotient of a semi-path Hopf algebra as Hopf algebras.*

Proof. Let k be the complex field and L a complex semisimple Lie algebra determined by $A = (a_{ij})_{n \times n}$. So A is a symmetrizable Cartan matrix with $d_i \in \{1, 2, 3\}$ such that $d_i a_{ij} = d_j a_{ji}$ for any $i, j \in J^{(1)} = \{1, 2, \dots, n\}$. Let $|N| = 1$, $J^{(2)} = \{n+1, n+2, \dots, n+n\}$, $J = J^{(1)} \cup J^{(2)}$ and $\sigma : J^{(1)} \rightarrow J^{(2)}$ by sending i to $i+n$. Let G be a free commutative group generated by generator set $\{\xi_i \mid i \in J^{(1)}\}$. Set $\xi_{\sigma(i)} = \xi_i^{-1}$, $g_i = g_{\sigma(i)} := \xi_i^2$, $r_{ij} = r_{\sigma(i), \sigma(j)} = 1 - a_{ij}$ for any $i, j \in J^{(1)}$, $i \neq j$. Define $\chi_i(\xi_j) = q^{-d_i a_{ij}}$ and $\chi_{\sigma(i)}(\xi_j) = \chi_i^{-1}(\xi_j)$ for any $i, j \in J^{(1)}$, where q is not a root of 1 with $0 \neq q \in k$. It is easy to check (FL1)–(FL7) hold, i.e. $\text{ESC}(G, g_i, \chi_i; i \in J)$ is an FL-quantum group type. U in Theorem 14.3.12 exactly is the quantum enveloping algebra $U_q(L)$ of L (see [87, p.218] or [72]). Therefore the conclusion follows from Theorem 14.3.12. $\square$

In fact, for any a generalized Cartan matrix A , we can obtain an FL-quantum group type $\text{ESC}(G, g_i, \chi_i; i \in J)$ as in the proof above. By the way, if $\text{ESC}(G, g_i, \chi_i; i \in J)$ is a Local FL-type, then there exist the Hopf ideals, which generated by (14.28) and (14.29), in co-path Hopf algebra $kQ^c(G, g_i, \chi_i; i \in J)$ and multiple Taft algebra $kQ^c(G, g_i, \chi_i; i \in J)$, respectively.

14.4 Classification of ramification systems for the symmetric group

Unless specified otherwise, in the section we have the following assumptions and notations. n is a positive integer with $n \neq 6$; F is a field containing a primitive n -th root of 1; $G = S_n$, the symmetric group; $\text{Aut}G$ and $\text{Inn}G$ denote the automorphism group and inner automorphism group, respectively; S_I denotes the group of full permutations of set I ; $Z(G)$ denotes the center of G and Z_x the centralizer of x ; 1 denotes the unity element of G ; $\widehat{G}$ denotes the set of characters

of all one-dimensional representations of G . ϕ_h denotes the inner automorphism induced by h given by $\phi_h(x) = h x h^{-1}$, for any $x \in G$. C_n is cyclic group of order n ; D^B denotes the cartesian product $\prod_{i \in B} D_i$, where $D_i = D$ for any $i \in B$.

Proposition 14.4.1. (i) *There are $\frac{n!}{\lambda_1! \lambda_2! \dots \lambda_n! 1^{\lambda_1} 2^{\lambda_2} \dots n^{\lambda_n}}$ permutations of type $1^{\lambda_1} 2^{\lambda_2} \dots n^{\lambda_n}$ in S_n ;*

(ii) *If $n \neq 6$, then $\text{Aut}(S_n) = \text{Inn}(S_n)$; If $n \neq 2, 6$, then $\text{Inn}(S_n) \cong S_n$;*

(iii) *$S'_n = A_n$, where A_n is the alternating group. In addition,*

$$S_n/A_n \cong \begin{cases} C_2 & n \geq 2 \\ C_1 & n = 1 \end{cases}.$$

Proof. (i) It follows from [57, Exercise 2.7.7].

(ii) By [141, Theorem 1.12.7], $\text{Aut}(A_n) \cong S_n$ when $n > 3$ and $n \neq 6$. By [56, Theorem 1.6.10], A_n is a non-commutative simple group when $n \neq 4$ and $n > 1$. Thus it follows from [141, Theorem 1.12.6] that $\text{Aut}(S_n) = \text{Inn}(S_n) \cong S_n$ when $n \neq 6$ and $n > 4$. Applying [141, Example in Page 100], we obtain $\text{Aut}(S_n) = \text{Inn}(S_n) \cong S_n$ when $n = 3, 4$. Obviously, $\text{Aut}(S_n) = \text{Inn}(S_n)$ when $n = 1, 2$.

(iii) It follows from [122, Theorem 2.3.10]. $\square$

Definition 14.4.2. *A group G is said to act on a non-empty set Ω , if there is a map $G \times \Omega \rightarrow \Omega$, denoted by $(g, x) \mapsto g \circ x$, such that for all $x \in \Omega$ and $g_1, g_2 \in G$:*

$$(g_1 g_2) \circ x = g_1 \circ (g_2 \circ x) \quad \text{and} \quad 1 \circ x = x,$$

where 1 denote the unity element of G .

For each $x \in \Omega$, let

$$G_x := \{g \circ x \mid g \in G\},$$

called the orbit of G on Ω . For each $g \in G$, let

$$F_g := \{x \in \Omega \mid g \circ x = x\},$$

called the fixed point set of g .

Burnside's lemma, sometimes also called Burnside's counting theorem, which is useful to compute the number of orbits.

Theorem 14.4.3. (See [57, Theorem 2.9.3.1]) (Burnside's Lemma) *Let G be a finite group that acts on a finite set Ω , then the number $\mathcal{N}$ of orbits is given by the following formula:*

$$\mathcal{N} = \frac{1}{|G|} \sum_{g \in G} |F_g|.$$

Definition 14.4.4. *Let G be a group. A character of a one-dimensional representation of G is a homomorphism from G to $F - \{0\}$, that is, a map $\chi : G \rightarrow F$ such that $\chi(xy) = \chi(x)\chi(y)$ and $\chi(1) = 1$ for any $x, y \in G$. The set of all characters of all one-dimensional representations of G is denoted by $\hat{G}$.*

For convenience, a character of a one-dimensional representation of G is called a character of G in short. [22, Page 36, example 4] indicated that there is a one-to-one correspondence between characters of G and characters of G/G' .

Proposition 14.4.5. *Let G be a finite abelian group and F contains $|G|$ th primitive root of 1, then there are exactly $|G|$ characters of G , i.e. $|\widehat{G}| = |G|$.*

Corollary 14.4.6. *Let G be a finite group and F contains $|G|$ th primitive root of 1, then there are exactly $|G/G'|$ characters of G , i.e. $|\widehat{G}| = |G/G'|$.*

14.4.1 Ramification Systems and Isomorphisms

We begin with defining ramification and ramification system.

Definition 14.4.7. *Let G be a finite group and $\mathcal{K}(G)$ the set of all conjugate classes of G . r is called a ramification data or ramification of G if $r = \sum_{C \in \mathcal{K}(G)} r_C C$, where r_C is a non-negative integer for any C in $\mathcal{K}(G)$. For convenience, we choose a set $I_C(r)$ such that $r_C = |I_C(r)|$ for any C in $\mathcal{K}(G)$.*

Let $\mathcal{K}_r(G) = \{C \in \mathcal{K}(G) \mid r_C \neq 0\} = \{C \in \mathcal{K}(G) \mid I_C(r) \neq \emptyset\}$.

Definition 14.4.8. *$(G, r, \overrightarrow{\chi}, u)$ is called a ramification system with characters (or RSC in short), if r is a ramification of G , u is a map from $\mathcal{K}(G)$ to G with $u(C) \in C$ for any $C \in \mathcal{K}(G)$, and $\overrightarrow{\chi} = \{\chi_C^{(i)}\}_{i \in I_C(r), C \in \mathcal{K}_r(G)} \in \prod_{C \in \mathcal{K}_r(G)} (\widehat{Z_{u(C)}})^{r_C}$.*

Definition 14.4.9. *$RSC(G, r, \overrightarrow{\chi}, u)$ and $RSC(G', r', \overrightarrow{\chi'}, u')$ are said to be isomorphic (denoted $RSC(G, r, \overrightarrow{\chi}, u) \cong RSC(G', r', \overrightarrow{\chi'}, u')$) if the following conditions are satisfied:*

- (i) *There exists a group isomorphism $\phi : G \rightarrow G'$;*
- (ii) *For any $C \in \mathcal{K}(G)$, there exists an element $h_C \in G$ such that $\phi(h_C u(C) h_C^{-1}) = u'(\phi(C))$;*
- (iii) *For any $C \in \mathcal{K}_r(G)$, there exists a bijective map $\phi_C : I_C(r) \rightarrow I_{\phi(C)}(r')$ such that $\chi_{\phi(C)}^{(\phi_C(i))}(\phi(h_C h h_C^{-1})) = \chi_C^{(i)}(h)$, for all $h \in Z_{u(C)}$, for all $i \in I_C(r)$.*

Remark. Obviously, if $u(C) = u'(C)$ for any $C \in \mathcal{K}_r(G)$, then $RSC(G, r, \overrightarrow{\chi}, u) \cong RSC(G, r, \overrightarrow{\chi}, u')$. In fact, let $\phi = id_G$ and $\phi_C = id_{I_C(r)}$ for any $C \in \mathcal{K}_r(G)$. It is straightforward to check the conditions in Definition 14.4.9 hold.

Lemma 14.4.10. *Given an $RSC(G, r, \overrightarrow{\chi}, u)$. If $RSC(G', r', \overrightarrow{\chi'}, u') \cong RSC(G, r, \overrightarrow{\chi}, u)$, then there exists a group isomorphism $\phi : G \rightarrow G'$ such that*

- (i) $r'_{\phi(C)} = r_C$ for any $C \in \mathcal{K}(G)$;
- (ii) $\mathcal{K}_{r'}(G') = \phi(\mathcal{K}_r(G)) = \{\phi(C) \mid C \in \mathcal{K}_r(G)\}$.

Proof. (i) By Definition 14.4.9, there exists a group isomorphism $\phi : G \rightarrow G'$ such that $r'_{\phi(C)} = r_C$ for any $C \in \mathcal{K}_r(G)$. If $C \notin \mathcal{K}_r(G)$, i.e. $r_C = 0$, then $r'_{\phi(C)} = r_C = 0$;
(ii) is an immediate consequence of (i). $\square$

Proposition 14.4.11. *Given a $RSC(G, r, \vec{\chi}, u)$, then $RSC(G, r', \vec{\chi}', u') \cong RSC(G, r, \vec{\chi}, u)$ if and only if there exists a group isomorphism $\phi \in \text{Aut}G$ such that:*

- (i) $r' = \sum_{C \in \mathcal{K}(G)} r_{\phi^{-1}(C)} C$;
- (ii) $\chi'_{\phi(C)} = \chi_C^{(\phi_C^{-1}(i))} (\phi \phi_{h_C})^{-1}$ and $u'(\phi(C)) = \phi \phi_{h_C}(u(C))$ for any $C \in \mathcal{K}_r(G)$, $i \in I_C(r)$, $\phi \in \text{Aut}G$ and $\phi_C \in S_{r_C}$, $h_C \in G$.

Proof. If $RSC(G, r', \vec{\chi}', u') \cong RSC(G, r, \vec{\chi}, u)$, then, by Lemma 14.4.10(i), there exists $\phi \in \text{Aut}G$ such that

$$r' = \sum_{C \in \mathcal{K}(G)} r'_C C = \sum_{C \in \mathcal{K}(G)} r'_{\phi(C)} \phi(C) = \sum_{C \in \mathcal{K}(G)} r_C \phi(C) = \sum_{C \in \mathcal{K}(G)} r_{\phi^{-1}(C)} C.$$

This shows Part (i).

By Definition 14.4.9 (ii), there exists $h_C \in G$ such that $u'(\phi(C)) = \phi(h_C u(C) h_C^{-1}) = \phi \phi_{h_C}(u(C))$ for any $C \in \mathcal{K}_r(G)$; By Definition 14.4.9 (iii), there exists $\phi_C \in S_{I_C(r)} = S_{r_C}$ such that $\chi'_{\phi(C)}(\phi(h_C h h_C^{-1})) = \chi'_{\phi(C)}(\phi \phi_{h_C}(h)) = \chi_C^{(i)}(h)$ for any $h \in Z_{u(C)}$. Consequently,

$$\chi'_{\phi(C)}(\phi_C^{(i)}) = \chi_C^{(i)} (\phi \phi_{h_C})^{-1},$$

i.e.

$$\chi'_{\phi(C)} = \chi_C^{(\phi_C^{-1}(i))} (\phi \phi_{h_C})^{-1},$$

where $C \in \mathcal{K}_r(G)$. That is, (ii) holds.

Conversely, if (i) and (ii) hold, let

$$r' = \sum_{C \in \mathcal{K}(G)} r_{\phi^{-1}(C)} C$$

and

$$I_{\phi(C)}(r') = I_C(r)$$

for any $C \in \mathcal{K}(G)$, $\phi \in \text{Aut}G$. Obviously, $\mathcal{K}_{r'}(G) = \{\phi(C) \mid C \in \mathcal{K}_r(G)\}$. For any $C \in \mathcal{K}_r(G)$, $h_C \in G$, $\phi_C \in S_{I_C(r)} = S_{r_C}$, let

$$u'(\phi(C)) = \phi(h_C u(C) h_C^{-1})$$

and

$$\chi'_{\phi(C)} = \chi_C^{(\phi_C^{-1}(i))} (\phi \phi_{h_C})^{-1}$$

for any $i \in I_{\phi(C)}(r')$. It is easy to check $RSC(G, r', \vec{\chi}', u') \cong RSC(G, r, \vec{\chi}, u)$. $\square$

Corollary 14.4.12. *If $G = S_n$ with $n \neq 6$, then $RSC(G, r, \vec{\chi}, u) \cong RSC(G, r', \vec{\chi}', u')$ if and only if*

- (i) $r' = r$;
- (ii) $\chi'_C = \chi_C^{(\phi_C^{-1}(i))} \phi_{g_C}^{-1}$ and $u'(C) = \phi_{g_C}(u(C))$ for any $C \in \mathcal{K}_r(G)$, $i \in I_C(r)$, $\phi \in \text{Aut}G$ and $g_C \in G$.

Proof. By Proposition 14.4.1(ii), $\text{Aut}G = \text{Inn}G$ and $\phi(C) = C$ for any $\phi \in \text{Aut}G, C \in \mathcal{K}(G)$. There exists $g_C \in G$ such that $\phi\phi_{h_C} = \phi_{g_C}$ since $\phi\phi_{h_C} \in \text{Aut}G = \text{Inn}G$ for any $h_C \in G, \phi \in \text{Aut}G, C \in \mathcal{K}(G)$. Considering Proposition 14.4.11, we complete the proof. $\square$

From now on, we suppose that $G = S_n (n \neq 6)$. For a given ramification r of G , let $\Omega(G, r)$ be the set of all RSC's of G with the ramification r , namely, $\Omega(G, r) := \{(G, r, \vec{\chi}, u) \mid (G, r, \vec{\chi}, u) \text{ is an RSC}\}$. Let $\mathcal{N}(G, r)$ be the number of isomorphic classes in $\Omega(G, r)$. This article is mostly devoted to investigate the formula of $\mathcal{N}(G, r)$.

14.4.2 Action of Group

Denote $\text{RSC}(G, r, \vec{\chi}, u)$ by $\left(\left\{\chi_C^{(i)}\right\}_i, u(C)\right)_{C \in \mathcal{K}_r(G)}$ for short. Let $M := \prod_{C \in \mathcal{K}_r(G)} (\text{Aut}G \times S_{I_C(r)})$
 $= \prod_{C \in \mathcal{K}_r(G)} (\text{Inn}G \times S_{r_C})$. Define the action of M on $\Omega(G, r)$ as follows:

$$(\phi_{g_C}, \phi_C)_C \circ \left(\left\{\chi_C^{(i)}\right\}_i, u(C)\right)_C = \left(\left\{\chi_C^{(\phi_C^{-1}(i))} \phi_{g_C}^{-1}\right\}_i, \phi_{g_C} u(C)\right)_C \quad (14.33)$$

In term of Corollary 14.4.12, each orbit of $\Omega(G, r)$ represents an isomorphic class of RSC's. As a result, $\mathcal{N}(G, r)$ is equal to the number of orbits in $\Omega(G, r)$.

We compute the the number of orbits by following steps.

14.4.3 The structure of centralizer in S_n

We recall the semidirect product and the wreath product of groups (see [22, Page 21], [111, Page 268-272],).

Definition 14.4.13. Let N and K be groups, and assume that there exists a group homomorphism $\alpha : K \rightarrow \text{Aut}N$. The semidirect product $N \rtimes_{\alpha} K$ of N and K with respect to α is defined as follows:

- (1) As set, $N \rtimes_{\alpha} K$ is the Cartesian product of N and K ;
- (2) The multiplication is given by

$$(a, x)(b, y) = (a\alpha(x)(b), xy)$$

for any $a \in N$ and $x \in N$.

Remark. Let $L = N \rtimes_{\alpha} K$. If we set

$$\bar{N} = \{(a, 1) \mid a \in N\}, \bar{K} = \{(1, x) \mid x \in K\},$$

then $\bar{N}$ and $\bar{K}$ are subgroups of L which are isomorphic to N and K , respectively, and will be identified with N and K .

Definition 14.4.14. Let A and H be two groups and H act on the set X . Assume that B is the set of all maps from X to A . Define the multiplication of B and the group homomorphism α from H to $\text{Aut} B$ as follows:

$$(bb')(x) = b(x)b'(x) \quad \text{and} \quad \alpha(h)(b)(x) = b(h^{-1} \cdot x)$$

for any $x \in X, h \in H, b, b' \in B$.

The semidirect product $B \rtimes_{\alpha} H$ is called the (general) wreath product of A and H , written as $A \wr H$.

Remark. [111, Page 272] indicated that in Definition 14.4.14 $B \cong \prod_{x \in X} A_x$, where $A_x = A$ for any $x \in X$. Hence $W = A \wr H = B \rtimes H = (\prod_{x \in X} A_x) \rtimes H$. In particular, the wreath product plays an important role in the structure of centralizer in S_n when $H = S_n$ and $X = \{1, 2, \dots, n\}$.

Now we keep on the work in [111, Page 295-299]. Let Y_i be the set of all letters which belong to those cycles of length i in the independent cycle decomposition of σ . Clearly, $Y_i \cap Y_j = \emptyset$ for $i \neq j$ and $\bigcup_i Y_i = \{1, 2, \dots, n\}$.

Lemma 14.4.15. $\rho \in Z_{\sigma}$ if and only if Y_i is ρ -invariant, namely $\rho(Y_i) \subset Y_i$, and the restriction ρ_i of ρ on Y_i commutes with the restriction σ_i of σ on Y_i for $i = 1, 2, \dots, n$.

Proof. It follows from [111, Page 295]. $\square$

Proposition 14.4.16. If σ and σ_i is the same as above, then

- (i) $Z_{\sigma} = \prod_i Z_{\sigma_i}$;
- (ii) $Z_{\sigma_i} \cong C_i \wr S_{\lambda_i} \cong (C_i)^{\lambda_i} \rtimes S_{\lambda_i}$, where Z_{σ_i} is the centralizer of σ_i in S_{Y_i} ; $(C_i)^{\lambda_i} = \overbrace{C_i \times \dots \times C_i}^{\lambda_i}$; define $Z_{\sigma_i} = 1$ and $S_{\lambda_i} = 1$ when $\lambda_i = 0$.

Proof. (i) It follows from Lemma 14.4.15.

(ii) Assume

$$\tau = (a_{10}a_{11} \dots a_{1,l-1})(a_{20} \dots a_{2,l-1}) \dots (a_{m0} \dots a_{m,l-1})$$

and

$$X = \{1, 2, \dots, m\}, \quad Y = \{a_{ij} \mid 1 \leq i \leq m, 0 \leq j \leq l-1\}.$$

Obviously, in order to prove (ii), it suffices to show that $Z_{\tau} \cong C_l \wr S_m$, where Z_{τ} is the centralizer of τ in S_Y . The second index of a_{ij} ranges over $\{0, 1, \dots, l-1\}$. It is convenient to identify this set with $C_l = \mathbb{Z}/(l)$, and to use notation such as $a_{i,l+j} = a_{ij}$. We will define an isomorphism φ from Z_{τ} onto $C_l \wr S_m$. For any element ρ of Z_{σ} , we will define a permutation $\theta(\rho) \in S_m$ and a function $f(\rho)$ from X into C_l by

$$\rho^{-1}(a_{i0}) = a_{jk}, \quad j = \theta(\rho)^{-1}(i), \quad f_{\rho}(i) = g^k,$$

where g is the generator of C_l , $1 \leq i \leq m$. Let

$$\varphi(\rho) = (f_{\rho}, \theta(\rho)).$$

Then φ maps Z_{σ} into $C_l \wr S_m$. First we show that φ is surjective. In fact, for any $(f, \theta) \in C_l \wr S_m$ and $i \in X$, if $f(i) = g^k$, $\theta(j) = i$, then there exists a permutation ρ such that $\rho^{-1}(a_{ir}) = a_{j,k+r}$. It is easy to verify that $\rho \in Z_{\sigma}$ and $\varphi(\rho) = (f, \theta)$. Next we show that φ is injective. If $\rho \in \text{Ker} \varphi$, then $\theta(\rho) = 1$ and $f_{\rho}(i) = g^0$, this means that $\rho(a_{i0}) = a_{i0}$ for all i .

Hence, $\rho(a_{ij}) = a_{ij}$ for all i and j and $\rho = 1_Y$, which implies that φ is bijective. We complete the proof if φ is homomorphism, which is proved below. For any $\rho, \pi \in Z_\tau$, suppose that

$$(\rho\pi)^{-1}(a_{i0}) = \pi^{-1}(a_{jk}) = a_{s,t+k},$$

where $j = \theta(\rho)^{-1}(i), g^k = f_\rho(i), s = \theta(\pi)^{-1}(j) = \theta(\pi)^{-1}\theta(\rho)^{-1}(i), g^t = f_\pi(j) = f_\pi(\theta(\rho)^{-1}(i))$. Consequently,

$$\theta(\rho\pi)^{-1}(i) = s = (\theta(\rho)\theta(\pi))^{-1}(i), \quad f_{\rho\pi}(i) = g^{t+k} = f_\rho(i)f_\pi(j) = f_\rho(i)f_\pi(\theta(\rho)^{-1}(i)).$$

On the other hand, in view of Definition 14.4.14, $\theta(\rho)(f_\pi)(i) = f_\pi(\theta(\rho)^{-1}(i)) = f_{\rho\pi}(i)$, whence

$$(f_\rho, \theta(\rho))(f_\pi, \theta(\pi)) = (f_\rho\theta(\rho)(f_\pi), \theta(\rho)\theta(\pi)) = (f_{\rho\pi}, \theta(\rho\pi)).$$

It follows that

$$\varphi(\rho\pi) = \varphi(\rho)\varphi(\pi),$$

which is just what we need. $\square$

Corollary 14.4.17. (i) $|Z_\sigma| = \lambda_1!\lambda_2!\cdots\lambda_n!1^{\lambda_1}2^{\lambda_2}\cdots n^{\lambda_n}$, where σ is the same as above;
(ii) $\sum_{h \in C} |Z_h| = |G| = n!$ for any $C \in \mathcal{K}(G)$,

Proof. (i) It follows from 14.4.16 that

$$|Z_\sigma| = \prod_i |Z_{\sigma_i}| = \prod_i |(C_i)^{\lambda_i} \rtimes S_{\lambda_i}| = \prod_i |(C_i)^{\lambda_i}| |S_{\lambda_i}| = \prod_i (i^{\lambda_i} \lambda_i!).$$

(ii) Assume the type of C is $1^{\lambda_1}2^{\lambda_2}\cdots n^{\lambda_n}$. Combining (i) and Proposition 14.4.1(i), we have

$$\sum_{h \in C} |Z_h| = (\lambda_1!\lambda_2!\cdots\lambda_n!1^{\lambda_1}2^{\lambda_2}\cdots n^{\lambda_n}) \frac{n!}{\lambda_1!\lambda_2!\cdots\lambda_n!1^{\lambda_1}2^{\lambda_2}\cdots n^{\lambda_n}} = n! . \quad \square$$

Proposition 14.4.16 shows that Z_σ is a direct product of some wreath products such as $C_l \wr S_m$. Denote by $((b_i)_i, \sigma)$ the element of $C_l \wr S_m = (C_l)^m \rtimes S_m$ where $b_i \in C_l, \sigma \in S_m$. By Definition 14.4.13 and 14.4.14, we have

$$\begin{aligned} ((b_i)_i, \sigma)((b'_i)_i, \sigma') &= ((b_i)_i \cdot \sigma((b'_i)_i), \sigma\sigma') \\ &= ((b_i)_i \cdot (b'_{\sigma^{-1}(i)})_i, \sigma\sigma') \\ &= ((b_i b'_{\sigma^{-1}(i)})_i, \sigma\sigma') \end{aligned}$$

and

$$((b_i)_i, \sigma)^{-1} = ((b_{\sigma(i)}^{-1})_i, \sigma^{-1}).$$

14.4.4 Characters of Z_σ

Lemma 14.4.18. *If $H = H_1 \times \cdots \times H_r$, then*

$$(i) \ H' = H'_1 \times \cdots \times H'_r;$$

$$(ii) \ H/H' \cong \prod_{i=1}^r H_i/H'_i.$$

Proof. (i) It follows from [141, Page 77];

(ii) It follows from [56, Corollary 8.11] and (i). $\square$

According to Proposition 14.4.16(i) and Lemma 14.4.18(ii), to investigate the structure of $Z_{\sigma_i}/Z'_{\sigma_i}$, we have to study $(C_l \wr S_m)/(C_l \wr S_m)'$.

Lemma 14.4.19. *Let $B = (C_l)^m$, $\bar{B} = \left\{ (b_i)_i \in B \mid \prod_{i=1}^m b_i = 1 \right\}$ and $W = C_l \wr S_m = B \rtimes S_m$. Then $W' = \bar{B} \rtimes S'_m = \bar{B} \rtimes A_m$.*

Proof. Obviously, $\bar{B} \triangleleft B$.

For any $((b_i)_i, \sigma), ((b'_i)_i, \sigma') \in W$, see

$$\begin{aligned} & ((b_i)_i, \sigma)^{-1} ((b'_i)_i, \sigma')^{-1} ((b_i)_i, \sigma) ((b'_i)_i, \sigma') \\ &= ((b_{\sigma(i)}^{-1})_i, \sigma^{-1}) ((b'_{\sigma'(i)}^{-1})_i, \sigma'^{-1}) ((b_i b'_{\sigma^{-1}(i)})_i, \sigma \sigma') \\ &= ((b_{\sigma(i)}^{-1} b'_{\sigma' \sigma(i)}^{-1})_i, \sigma^{-1} \sigma'^{-1}) ((b_i b'_{\sigma^{-1}(i)})_i, \sigma \sigma') \\ &= ((b_{\sigma(i)}^{-1} b'_{\sigma' \sigma(i)}^{-1} b'_{\sigma' \sigma(i)} b'_{\sigma^{-1} \sigma' \sigma(i)})_i, \sigma^{-1} \sigma'^{-1} \sigma \sigma'). \end{aligned}$$

Considering $\prod_i b_{\tau(i)} = \prod_i b_i$ for any $\tau \in S_m$, we have

$$\begin{aligned} & \prod_i (b_{\sigma(i)}^{-1} b'_{\sigma' \sigma(i)}^{-1} b'_{\sigma' \sigma(i)} b'_{\sigma^{-1} \sigma' \sigma(i)}) \\ &= \prod_i b_i^{-1} \prod_i b_i'^{-1} \prod_i b_i \prod_i b_i' \\ &= 1 \end{aligned}$$

and $b_{\sigma(i)}^{-1} b'_{\sigma' \sigma(i)}^{-1} b'_{\sigma' \sigma(i)} b'_{\sigma^{-1} \sigma' \sigma(i)} \in \bar{B}$. By Proposition 14.4.1(iii), $\sigma^{-1} \sigma'^{-1} \sigma \sigma' \in S'_m = A_m$. Thus

$$((b_i)_i, \sigma)^{-1} ((b'_i)_i, \sigma')^{-1} ((b_i)_i, \sigma) ((b'_i)_i, \sigma') \in \bar{B} \rtimes A_m,$$

which implies $W' \subset \bar{B} \rtimes A_m$.

Conversely, for any $b = ((b_i)_i, 1) \in \bar{B}$, then $\prod_{i=1}^m b_i = 1$. Let $\sigma = (1, 2, \dots, m)$, $b'_1 = 1$, $b'_i = \prod_{j=1}^{i-1} b_j$ ($2 \leq i \leq m$), See

$$\begin{aligned} ((b'_i)_i, 1)^{-1} (1, \sigma)^{-1} ((b'_i)_i, 1) (1, \sigma) &= ((b_i'^{-1})_i, 1) (1, \sigma^{-1}) ((b'_i)_i, \sigma) \\ &= ((b_i'^{-1}, b'_{\sigma(i)})_i, 1) \\ &= ((\prod_{j=1}^{i-1} b_j^{-1} \prod_{j=1}^i b_j)_i, 1) \\ &= ((b_i)_i, 1) = b. \end{aligned}$$

Therefore, b is a commutator of $((b'_i)_i, 1)$ and $(1, \sigma)$, i.e. $b \in W'$. This show $\bar{B} \subset W'$.

Furthermore, for any $\sigma, \sigma' \in S_m$, we have

$$(1, \sigma)^{-1}(1, \sigma')^{-1}(1, \sigma)(1, \sigma') = (1, \sigma^{-1}\sigma'^{-1}\sigma\sigma'),$$

which implies $(1, \sigma^{-1}\sigma'^{-1}\sigma\sigma') \in W'$ and $S'_m = A_m \subset W'$.

Consequently, $\bar{B} \rtimes A_m \subset W'$. $\square$

Proposition 14.4.20. (i) Let B and $\bar{B}$ be the same as in Lemma 14.4.19. Then $B/\bar{B} \cong C_l$;

(ii) Let W and W' be the same as in Lemma 14.4.19. Then

$$W/W' \cong \begin{cases} C_l \times C_2 & m \geq 2 \\ C_l & m = 1 \\ 1 & m = 0 \end{cases};$$

(iii) Let σ be the same as in Proposition 14.4.16. Then

$$Z_\sigma/Z'_\sigma \cong \left(\prod_{\lambda_i=1} C_i \right) \left(\prod_{\lambda_i \geq 2} (C_i \times C_2) \right).$$

Proof. (i) It is clear that $\nu : B \rightarrow C_l$ by sending $(b_i)_i$ to $\prod_i b_i$ is a group homomorphism with $\text{Ker}\nu = \bar{B}$. Thus (i) holds.

(ii) By (i) and Proposition 14.4.1(iii), we have to show $W/W' \cong (B/\bar{B}) \times (S_m/A_m)$. Define

$$\begin{aligned} \psi : W &\rightarrow (B/\bar{B}) \times (S_m/A_m) \\ (b, \sigma) &\mapsto (\bar{b}, \bar{\sigma}), \end{aligned}$$

where $\bar{b}$ and $\bar{\sigma}$ are the images of b and σ under canonical epimorphisms $B \rightarrow B/\bar{B}$ and $S_m \rightarrow S_m/A_m$, respectively. Therefore

$$\psi((b', \sigma')(b, \sigma)) = (\overline{b'\sigma'(b)}, \overline{\sigma'\sigma}) = (\bar{b}'\overline{\sigma'(b)}, \bar{\sigma}'\bar{\sigma}).$$

Let $b = (b_i)_i$. Since $\nu(\sigma'(b)) = \nu((b_{\sigma'^{-1}(i)})_i) = \prod_i b_{\sigma'^{-1}(i)} = \prod_i b_i = \nu(b)$, we have $\overline{\sigma'(b)} = \bar{b}$. Consequently,

$$\psi((\sigma', b')(\sigma, b)) = (\bar{b}'\overline{\sigma(b)}, \bar{\sigma}'\bar{\sigma}) = (\bar{b}'\bar{b}, \bar{\sigma}'\bar{\sigma}) = \psi((b', \sigma'))\psi((b, \sigma)).$$

This show that ψ is a group homomorphism. Obviously, it is surjective and $\text{Ker}\psi = \bar{B} \rtimes A_m = W'$. we complete the proof of (ii).

(iii) By Proposition 14.4.16(i) and Lemma 14.4.18(ii), $Z_\sigma/Z'_\sigma = (\prod_i Z_{\sigma_i})/(\prod_i Z'_{\sigma_i}) \cong \prod_i (Z_{\sigma_i}/Z'_{\sigma_i})$.

Applying (ii), we complete the proof of (iii). $\square$

Corollary 14.4.21. If $\sigma \in C$ is a permutation of type $1^{\lambda_1}2^{\lambda_2}\dots n^{\lambda_n}$, then $|\widehat{Z}_\sigma| = \left(\prod_{\lambda_i=1} i \right) \left(\prod_{\lambda_i \geq 2} 2i \right)$, written as γ_C .

Proof. It follows from Corollary 14.4.6 and Proposition 14.4.20(iii). $\square$

Remark. Obviously, γ_C only depends on the conjugate class C

14.4.5 Fixed Point Set $F_{(\phi_{g_C}, \phi_C)_C}$

Let $(\phi_{g_C}, \phi_C)_C \in M$. If $\left(\left\{\chi_C^{(i)}\right\}_i, u(C)\right)_C \in F_{(\phi_{g_C}, \phi_C)_C}$, then, according to (14.33), we have

$$\begin{aligned} (\phi_{g_C}, \phi_C)_C \circ \left(\left\{\chi_C^{(i)}\right\}_i, u(C)\right)_C &= \left(\left\{\chi_C^{(\phi_C^{-1}(i))} \phi_{g_C}^{-1}\right\}_i, \phi_{g_C} u(C)\right)_C \\ &= \left(\left\{\chi_C^{(i)}\right\}_i, u(C)\right)_C. \end{aligned}$$

Consequently,

$$\phi_{g_C}(u(C)) = u(C) \quad (14.34)$$

and

$$\chi_C^{(\phi_C^{-1}(i))} \phi_{g_C}^{-1} = \chi_C^{(i)} \quad (14.35)$$

for any $C \in \mathcal{K}_r(G)$, $i \in I_C(r)$.

(14.34) implies $u(C) \in Z_{g_C}$ and $g_C \in Z_{u(C)}$. Considering $u(C) \in C$, we have $u(C) \in Z_{g_C} \cap C$. Conversely, if $\sigma \in Z_{g_C} \cap C$, then $u(C) = \sigma$ satisfies (14.34). Consequently, the number of $u(C)$'s which satisfy (14.34) is $|Z_{g_C} \cap C|$, written

$$\beta_{g_C} := |Z_{g_C} \cap C|.$$

(14.35) is equivalent to

$$\chi_C^{(i)} \phi_{g_C}^{-1} = \chi_C^{(\phi_C(i))}, \quad (14.36)$$

where $\phi_C \in S_{I_C(r)} = S_{r_C}$. It is clear that (14.36) holds if and only if the following (14.36') holds for every cycle, such as $\tau = (i_1, \dots, i_r)$, of σ :

$$\chi_C^{(i)} \phi_{g_C}^{-1} = \chi_C^{(\tau(i))}, \quad i = i_1, \dots, i_r. \quad (14.36')$$

By (14.36'), we have

$$\begin{aligned} \chi_C^{(i_2)} &= \chi_C^{(i_1)} \phi_{g_C}^{-1}, \\ \chi_C^{(i_3)} &= \chi_C^{(i_2)} \phi_{g_C}^{-1} = \chi_C^{(i_1)} (\phi_{g_C}^{-1})^2, \\ &\dots \\ \chi_C^{(i_r)} &= \chi_C^{(i_{r-1})} \phi_{g_C}^{-1} = \chi_C^{(i_1)} (\phi_{g_C}^{-1})^{r-1}, \\ \chi_C^{(i_1)} &= \chi_C^{(i_r)} \phi_{g_C}^{-1} = \chi_C^{(i_1)} (\phi_{g_C}^{-1})^r. \end{aligned} \quad (14.37)$$

We can obtain $\chi_C^{(i_2)}, \dots, \chi_C^{(i_r)}$ when we obtain $\chi_C^{(i_1)}$. Notice $(\phi_{g_C}^{-1})^r = (\phi_{g_C}^r)^{-1} = (\phi_{g_C^r})^{-1}$. Therefore (14.37) is equivalent to

$$\chi_C^{(i_1)} \phi_{g_C^r} = \chi_C^{(i_1)},$$

i.e.

$$\chi_C^{(i_1)} (g_C^r h(g_C^r)^{-1}) = \chi_C^{(i_1)}(h) \quad (14.38)$$

for any $h \in Z_{u(C)}$. Since $g_C \in Z_{u(C)}$ implies $g_C^r \in Z_{u(C)}$, we have

$$\chi_C^{(i_1)} (g_C^r h (g_C^r)^{-1}) = \chi_C^{(i_1)} (g_C^r) \chi_C^{(i_1)} (h) \chi_C^{(i_1)} ((g_C^r)^{-1}) = \chi_C^{(i_1)} (h).$$

This shows (14.38) holds for any characters in $Z_{u(C)}$. By Corollary 14.4.21, there exist γ_C characters $\chi_C^{(i_1)}$'s such that (14.37) holds. Assume that ϕ_C is written as multiplication of k_{ϕ_C} independent cycles. Thus given a $u(C)$, there exist $\gamma_C^{k_{\phi_C}}$ elements $\{\chi_C^{(i)}\}_i$'s such that (14.36) holds. Consequently, for any $C \in \mathcal{K}_r(G)$, there exist $\gamma_C^{k_{\phi_C}} \beta_{g_C}$ distinct elements $\left(\left\{\chi_C^{(i)}\right\}_i, u(C)\right)$'s such that both (14.34) and (14.35) hold.

Consequently, we have

$$|F_{(\phi_{g_C}, \phi_C)_C}| = \prod_{C \in \mathcal{K}_r(G)} \left(\gamma_C^{k_{\phi_C}} \beta_{g_C} \right) \quad (14.39)$$

for any $\phi_{g_C}, \phi_C)_C \in M$.

14.4.6 The Formula Computing the Number of Isomorphic Classes

Applying Burnside's Lemma and (14.39), we have

$$\begin{aligned} \mathcal{N}(G, r) &= \frac{1}{|M|} \sum_{(\phi_{g_C}, \phi_C)_C \in M} |F_{(\phi_{g_C}, \phi_C)_C}| \\ &= \frac{1}{|M|} \sum_{(\phi_{g_C}, \phi_C)_C \in M} \prod_{C \in \mathcal{K}_r(G)} \left(\gamma_C^{k_{\phi_C}} \beta_{g_C} \right) \\ &= \frac{1}{|M|} \prod_{C \in \mathcal{K}_r(G)} \sum_{\substack{\phi_{g_C} \in \text{Inn} G \\ \phi_C \in S_{r_C}}} \left(\gamma_C^{k_{\phi_C}} \beta_{g_C} \right) \\ &= \frac{1}{|M|} \prod_{C \in \mathcal{K}_r(G)} \sum_{\phi_{g_C} \in \text{Inn} G} \sum_{\phi_C \in S_{r_C}} \left(\gamma_C^{k_{\phi_C}} \beta_{g_C} \right) \\ &= \frac{1}{|M|} \prod_{C \in \mathcal{K}_r(G)} \left(\sum_{\phi_{g_C} \in \text{Inn} G} \beta_{g_C} \sum_{\phi_C \in S_{r_C}} \gamma_C^{k_{\phi_C}} \right) \end{aligned} \quad (14.40)$$

where β_{g_C} and k_{ϕ_C} are the same as above. $|M| = \prod_{C \in \mathcal{K}_r(G)} (n! r_C!)$ when $n \neq 2, 6$; $|M| =$

$\prod_{C \in \mathcal{K}_r(G)} (r_C!)$ when $n = 2$.

Next we simplify formula (14.40). In (14.40), since γ_C depends only on conjugate class, it can be computed by means of Corollary 14.4.21. Notice that k_{ϕ_C} denotes the number of independent cycles in independent cycle decomposition of ϕ_C in S_{r_C} .

Definition 14.4.22. (See [123, pages 292-295]) Let

$$[x]_n = x(x-1)(x-2) \cdots (x-n+1) \quad n = 1, 2, \dots$$

Obviously, $[x]_n$ is a polynomial with degree n . Let $(-1)^{n-k} s(n, k)$ denote the coefficient of x^k in $[x]_n$ and $s(n, k)$ is called the 1st Stirling number. That is,

$$[x]_n = \sum_{k=1}^n (-1)^{n-k} s(n, k) x^k.$$

[123, Theorem 8.2.9] obtained the meaning about the 1st Stirling number:

Lemma 14.4.23. (*[123, Theorem 8.2.9]*) *There exactly exist $s(n, k)$ permutations in S_n , of which the numbers of independent cycles in independent cycle decomposition are k . That is, $s(n, k) = |\{\tau \in S_n \mid \text{the number of independent cycles in independent cycle decomposition of } \tau \text{ is } k\}|$.*

About β_g , we have

Lemma 14.4.24. *Let $G = S_n$. Then $\sum_{g \in G} \beta_g = \sum_{g \in G} |Z_g \cap C| = |G| = n!$ for any $C \in \mathcal{K}(G)$.*

Proof. For any $C \in \mathcal{K}(G)$, we have

$$\begin{aligned}
 \sum_{g \in G} |Z_g \cap C| &= \sum_{g \in G} \left| \bigcup_{h \in C} (Z_g \cap \{h\}) \right| \\
 &= \sum_{g \in G} \sum_{h \in C} |Z_g \cap \{h\}| \\
 &= \sum_{h \in C} \sum_{g \in G} |Z_g \cap \{h\}| \\
 &= \sum_{h \in C} |Z_h| \\
 &= n! \quad (\text{by Corollary 14.4.17(ii)}). \quad \square
 \end{aligned}$$

Theorem 14.4.25. *Let $G = S_n$ with $n \neq 6$. Then*

$$\mathcal{N}(G, r) = \prod_{C \in \mathcal{K}_r(G)} \binom{\gamma_C + r_C - 1}{r_C}. \quad (14.41)$$

Proof. We shall show in following two cases.

(i) $n \neq 2, 6$. By Proposition 14.4.1(ii), $\text{Inn}G = G$. Applying (14.40), we have

$$\begin{aligned}
\mathcal{N}(S_n, r) &= \frac{1}{\prod_{C \in \mathcal{K}_r(G)} (n! r_C!)} \prod_{C \in \mathcal{K}_r(G)} \left(\sum_{g_C \in G} \beta_{g_C} \sum_{\phi_C \in S_{r_C}} \gamma_C^{k_{\phi_C}} \right) \\
&= \prod_{C \in \mathcal{K}_r(G)} \left(\frac{1}{n!} \sum_{g_C \in G} \beta_{g_C} \right) \prod_{C \in \mathcal{K}_r(G)} \left(\frac{1}{r_C!} \sum_{\phi_C \in S_{r_C}} \gamma_C^{k_{\phi_C}} \right) \\
&= \prod_{C \in \mathcal{K}_r(G)} \left(\frac{1}{n!} n! \right) \prod_{C \in \mathcal{K}_r(G)} \left(\frac{1}{r_C!} \sum_{k=1}^{r_C} s(r_C, k) \gamma_C^k \right) \quad (\text{by Lemma 14.4.24, 14.4.23}) \\
&= \prod_{C \in \mathcal{K}_r(G)} \left(\frac{1}{r_C!} (-1)^{r_C} \sum_{k=1}^{r_C} (-1)^{r_C-k} s(r_C, k) (-\gamma_C)^k \right) \\
&= \prod_{C \in \mathcal{K}_r(G)} \left(\frac{1}{r_C!} (-1)^{r_C} (-\gamma_C) (-\gamma_C - 1) \cdots (-\gamma_C - r_C + 1) \right) \quad (\text{by Definition 14.4.22}) \\
&= \prod_{C \in \mathcal{K}_r(G)} \frac{(\gamma_C + r_C - 1) \cdots (\gamma_C + 1) \gamma_C}{r_C!} \\
&= \prod_{C \in \mathcal{K}_r(G)} \binom{\gamma_C + r_C - 1}{r_C}. \quad \square
\end{aligned}$$

(ii) $n = 2$. In this case we have $\text{Inn}G = \{1\}$, $\phi_{g_C} = 1$ and $\beta_{g_C} = 1$ in (14.40). Thus

$$\begin{aligned}
\mathcal{N}(S_n, r) &= \frac{1}{\prod_{C \in \mathcal{K}_r(G)} (r_C!)} \prod_{C \in \mathcal{K}_r(G)} \sum_{\phi_C \in S_{r_C}} \left(\gamma_C^{k_{\phi_C}} \right) \\
&= \prod_{C \in \mathcal{K}_r(G)} \left(\frac{1}{r_C!} \sum_{\phi_C \in S_{r_C}} \gamma_C^{k_{\phi_C}} \right) \quad (\text{similar to the computation of (i)}) \\
&= \prod_{C \in \mathcal{K}_r(G)} \binom{\gamma_C + r_C - 1}{r_C}. \quad \square
\end{aligned}$$

Corollary 14.4.26. *If $r = \sum_{C \in \mathcal{K}(G)} C$, i.e. $r_C = 1$ for any $C \in \mathcal{K}(G)$, then*

$$\mathcal{N}(G, r) = \prod_{C \in \mathcal{K}(G)} \gamma_C. \quad (14.42)$$

Proof. Considering $\mathcal{K}_r(G) = \mathcal{K}(G)$ and applying Theorem 14.4.25, we can complete the proof. $\square$

Corollary 14.4.27. *If $C_0 = \{(1)\}$ and $\mathcal{K}_r(G) = \{C_0\}$, then*

$$\mathcal{N}(G, r) = \begin{cases} 1 & n = 1 \\ r_{C_0} + 1 & n \geq 2 \end{cases}.$$

Proof. By Corollary 14.4.21, we have $\gamma_{C_0} = 1$ when $n = 1$; $\gamma_{C_0} = 2$ when $n \geq 2$. Applying Theorem 14.4.25, we can complete the proof. $\square$

14.4.7 Represetative

We have obtained the number of isomorphic classes, now we give the representatives

Definition 14.4.28. Given an $RSC(G, r, \vec{\chi}, u)$, written $\widehat{Z_{u(C)}} = \{\xi_{u(C)}^{(i)} \mid i = 1, 2, \dots, \gamma_C\}$ and $n_i := |\{j \mid \chi_C^{(j)} = \xi_{u(C)}^{(i)}\}|$, for any $C \in \mathcal{K}_r(G)$ and $1 \leq i \leq \gamma_{u(C)}$, then $\{(n_1, n_2, \dots, n_{\gamma_C})\}_{C \in \mathcal{K}_r(G)}$ is called the type of $RSC(G, r, \vec{\chi}, u)$.

Lemma 14.4.29. $RSC(G, r, \vec{\chi}, u)$ and $RSC(G, r, \vec{\chi}', u)$ are isomorphic if and only if they have the same type.

Proof. If $RSC(G, r, \vec{\chi}, u)$ and $RSC(G, r, \vec{\chi}', u)$ have the same type, then there exists a bijective map $\phi_C \in S_{r_C}$ such that $\chi_C'^{(\phi_C(i))} = \chi_C^{(i)}$ for any $i \in I_C(r)$, $C \in \mathcal{K}_r(G)$. Let $\phi = id_G$ and $g_C = 1$. It follows from Corollary 14.4.12 that they are isomorphic.

Conversely, if $RSC(G, r, \vec{\chi}, u)$ and $RSC(G, r, \vec{\chi}', u)$ are isomorphic, then, by Corollary 14.4.12, there exist $\phi \in \text{Aut}G$ and $\phi_C \in S_{r_C}$, as well as $g_C \in G$ such that $u(C) = \phi_{g_C}(u(C))$ and $\chi_C'^{(\phi_C(i))}\phi_{g_C}(h) = \chi_C^{(i)}(h)$ for any $C \in \mathcal{K}_r(G)$, $h \in Z_{u(C)}$, $i \in I_C(r)$. Thus $g_C \in Z_{u(C)}$ and $\chi_C'^{(\phi_C(i))}\phi_{g_C}(h) = \chi_C'^{(\phi_C(i))}(g_C h g_C^{-1}) = \chi_C'^{(\phi_C(i))}(h) = \chi_C^{(i)}(h)$ for any $h \in Z_{u(C)}$, i.e. $\chi_C'^{(\phi_C(i))} = \chi_C^{(i)}$. This implies that $RSC(G, r, \vec{\chi}, u)$ and $RSC(G, r, \vec{\chi}', u)$ have the same type. $\square$

Theorem 14.4.30. Let $G = S_n$ ($n \neq 6$) and u_0 be a map from $\mathcal{K}(G) \rightarrow G$ with $u_0(C) \in C$ for any $C \in \mathcal{K}(G)$. Let $\bar{\Omega}(G, r, u_0)$ denote the set consisting of all elements with distinct type in $\{(G, r, \vec{\chi}, u_0) \mid (G, r, \vec{\chi}, u_0) \text{ is an } RSC\}$. Then $\bar{\Omega}(G, r, u_0)$ becomes the representative set of $\Omega(G, r)$.

Proof. According to Theorem 14.4.25 and [35], the number of elements in $\bar{\Omega}(G, r, u_0)$ is the same as the number of isomorphic classes in $\Omega(G, r)$. Applying Lemma 14.4.29 we complete the proof. $\square$

For example, applying Corollary 14.4.26 we compute $\mathcal{N}(G, r)$ for $G = S_3, S_4, S_5$ and $r = \sum_{C \in \mathcal{K}(G)} C$.

Example 14.4.31. (i) Let $G = S_3$ and $r = \sum_{C \in \mathcal{K}(G)} C$. Then

type	1^3	$1^1 2^1$	3^1
γ_C	2	2	3

By (14.42), $\mathcal{N}(S_3, r) = 12$. We explicitly write the type of representative elements as follows:
 $\{(1, 0)_{1^3}, (1, 0)_{1^1 2^1}, (1, 0, 0)_{3^1}\}, \{(0, 1)_{1^3}, (1, 0)_{1^1 2^1}, (1, 0, 0)_{3^1}\}, \{(1, 0)_{1^3}, (0, 1)_{1^1 2^1}, (1, 0, 0)_{3^1}\},$
 $\{(0, 1)_{1^3}, (0, 1)_{1^1 2^1}, (1, 0, 0)_{3^1}\}, \{(1, 0)_{1^3}, (1, 0)_{1^1 2^1}, (0, 1, 0)_{3^1}\}, \{(1, 0)_{1^3}, (1, 0)_{1^1 2^1}, (0, 0, 1)_{3^1}\},$
 $\{(0, 1)_{1^3}, (1, 0)_{1^1 2^1}, (0, 1, 0)_{3^1}\}, \{(0, 1)_{1^3}, (1, 0)_{1^1 2^1}, (0, 0, 1)_{3^1}\}, \{(1, 0)_{1^3}, (0, 1)_{1^1 2^1}, (0, 1, 0)_{3^1}\},$
 $\{(1, 0)_{1^3}, (0, 1)_{1^1 2^1}, (0, 0, 1)_{3^1}\}, \{(0, 1)_{1^3}, (0, 1)_{1^1 2^1}, (0, 1, 0)_{3^1}\}, \{(0, 1)_{1^3}, (0, 1)_{1^1 2^1}, (0, 0, 1)_{3^1}\}.$

(ii) Let $G = S_4$ and $r = \sum_{C \in \mathcal{K}(G)} C$. Then

type	1^4	$1^1 3^1$	$1^2 2^1$	2^2	4^1
γ_C	2	3	4	4	4

By (14.42), $\mathcal{N}(S_4, r) = 384$.

(iii) Let $G = S_4$ and $r = \sum_{C \in \mathcal{K}(G)} C$. Then

type	1^5	$1^1 4^1$	$1^2 3^1$	$1^3 2^1$	$1^1 2^2$	$2^1 3^1$	5^1
γ_C	2	4	6	4	4	6	5

By (14.42), $\mathcal{N}(S_5, r) = 23040$.

Chapter 15

Appendix

The proof of Theorem 14.1.1: We define two functors W and V as follows.

$$\begin{aligned} W : {}^{kG}_{kG}\mathcal{M}^{kG}_{kG} &\rightarrow \prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}} \\ B &\mapsto \{u(C)B^1\}_{C \in \mathcal{K}(G)} \\ f &\mapsto W(f), \end{aligned}$$

where for any morphism $f : B \rightarrow B'$ in ${}^{kG}_{kG}\mathcal{M}^{kG}_{kG}$, $W(f) = \{u(C)f^1\}_{C \in \mathcal{K}(G)}$.

$$\begin{aligned} V : \prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}} &\rightarrow {}^{kG}_{kG}\mathcal{M}^{kG}_{kG} \\ \{M(C)\}_{C \in \mathcal{K}(G)} &\mapsto \bigoplus_{y=xg_\theta^{-1}u(C)g_\theta, x,y \in G} x \otimes M(C) \otimes_{kZ_{u(C)}} g_\theta, \\ f &\mapsto V(f), \end{aligned}$$

where for any morphism $f = \{f_C\}_{C \in \mathcal{K}(G)} : \{M(C)\}_{C \in \mathcal{K}(G)} \rightarrow \{N(C)\}_{C \in \mathcal{K}(G)}$, $V(f)(x \otimes m \otimes_{kZ_{u(C)}} g_\theta) = x \otimes f_C(m) \otimes_{kZ_{u(C)}} g_\theta$ for any $m \in M_C$, $x, y \in G$ with $x^{-1}y = g_\theta^{-1}u(C)g_\theta$. That is, ${}^yV(f)^x = \text{id} \otimes f_C \otimes \text{id}$.

We will show that W and V are mutually inverse functors by three steps.

(i) Let B be a kG -Hopf bimodule. Then it is easy to see that $u(C)B^1$ is a right $kZ_{u(C)}$ -module with the module action given by

$$b \triangleleft h = h^{-1} \cdot b \cdot h, \quad b \in {}^{u(C)}B^1, \quad h \in Z_{u(C)}. \quad (15.1)$$

Hence $W(B)$ is an object in $\prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$. Moreover, one can see that $W(f)$ is a morphism from $W(B)$ to $W(B')$ in $\prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$ if $f : B \rightarrow B'$ is a morphism in ${}^{kG}_{kG}\mathcal{M}^{kG}_{kG}$. Now it is straightforward to check that W is a functor from ${}^{kG}_{kG}\mathcal{M}^{kG}_{kG}$ to $\prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$.

(ii) Let $M = \{M(C)\}_{C \in \mathcal{K}(G)} \in \prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$, we define

$$\begin{aligned} {}^yV(M)^x &= x \otimes M(C) \otimes_{kZ_{u(C)}} g_\theta, \\ h \cdot (x \otimes m \otimes_{kZ_{u(C)}} g_\theta) &= hx \otimes m \otimes_{kZ_{u(C)}} g_\theta, \\ (x \otimes m \otimes_{kZ_{u(C)}} g_\theta) \cdot h &= xh \otimes m \otimes_{kZ_{u(C)}} g_\theta h = xh \otimes (m \triangleleft \zeta_\theta(h)) \otimes_{kZ_{u(C)}} g_{\theta'}, \end{aligned} \quad (15.2)$$

where $h, x, y \in G$ with $x^{-1}y \in C$ and the relation (14.2) and (14.3), $m \in M(C)$. It is easy to check that $V(M)$ is a kG -Hopf bimodule. If $f : \{M(C)\}_{C \in \mathcal{K}(G)} \rightarrow \{N(C)\}_{C \in \mathcal{K}(G)}$ is a

morphism in $\prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$, then one can see that $V(f) : V(M) \rightarrow V(N)$ is a morphism in ${}^{kG}_{kG}\mathcal{M}_{kG}^{kG}$. Now let $f : \{M(C)\}_{C \in \mathcal{K}(G)} \rightarrow \{N(C)\}_{C \in \mathcal{K}(G)}$ and $f' : \{N(C)\} \rightarrow \{L(C)\}_{C \in \mathcal{K}(G)}$ be two morphisms in $\prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$. Then for any $x, y \in G$ with $x^{-1}y = g_\theta^{-1}u(C)g_\theta$ and $m \in M(C)$, we have

$$\begin{aligned} V(f'f)(x \otimes m \otimes_{kZ_{u(C)}} g_\theta) &= x \otimes (f'f)_C(m) \otimes_{kZ_{u(C)}} g_\theta \\ &= x \otimes f'_C(f_C(m)) \otimes_{kZ_{u(C)}} g_\theta \\ &= V(f')(V(f)(x \otimes m \otimes_{kZ_{u(C)}} g_\theta)). \end{aligned}$$

Therefore, $V(f'f) = V(f')V(f)$. It follows that V is a functor from $\prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$ to ${}^{kG}_{kG}\mathcal{M}_{kG}^{kG}$.

(iii) We now construct two natural isomorphisms φ and ψ as follows. Let $B \in {}^{kG}_{kG}\mathcal{M}_{kG}^{kG}$. Define a k -linear map φ_B from B to $VW(B) = \bigoplus_{y=xg_\theta^{-1}u(C)g_\theta, x, y \in G} x \otimes {}^{u(C)}B^1 \otimes_{kZ_{u(C)}} g_\theta$ by

$$\varphi_B(b) = x \otimes (g_\theta x^{-1}) \cdot b \cdot g_\theta^{-1} \otimes_{kZ_{u(C)}} g_\theta, \quad (15.3)$$

for any $x, y \in G$ satisfying Eq.(14.2), $b \in {}^yB^x$. Now we have

$$\begin{aligned} \delta^-(g_\theta x^{-1} \cdot b \cdot g_\theta^{-1}) &= (g_\theta x^{-1}) \cdot (\delta^-(b)) \cdot g_\theta^{-1} \quad (B \text{ is a } kG\text{-Hopf bimodule.}) \\ &= (g_\theta x^{-1}) \cdot (y \otimes b) \cdot g_\theta^{-1} \\ &= g_\theta x^{-1} y g_\theta^{-1} \otimes g_\theta x^{-1} \cdot b \cdot g_\theta^{-1} \\ &= u(C) \otimes g_\theta x^{-1} \cdot b \cdot g_\theta^{-1} \quad (\text{by Eq.(14.2)}) \end{aligned}$$

and

$$\delta^+(g_\theta x^{-1} \cdot b \cdot g_\theta^{-1}) = g_\theta x^{-1} \cdot b \cdot g_\theta^{-1} \otimes 1.$$

It follows that $g_\theta x^{-1} \cdot b \cdot g_\theta^{-1} \in {}^{u(C)}B^1$, and hence $x \otimes g_\theta x^{-1} \cdot b \cdot g_\theta^{-1} \otimes_{kZ_{u(C)}} g_\theta \in {}^yVW(B)^x$. This shows that φ_B is a kG -bicomodule map.

Next, let $x, y \in G$ with $x^{-1}y = g_\theta^{-1}u(C)g_\theta$, $C \in \mathcal{K}(G)$, and let $b \in {}^yB^x$ and $h \in G$. Then $h \cdot b \in {}^{hy}B^{hx}$ and $(hx)^{-1}(hy) = x^{-1}y = g_\theta^{-1}u(C)g_\theta$. Hence

$$\begin{aligned} \varphi_B(h \cdot b) &= hx \otimes g_\theta (hx)^{-1} \cdot (h \cdot b) \cdot g_\theta^{-1} \otimes_{kZ_{u(C)}} g_\theta \\ &= hx \otimes g_\theta x^{-1} \cdot b \cdot g_\theta^{-1} \otimes_{kZ_{u(C)}} g_\theta \\ &= h \cdot (x \otimes g_\theta x^{-1} \cdot b \cdot g_\theta^{-1} \otimes_{kZ_{u(C)}} g_\theta) \\ &= h \cdot \varphi_B(b). \end{aligned}$$

This shows that φ_B is a left kG -module map. On the other hand, $b \cdot h \in {}^{yh}B^{xh}$ and $(xh)^{-1}(yh) = h^{-1}x^{-1}yh = h^{-1}g_\theta^{-1}u(C)g_\theta h$. By Eq.(14.3), one gets $g_\theta h = \zeta_\theta(h)g_{\theta'}$ with $\zeta_\theta(h) \in Z_{u(C)}$ and $\theta' \in \Theta_C$. Hence $(xh)^{-1}(yh) = g_{\theta'}^{-1}\zeta_\theta(h)^{-1}u(C)\zeta_\theta(h)g_{\theta'} = g_{\theta'}^{-1}u(C)g_{\theta'}$. Thus we have

$$\begin{aligned} \varphi_B(b \cdot h) &= xh \otimes g_{\theta'}(xh)^{-1} \cdot (b \cdot h) \cdot g_{\theta'}^{-1} \otimes_{kZ_{u(C)}} g_{\theta'} \\ &= xh \otimes g_{\theta'} h^{-1} x^{-1} \cdot b \cdot h g_{\theta'}^{-1} \otimes_{kZ_{u(C)}} g_{\theta'} \\ &= xh \otimes \zeta_\theta(h)^{-1} g_\theta x^{-1} \cdot b \cdot g_\theta^{-1} \zeta_\theta(h) \otimes_{kZ_{u(C)}} g_{\theta'} \\ &= xh \otimes ((g_\theta x^{-1} \cdot b \cdot g_\theta^{-1}) \triangleleft \zeta_\theta(h)) \otimes_{kZ_{u(C)}} g_{\theta'} \quad (\text{by (14.5)}) \\ &= (x \otimes g_\theta x^{-1} \cdot b \cdot g_\theta^{-1} \otimes_{kZ_{u(C)}} g_\theta) \cdot h \\ &= \varphi_B(b) \cdot h. \end{aligned}$$

It follows that φ_B is a right kG -module map, and hence φ_B is a kG -Hopf bimodule map.

Suppose that $\varphi_B(b) = 0$ with $b \in {}^y B^x$. Then $x \otimes g_\theta x^{-1} \cdot b \cdot g_\theta^{-1} \otimes_{kZ_{u(C)}} g_\theta = 0$, and hence $g_\theta x^{-1} \cdot b \cdot g_\theta^{-1} \otimes_{kZ_{u(C)}} g_\theta = 0$. Since $M(C) \otimes_{kZ_{u(C)}} kZ_{u(C)} g_\theta \cong M(C) \otimes_{kZ_{u(C)}} kZ_{u(C)} \cong M(C)$ as k -vector spaces for any right $kZ_{u(C)}$ -module $M(C)$, we have $g_\theta x^{-1} \cdot b \cdot g_\theta^{-1} = 0$. This implies $b = 0$. It follows that φ_B is injective.

Let $x \otimes b \otimes_{kZ_{u(C)}} g_\theta \in {}^y VW(B)^x$ with $b \in {}^{u(C)} B^1$ and $x^{-1}y = g_\theta^{-1}u(C)g_\theta$. Put $b' := xg_\theta^{-1} \cdot b \cdot g_\theta$. Then

$$\begin{aligned} \delta^-(b') &= xg_\theta^{-1} \cdot \delta^-(b) \cdot g_\theta \\ &= xg_\theta^{-1} \cdot (u(C) \otimes b) \cdot g_\theta \\ &= y \otimes b', \\ \delta^+(b') &= b' \otimes x. \end{aligned}$$

Hence $b' \in {}^y B^x$ and $\varphi_B(b') = x \otimes b \otimes_{kZ_{u(C)}} g_\theta$. This shows that φ_B is surjective.

If $f : B \rightarrow B'$ is a kG -Hopf bimodule map, then it is easy to see that $\varphi_{B'} \circ f = (VW)(f) \circ \varphi_B$. Thus we have proven that φ is a natural equivalence from the identity functor $id_{\mathcal{B}(kG)}$ to the composition functor VW .

Now for any $M = \{M(C)\}_{C \in \mathcal{K}(G)} \in \prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$, it follows from Eq.(14.2) that ${}^{u(C)}V(M)^1 = 1 \otimes M(C) \otimes_{kZ_{u(C)}} 1$. Hence $WV(M) = \{1 \otimes M(C) \otimes_{kZ_{u(C)}} 1\}_{C \in \mathcal{K}(G)}$. Define a map ψ_M from M to $WV(M)$ by

$$(\psi_M)_C(m) = 1 \otimes m \otimes_{kZ_{u(C)}} 1,$$

where $C \in \mathcal{K}(G), m \in M(C)$. Clearly, $(\psi_M)_C$ is k -linear. Let $m \in M(C), h \in Z_{u(C)}$. Then

$$\begin{aligned} ((\psi_M)_C(m)) \triangleleft h &= (1 \otimes m \otimes_{kZ_{u(C)}} 1) \triangleleft h \\ &= h^{-1} \cdot (1 \otimes m \otimes_{kZ_{u(C)}} 1) \cdot h \\ &= 1 \otimes m \triangleleft h \otimes_{kZ_{u(C)}} 1 \\ &= (\psi_M)_C(m \triangleleft h). \end{aligned}$$

That is, $(\psi_M)_C$ is a $kZ_{u(C)}$ -module map. Obviously, $(\psi_M)_C$ is bijective and ψ is a natural transformation. Thus ψ is a natural equivalence from the identity functor $id_{\prod_{C \in \mathcal{K}(G)} \mathcal{M}_{kZ_{u(C)}}$ to the composition functor WV . $\square$

The proof of Lemma 14.1.3: (i) We first show that every irreducible representation of G is one dimensional. In fact, this is a well-known conclusion. For the completeness, we give a proof. Assume that ρ is an irreducible representation of G corresponding to V . For any $g \in G$, since $\rho(g^m) = id_V$, all eigenvalues of $\rho(g)$ is m -th roots of 1, which belong to k . Say that λ_g is an eigenvalue of $\rho(g)$. It is clear that $Ker(\rho(g) - \lambda_g id_V) := W$ is a kG -submodule of V and $W \neq 0$ since λ_g is an eigenvalue of $\rho(g)$. Therefore, $W = V$, which implies $\rho(g) = \lambda_g id_V$. Considering V is irreducible, we have that $\dim V = 1$.

Next we show that the order n of G is not divisible by $\text{char}(k) = p$. Assume that $\beta \in k$ is a primitive m -root of 1. If $p \mid n$, then $p \mid m$ since there exists an element with order p . Say $m = pt$. See $(\beta^m - 1) = (\beta^t - 1)^p = 0$, which implies $\beta^t = 1$. We get a contradiction.

Using Maschke's Theorem, we have that every kG -module is a pointed module.

(ii) Assume M is a kG -Hopf bimodule. By Theorem 14.1.1, $M \cong \bigoplus_{y=g\theta^{-1}u(C)g\theta, x,y \in G} x \otimes M(C) \otimes_{kZ_{u(C)}} g\theta$ as kG -Hopf bimodules with $M(C) = {}^{u(C)}M^1$ for any $C \in \mathcal{K}(G)$. It follows from (i) that $M(C)$ is a right pointed kG -module. Therefore, M is a PM kG -Hopf bimodule. $\square$

The proof of Lemma 14.1.4: It is clear that If ξ is a graded linear map from graded vector space $V = \bigoplus_{i \geq 0} V_{(i)}$ to graded vector space $V' = \bigoplus_{i \geq 0} V'_{(i)}$, then

$$\xi \iota_i \pi_i = \iota_i \pi_i \xi, \quad \xi \iota_i = \iota_i \pi_i \xi \iota_i, \quad \pi_i \xi = \pi_i \xi \iota_i \pi_i. \quad (15.4)$$

for $i = 1, 2, \dots$.

Observe that (M, α^-, α^+) is a B -bimodule. Since Δ_H and μ_H are graded, we have the following equation by (15.4):

$$\Delta \iota_1 \pi_1 = (\iota_1 \pi_1 \otimes \iota_0 \pi_0) \Delta + (\iota_0 \pi_0 \otimes \iota_1 \pi_1) \Delta. \quad (15.5)$$

Since $\pi_1 \iota_1 = \text{id}$, it follows from Eq.(15.5) that

$$\Delta \iota_1 = (\iota_1 \pi_1 \otimes \iota_0 \pi_0) \Delta \iota_1 + (\iota_0 \pi_0 \otimes \iota_1 \pi_1) \Delta \iota_1. \quad (15.6)$$

By a straightforward computation, one can show the following two equations:

$$\delta^- \pi_1 = (\pi_0 \otimes \pi_1) \Delta, \quad \delta^+ \pi_1 = (\pi_1 \otimes \pi_0) \Delta. \quad (15.7)$$

Now we have

$$\begin{aligned} (\text{id} \otimes \delta^-) \delta^- &= (\text{id} \otimes \delta^-) (\pi_0 \otimes \pi_1) \Delta \iota_1 \\ &= (\pi_0 \otimes \pi_0 \otimes \pi_1) (\text{id} \otimes \Delta) \Delta \iota_1 \quad (\text{by Eq.(15.7)}), \end{aligned}$$

and

$$\begin{aligned} (\Delta_B \otimes \text{id}) \delta^- &= (\Delta_B \otimes \text{id}) (\pi_0 \otimes \pi_1) \Delta \iota_1 \\ &= (\pi_0 \otimes \pi_0 \otimes \pi_1) (\Delta \otimes \text{id}) \Delta \iota_1. \end{aligned}$$

This shows that $(\text{id} \otimes \delta^-) \delta^- = (\Delta_B \otimes \text{id}) \delta^-$. Clearly, $(\varepsilon \otimes \text{id}) \delta^- = \text{id}$. Hence (M, δ^-) is a left B -comodule. Similarly, one can show that (M, δ^+) is a right B -comodule and (M, δ^-, δ^+) is a B -bicomodule. Next, we have

$$\begin{aligned} \delta^- \alpha^+ &= (\pi_0 \otimes \pi_1) \Delta \iota_1 \pi_1 \mu (\iota_1 \otimes \iota_0) \\ &= (\pi_0 \otimes \pi_1) \Delta \mu (\iota_1 \otimes \iota_0) \quad (\text{by Eq.(15.5)}) \\ &= (\pi_0 \mu \otimes \pi_1 \mu) (\text{id} \otimes \tau \otimes \text{id}) (\Delta \iota_1 \otimes \Delta \iota_0) \\ &= (\mu_B \otimes \pi_1) (\pi_0 \otimes \pi_0 \otimes \mu) (\text{id} \otimes \tau \otimes \text{id}) (\Delta \otimes \iota_0 \otimes \iota_0) (\iota_1 \otimes \Delta_B) \\ &= (\mu_B \otimes \pi_1) (\text{id} \otimes \text{id} \otimes \mu) (\text{id} \otimes \text{id} \otimes \iota_1 \otimes \iota_0) (\text{id} \otimes \tau \otimes \text{id}) (\pi_0 \otimes \pi_1 \otimes \text{id} \otimes \text{id}) \\ &\quad (\Delta \iota_1 \otimes \Delta_B) \quad (\text{by Eq.(15.6)}) \\ &= (\mu \otimes \alpha^+) (\text{id} \otimes \tau \otimes \text{id}) (\delta^- \otimes \Delta_B), \end{aligned}$$

where τ is the usual twist map. This shows that δ^- is a right B -module map. Similarly, one can show that δ^- is a left B -module map, and hence a B -bimodule map. A similar argument shows that δ^+ is also a B -bimodule map. Hence $(M, \alpha^-, \alpha^+, \delta^-, \delta^+)$ is a B -Hopf bimodule. $\square$

Proof of Lemma 14.1.5: (i) Suppose that the tensor algebra $T_B(M)$ of M over B admits a graded Hopf algebra structure. Obviously, B is a Hopf algebra. It follows from Lemma 14.1.4 that M is a B -Hopf bimodule.

Conversely, suppose that B admits a Hopf algebra structure and M admits a B -Hopf bimodule structure. By [91, Section 1.4], $T_B(M)$ is a bialgebra with the counit $\varepsilon = \varepsilon_B \pi_0$ and the comultiplication $\Delta = (\iota_0 \otimes \iota_0)\Delta_B + \sum_{n>0} \mu^{n-1} T_n(\Delta_M)$, where ε_B is the counit of B , $\Delta_M = (\iota_0 \otimes \iota_1)\delta_M^- + (\iota_1 \otimes \iota_0)\delta_M^+$. It is easy to see that $T_B(M)$ is a graded bialgebra. Then it follows from [91, Proposition 1.5.1] that $T_B(M)$ is a graded Hopf algebra.

(ii) Assume that the cotensor coalgebra $T_B^c(M)$ of M over B admits a graded Hopf algebra structure. Clearly, B is a Hopf algebra. By Lemma 14.1.4 one gets that M is a B -Hopf bimodule.

Conversely, assume that B admits a Hopf algebra structure and M admits a B -Hopf bimodule structure. By [91, Section 1.4], $T_B^c(M)$ is a bialgebra containing B as a subbialgebra, and the multiplication of $T_B^c(M)$ is $\mu = \mu_B(\pi_0 \otimes \pi_0) + \sum_{n>0} T_n^c(\mu_M)\Delta_{n-1}$, where $\mu_M = \alpha_M^-(\pi_0 \otimes \pi_1) + \alpha_M^+(\pi_1 \otimes \pi_0)$. Since Δ preserves the grading, μ is a graded map. Thus $T_B^c(M)$ is a graded bialgebra. By [91, Proposition 1.5.1], $T_B^c(M)$ is a graded Hopf algebra. $\square$

The proof of Lemma 14.1.6: (i) Since ϕ is a Hopf algebra map, $\iota_0\phi : B \rightarrow T_{B'}(M')$ is an algebra map. Since ψ is a B -bimodule map from M to ${}^\phi M'_\phi$, $\iota_1\psi$ is a B -bimodule map from M to $\iota_0\phi T_{B'}(M')\iota_0\phi$. It follows that $T_B(\iota_0\phi, \iota_1\psi)$ is an algebra homomorphism from $T_B(M)$ to $T_{B'}(M')$. Clearly, $T_B(\iota_0\phi, \iota_1\psi)$ preserves the gradation of $T_B(M)$ and $T_{B'}(M')$.

Let $\Phi := T_B(\iota_0\phi, \iota_1\psi)$. Then both $\Delta_{T_{B'}(M')}\Phi$ and $(\Phi \otimes \Phi)\Delta_{T_B(M)}$ are graded algebra maps from $T_B(M)$ to $T_{B'}(M') \otimes T_{B'}(M')$. Now we show that $\Delta_{T_{B'}(M')}\Phi = (\Phi \otimes \Phi)\Delta_{T_B(M)}$. By the universal property of $T_B(M)$, we only have to prove that $\Delta_{T_{B'}(M')}\Phi\iota_n = (\Phi \otimes \Phi)\Delta_{T_B(M)}\iota_n$ for $n = 1, 2$. Since ϕ is a coalgebra map, one can see that $\Delta_{T_{B'}(M')}\Phi\iota_0 = (\Phi \otimes \Phi)\Delta_{T_B(M)}\iota_0$. Note that ψ is a B' -bicomodule map from ${}^\phi M^\phi$ to M' . Hence by the proof of Lemma 14.1.5 we have

$$\begin{aligned}
\Delta_{T_{B'}(M')}\Phi\iota_1 &= \Delta_{T_{B'}(M')}\iota_1\psi \\
&= (\iota_0 \otimes \iota_1)\delta_{M'}^-\psi + (\iota_1 \otimes \iota_0)\delta_{M'}^+\psi \\
&= (\iota_0 \otimes \iota_1)(\phi \otimes \psi)\delta_M^- + (\iota_1 \otimes \iota_0)(\psi \otimes \phi)\delta_M^+ \\
&= (\iota_0\phi \otimes \iota_1\psi)\delta_M^- + (\iota_1\psi \otimes \iota_0\phi)\delta_M^+ \\
&= (\Phi \otimes \Phi)(\iota_0 \otimes \iota_1)\delta_M^- + (\Phi \otimes \Phi)(\iota_1 \otimes \iota_0)\delta_M^+ \\
&= (\Phi \otimes \Phi)[(\iota_0 \otimes \iota_1)\delta_M^- + (\iota_1 \otimes \iota_0)\delta_M^+] \\
&= (\Phi \otimes \Phi)\Delta_{T_B(M)}\iota_1.
\end{aligned}$$

Furthermore, it is easy to see that $\varepsilon_{T_{B'}(M')}\Phi = \varepsilon_{T_B(M)}$. It follows that Φ is a coalgebra map, and hence a Hopf algebra map since any bialgebra map between two Hopf algebras is a Hopf algebra map.

(ii) Since ϕ is a Hopf algebra map, $\phi\pi_0 : T_B^c(M) \rightarrow B'$ is a coalgebra map. Since ψ is a B' -bicomodule map from ${}^\phi M^\phi$ to M' , $\psi\pi_1$ is a B' -bicomodule map from ${}^{\phi\pi_0}T_B^c(M)^{\phi\pi_0}$ to M' .

We also have $\text{corad}(T_B^c(M)) \subseteq \iota_0(B)$, and hence $(\psi\pi_1)(\text{corad}(T_B^c(M))) = 0$. It follows that $T_{B'}^c(\phi\pi_0, \psi\pi_1)$ is a coalgebra map from $T_B^c(M)$ to $T_{B'}^c(M')$. Clearly, $T_{B'}^c(\phi\pi_0, \psi\pi_1)$ is a graded map.

Put $\Psi := T_{B'}^c(\phi\pi_0, \psi\pi_1)$. Then both $\Psi\mu_{T_B^c(M)}$ and $\mu_{T_{B'}^c(M')}(\Psi \otimes \Psi)$ are graded coalgebra maps from $T_B^c(M) \otimes T_B^c(M)$ to $T_{B'}^c(M')$. Since $T_B^c(M) \otimes T_B^c(M)$ is a graded coalgebra, $\text{corad}(T_B^c(M) \otimes T_B^c(M)) \subseteq (T_B^c(M) \otimes T_B^c(M))_0 = \iota_0(B) \otimes \iota_0(B)$. It follows that $(\pi_1\Psi\mu_{T_B^c(M)})(\text{corad}(T_B^c(M) \otimes T_B^c(M))) = 0$ and $(\pi_1\mu_{T_{B'}^c(M')}(\Psi \otimes \Psi))(\text{corad}(T_B^c(M) \otimes T_B^c(M))) = 0$. Thus by the universal property of $T_{B'}^c(M')$, in order to prove $\Psi\mu_{T_B^c(M)} = \mu_{T_{B'}^c(M')}(\Psi \otimes \Psi)$, we only need to show $\pi_n\Psi\mu_{T_B^c(M)} = \pi_n\mu_{T_{B'}^c(M')}(\Psi \otimes \Psi)$ for $n = 1, 2$. However, this follows from a straightforward computation dual to Part (i). Furthermore, one can see $\Psi(1) = 1$. Hence Ψ is an algebra map, and so a Hopf algebra map. $\square$

The Proof of Lemma 1.6: (i) $\Rightarrow$ (ii) Assume that $\phi : B \rightarrow B'$ is a Hopf algebra isomorphism and $\psi : M \rightarrow {}^{\phi^{-1}}M'^{\phi^{-1}}$ is a B -Hopf bimodule isomorphism. Then $\phi^{-1} : B' \rightarrow B$ is also a Hopf algebra isomorphism and $\psi^{-1} : M' \rightarrow {}^{\phi}_{\phi^{-1}}M^{\phi}_{\phi^{-1}}$ is a B' -Hopf bimodule isomorphism. By Lemma 14.1.6(i), we have two graded Hopf algebra homomorphisms $T_B(\iota_0\phi, \iota_1\psi) : T_B(M) \rightarrow T_{B'}(M')$ and $T_{B'}(\iota_0\phi^{-1}, \iota_1\psi^{-1}) : T_{B'}(M') \rightarrow T_B(M)$. Then we have

$$\begin{aligned} T_{B'}(\iota_0\phi^{-1}, \iota_1\psi^{-1})T_B(\iota_0\phi, \iota_1\psi)\iota_0 &= T_{B'}(\iota_0\phi^{-1}, \iota_1\psi^{-1})\iota_0\phi \\ &= \iota_0\phi^{-1}\phi = \iota_0 \end{aligned}$$

and

$$\begin{aligned} T_{B'}(\iota_0\phi^{-1}, \iota_1\psi^{-1})T_B(\iota_0\phi, \iota_1\psi)\iota_1 &= T_{B'}(\iota_0\phi^{-1}, \iota_1\psi^{-1})\iota_1\psi \\ &= \iota_1\psi^{-1}\psi = \iota_1. \end{aligned}$$

It follows from the universal property of $T_B(M)$ that $T_{B'}(\iota_0\phi^{-1}, \iota_1\psi^{-1})T_B(\iota_0\phi, \iota_1\psi) = \text{id}_{T_B(M)}$. Similarly, one can show that $T_B(\iota_0\phi, \iota_1\psi)T_{B'}(\iota_0\phi^{-1}, \iota_1\psi^{-1}) = \text{id}_{T_{B'}(M')}$. Hence $T_B(M) \cong T_{B'}(M')$ as graded Hopf algebras.

(i) $\Rightarrow$ (iii) It is similar to the proof above.

(ii) $\Rightarrow$ (i) Assume that $f : T_B(M) \rightarrow T_{B'}(M')$ is a graded Hopf algebra isomorphism. Put $\phi := \pi_0 f \iota_0$ and $\psi := \pi_1 f \iota_1$. Then it is easy to check that ϕ is a Hopf algebra isomorphism from B to B' and ψ is a B -Hopf bimodule isomorphism from M to ${}^{\phi^{-1}}M'^{\phi^{-1}}$.

(iv) $\Rightarrow$ (i) Assume that φ is a graded Hopf algebra isomorphism from $B[M]$ to $B'[M']$. Let $\phi := \pi_0 \varphi \iota_0$ and $\psi := \pi_1 \varphi \iota_1$. Obviously, ϕ is a Hopf algebra isomorphism from B to B' and ψ is a bijection from M to M' . Now we show that ψ is a B -Hopf bimodule isomorphism from M to ${}^{\phi^{-1}}M'^{\phi^{-1}}$. Assume that $\alpha^-, \alpha^+, \delta^-$ and δ^+ are the module operations and comodule operations of M ; $\alpha'^-, \alpha'^+, \delta'^-$ and δ'^+ are the module operations and comodule operations of M' .

See that

$$\begin{aligned} \alpha'^-(\phi \otimes \psi) &= \pi_1 \mu(\iota_0 \pi_0 \varphi \iota_0 \otimes \iota_1 \pi_1 \varphi \iota_1) \\ &= \pi_1 \mu(\varphi \iota_0 \otimes \varphi \iota_1) \quad \text{and} \\ \psi \alpha^- &= \pi_1 \varphi \iota_1 \pi_1 \mu(\iota_0 \otimes \iota_1) \\ &= \pi_1 \varphi \mu(\iota_0 \otimes \iota_1) \\ &= \pi_1 \mu(\varphi \iota_0 \otimes \varphi \iota_1). \end{aligned}$$

Thus ψ is a left B -module homomorphism from M to ${}_{\phi}M'$. Similarly, we have that ψ is a right B -module homomorphism from M to M'_{ϕ} . See that

$$\begin{aligned} (\phi^{-1} \otimes id_{M'}) \delta'^{-} \psi &= (\phi^{-1} \otimes id_{M'}) (\pi_0 \otimes \pi_1) \Delta \iota_1 \pi_1 \varphi \iota_1 \\ &= (\phi^{-1} \pi_0 \otimes \pi_1) \Delta \varphi \iota_1 \quad \text{and} \\ (id_B \otimes \psi) \delta^{-} &= (\phi^{-1} \pi_0 \varphi \iota_0 \pi_0 \otimes \pi_1 \varphi \iota_0 \pi_0) \Delta \iota_1 \\ &= (\phi^{-1} \pi_0 \otimes \pi_1) \Delta \varphi \iota_1. \end{aligned}$$

Thus ψ is a left B -comodule homomorphism from M to ${}^{\phi^{-1}}M'$. Similarly, we have that ψ is a right B -comodule homomorphism from M to $M'{}^{\phi^{-1}}$.

(iii) $\Rightarrow$ (iv) Obvious. $\square$

The proof of Theorem 14.1.9: (i) $\Rightarrow$ (ii). Assume that (Q, G, r) is a Hopf quiver, where r is a ramification of G . For any $C \in \mathcal{K}(G)$, let $M(C)$ be a trivial right $kZ_{u(C)}$ -module with $\dim M(C) = r_C$. That is, $m \cdot h = m$ for any $m \in M(C)$ and $h \in Z_{u(C)}$. From the proof of Theorem 14.1.1, $M := \bigoplus_{y=xg_{\theta}^{-1}u(C)g_{\theta}, x,y \in G} (x \otimes M(C) \otimes_{kZ_{u(C)}} g_{\theta})$ is a kG -Hopf bimodule. Let $\{\xi_C^{(i)} \mid i \in I_C(r)\}$ be a basis of $M(C)$ over k . Let $\phi : kQ_1 \rightarrow M$ be a k -linear map defined by

$$\phi(a_{x,y}^{(i)}) = x \otimes \xi_C^{(i)} \otimes_{kZ_{u(C)}} g_{\theta},$$

where $x, y \in G$ with $x^{-1}y = g_{\theta}u(C)g_{\theta}^{-1}$, $C \in \mathcal{K}_r(G)$ and $i \in I_C(r)$. It is easy to see that ϕ is a k -linear isomorphism and a kG -bicomodule homomorphism from kQ_1^c to M . It follows that the arrow comodule kQ_1^c admits a kG -Hopf bimodule structure.

(ii) $\Rightarrow$ (i). Assume that the arrow comodule kQ_1^c admits a kG -Hopf bimodule structure. Then by Theorem 14.1.1 and its proof, there is a vector space isomorphism ${}^y(kQ_1^c)^x \cong x \otimes {}^{u(C)}(kQ_1^c)^1 \otimes_{kZ_{u(C)}} g_{\theta}$ for any $x, y \in G$ with $y = xg_{\theta}u(C)g_{\theta}^{-1}$. Hence for any $x, y \in G$ with $y = xg_{\theta}u(C)g_{\theta}^{-1}$, we have $\dim {}^y(kQ_1^c)^x = \dim {}^{u(C)}(kQ_1^c)^1$. This shows that Q is a Hopf quiver.

(ii) $\Leftrightarrow$ (iii) It follows from Lemma 14.1.8. $\square$

The proof of Lemma 14.2.7: We first show that ${}_{\phi}^{\phi^{-1}}\mathcal{B}(V)$ is a Nichols algebra of ${}_{\phi}^{\phi^{-1}}V$ by following steps. Let $R := {}_{\phi}^{\phi^{-1}}\mathcal{B}(V)$.

(i) Obviously, $R_0 = k$, $R_1 = P(R)$ and R is generated by R_1 as algebras. Since $C(x \otimes y) = C'(x' \otimes y')$ for any $x', y' \in R$, $x, y \in \mathcal{B}(V)$ with $x' = x, y' = y$, where C and C' denote the braidings in ${}^B_B\mathcal{YD}$ and ${}^{B'}_{B'}\mathcal{YD}$, respectively, we have that R is a graded braided Hopf algebra in ${}^{B'}_{B'}\mathcal{YD}$.

(ii) $R_1 = {}_{\phi}^{\phi^{-1}}\mathcal{B}(V)_1 \cong {}_{\phi}^{\phi^{-1}}V$ as YD B' -modules, since $\mathcal{B}(V)_1 \cong V$ as YD B -modules.

Consequently, ${}_{\phi}^{\phi^{-1}}\mathcal{B}(V)$ is a Nichols algebra of ${}_{\phi}^{\phi^{-1}}V$. By [7, Proposition 2.2 (iv)], $R \cong \mathcal{B}({}_{\phi}^{\phi^{-1}}V)$ as graded braided Hopf algebra in ${}^{B'}_{B'}\mathcal{YD}$. $\square$

Proposition 15.0.32. *If $(kQ_1^c, G, r, \vec{\chi}, u)$ is a kG -Hopf bimodule, then $(kG)^*$ -coactions on the $(kG)^*$ -Hopf bimodule $(kQ_1^a, G, r, \vec{\chi}, u)$ are given by*

$$\delta^{-}(a_{y,x}^{(i)}) = \sum_{h \in G} p_h \otimes a_{h^{-1}y, h^{-1}x}^{(i)}, \quad \delta^{+}(a_{y,x}^{(i)}) = \sum_{h \in G} \chi_C^{(i)}(\zeta_{\theta}(h^{-1})^{-1}) a_{yh^{-1}, xh^{-1}}^{(i)} \otimes p_h$$

where $x, y \in G$ with $x^{-1}y = g_{\theta}^{-1}u(C)g_{\theta}$, ζ_{θ} is given by (14.3), $C \in \mathcal{K}_r(G)$, $i \in I_C(r)$ and $p_h \in (kG)^*$ is defined by $p_h(g) = \delta_{h,g}$ for all $g, h \in G$.

Proof. Let $(kQ_1^c, G, r, \vec{\chi}, u) = (kQ_1^c, \alpha^-, \alpha^+)$ and $(kQ_1^a, G, r, \vec{\chi}, u) = (kQ_1^a, \delta^-, \delta^+)$. Since (kQ_1^c, kQ_1^a) is an arrow dual pairing, $\xi_{kQ_1^a}$ is a $(kG)^*$ -Hopf bimodule isomorphism from $(kQ_1^a, \delta^-, \delta^+)$ to $((kQ_1^c)^*, \alpha^{-*}, \alpha^{+*})$ (see Lemma 14.1.8). Now let $h, x, y, v, w \in G$ with $x^{-1}y = g_\theta^{-1}u(C)g_\theta$, where $C \in \mathcal{K}_r(G)$, $\theta \in \Theta_C$. By Proposition 14.1.11, we have

$$\begin{aligned} \langle \delta^-((a_{y,x}^{(i)})^*), h \otimes a_{w,v}^{(j)} \rangle &= \langle (a_{y,x}^{(i)})^*, h \cdot a_{w,v}^{(j)} \rangle \\ &= \langle (a_{y,x}^{(i)})^*, a_{hw,hv}^{(j)} \rangle \\ &= \delta_{x,hv} \delta_{y,hw} \delta_{i,j} \\ &= \delta_{h^{-1}x,v} \delta_{h^{-1}y,w} \delta_{i,j}. \end{aligned}$$

This shows that $\delta^-((a_{y,x}^{(i)})^*) = \sum_{h \in G} p_h \otimes (a_{h^{-1}y, h^{-1}x}^{(i)})^*$. Hence the left $(kG)^*$ -coaction of the $(kG)^*$ -Hopf bimodule $(kQ_1^a; G, r, \vec{\chi}, u)$ is given by

$$\delta^-((a_{y,x}^{(i)})^*) = \sum_{h \in G} p_h \otimes a_{h^{-1}y, h^{-1}x}^{(i)}.$$

In order to show the second equation, we only need to show that

$$\langle \delta^+((a_{y,x}^{(i)})^*), a_{wv}^{(j)} \otimes h \rangle = \sum_{g \in G} \langle \chi_C^{(i)}(\zeta_\theta(g^{-1})^{-1})(a_{yg^{-1}, xg^{-1}}^{(i)})^* \otimes p_g, a_{wv}^{(j)} \otimes h \rangle \quad (15.8)$$

for any $h, v, w \in G, D \in \mathcal{K}(G), j \in I_D(r)$ and $v^{-1}w \in D$, since $\xi_{kQ_1^a}$ is a $(kG)^*$ -Hopf bimodule isomorphism. If $v \neq xh^{-1}$ or $w \neq yh^{-1}$, then there exists $\beta \in k$ such that $a_{w,v}^{(j)} \cdot h = \beta a_{wh, vh}^{(j)}$ by Proposition 14.1.11. By simple computation, both the right hand side and the left hand side of (15.8) are zero.

Now assume $v = xh^{-1}$ and $w = yh^{-1}$. Assume $g_\theta h^{-1} = \zeta_\theta(h^{-1})g_{\theta'}$. Thus $v^{-1}w = hg_\theta^{-1}u(C)g_\theta h^{-1} = g_{\theta'}^{-1}\zeta_\theta(h^{-1})^{-1}u(C)\zeta_\theta(h^{-1})g_{\theta'} = g_{\theta'}^{-1}u(C)g_{\theta'}$. Since $g_{\theta'}h = \zeta_{\theta'}(h^{-1})^{-1}g_\theta$, $\zeta_{\theta'}(h) = \zeta_\theta(h^{-1})^{-1}$. Now we have

$$\begin{aligned} \text{the left hand side of (15.8)} &= \langle \delta^+((a_{y,x}^{(i)})^*), a_{w,v}^{(j)} \otimes h \rangle \\ &= \langle (a_{y,x}^{(i)})^*, a_{w,v}^{(j)} \cdot h \rangle \\ &= \chi_C^{(j)}(\zeta_{\theta'}(h)) \langle (a_{y,x}^{(i)})^*, a_{wh, vh}^{(j)} \rangle \\ &= \chi_C^{(j)}(\zeta_{\theta'}(h)) \delta_{i,j} \\ &= \chi_C^{(i)}(\zeta_\theta(h^{-1})^{-1}) \delta_{i,j} \\ &= \text{the right hand side of (15.8)}. \quad \square \end{aligned}$$

Now, we give an interesting quantum combinatoric formula by means of multiple Taft algebras. Let n be a positive integer and S_n be the symmetric group on the set $\{1, 2, \dots, n\}$. For any permutation $\sigma \in S_n$, let $\tau(\sigma)$ denote the number of reverse order of σ , i.e., $\tau(\sigma) = |\{(i, j) \mid 1 \leq i < j \leq n, \sigma(i) > \sigma(j)\}|$. For any $0 \neq q \in k$, let $S_n(q) := \sum_{\sigma \in S_n} q^{\tau(\sigma)}$.

Lemma 15.0.33. *In $kQ^c(G, r, \vec{\chi}, u)$, assume $g \in \mathcal{K}_r(G)$ and $j \in I_{\{g\}}(r)$. Let $q := \chi_{\{g\}}^{(j)}(g)$. If $i_1, i_2, \dots, i_m$ be non-negative integers, then*

$$a_{g^{i_m+1}, g^{i_m}}^{(j)} \cdot a_{g^{i_{m-1}+1}, g^{i_{m-1}}}^{(j)} \cdots a_{g^{i_1+1}, g^{i_1}}^{(j)} = q^{\beta_m + \frac{m(m-1)}{2}} S_m(q^{-1}) P_{g^{\alpha_m}}^{(j)}(g, m)$$

where $\alpha_m = i_1 + i_2 + \dots + i_m$, $P_h^{(j)}(g, m) = a_{g^m h, g^{m-1} h}^{(j)} a_{g^{m-1} h, g^{m-2} h}^{(j)} \cdots a_{g h, h}^{(j)}$, $\beta_1 = 0$ and $\beta_m = \sum_{j=1}^{m-1} (i_1 + i_2 + \dots + i_j)$ if $m > 1$.

Proof. Now let $a_l = a_{g^{i_l+1}, g^{i_l}}^{(j)}$ for $l = 1, 2, \dots, m$ and consider the sequence of m arrows $A = (a_m, \dots, a_1)$. We shall use the notations of [36, p.247]. For any $\sigma \in S_m$ and $1 \leq l \leq m$, let $m(\sigma, l) = |\{i \mid i < l, \sigma(i) < \sigma(l)\}|$. Then

$$m(\sigma, l) = (l-1) - |\{i \mid i < l, \sigma(i) > \sigma(l)\}|,$$

and hence

$$\begin{aligned} \sum_{l=1}^m m(\sigma, l) &= \frac{m(m-1)}{2} - \sum_{l=1}^m |\{i \mid i < l, \sigma(i) > \sigma(l)\}| \\ &= \frac{m(m-1)}{2} - \tau(\sigma). \end{aligned}$$

Now it follows from [36, Proposition 3.13] that

$$\begin{aligned} a_m \cdot a_{m-1} \cdot \dots \cdot a_1 &= \sum_{\sigma \in S_m} A_m^\sigma \cdots A_2^\sigma A_1^\sigma \\ &= \sum_{\sigma \in S_m} \left(\prod_{l=1}^m q^{i_1+i_2+\dots+i_{\sigma(l)-1}+m(\sigma, l)} \right) P_{g^{\alpha_m}}^{(j)}(g, m) \\ &= q^{\beta_m + \frac{m(m-1)}{2}} S_m(q^{-1}) P_{g^{\alpha_m}}^{(j)}(g, m). \quad \square \end{aligned}$$

Let $G \cong \mathbf{Z}$ be the infinite cyclic group with generator g and let r be a ramification of G given by $r_{\{g\}} = 1$ and $r_{\{g^n\}} = 0$ if $n \neq 1$. Let $0 \neq q \in k$. Define $\chi_g^{(1)} \in \widehat{G}$ by $\chi_g^{(1)}(g) = q$. Then $(G, r, \overrightarrow{\chi}, u)$ is an *RSC*. Using Lemma 14.3.3 and Lemma 15.0.33, one gets the following result.

Example 15.0.34. (i) For any $0 \neq q \in k$, $(m)_q! = q^{\frac{m(m-1)}{2}} S_m(q^{-1})$, and $S_m(q) = (m)_1! q^{\frac{m(m-1)}{2}}$, where m is a positive integer.

(ii) Assume q is a primitive n -th root of unity with $n > 1$. Then $S_m(q) = 0$ if $m \geq n$, and $S_m(q) \neq 0$ if $0 < m < n$.

Let R and B be algebras over field k and H be a Hopf algebra over k .

If B is a right H -comodule algebra with $R \subseteq B$ and $B^{coH} = R$, then B is called an H -extension of A . Furthermore, if there exists a convolution-invertible right H -comodule homomorphism γ from H to B , then B is called an H -cleft extension.

Theorem 15.0.35. (see [87, theorem 7.2.2]) An H -extension $R \subseteq B$ is an H -cleft iff

$$B \cong R \#_\sigma H \quad \text{as algebras.}$$

Sketch of proof. If $R \subseteq B$ is an H -cleft, then there is a convolution-invertible right H -comodule homomorphism γ from H to B . Set

$$\alpha(h, a) = \sum \gamma(h_1) a \gamma^{-1}(h_2)$$

and

$$\sigma(h, k) = \sum \gamma(h_1) \gamma(k_1) \gamma^{-1}(h_2 k_2)$$

for any $h, k \in H, a \in R$. We can show that the conditions in Theorem 4.1.1 are satisfied. Thus $R \#_\sigma H$ is an algebra. We can check that Φ is an algebra isomorphism from $R \#_\sigma H$ to B by sending $a \# h$ to $a \gamma(h)$ for any $a \in R, h \in H$.

Conversely, if $R\#_{\sigma}H$ is an algebra, we define that $B = R\#_{\sigma}H$ and $\gamma(h) = 1\#h$ for any $h \in H$. It is clear that the convolution-inverse of γ is

$$\gamma^{-1}(h) = \sum \sigma^{-1}(S(h_2), h_3)\#S(h_1)$$

for any $h \in H$. We can show that $R \subseteq B$ is an H -cleft extension. $\square$

Theorem 15.0.36. *(see [90, theorem 7.2.2]) Let H be a finite-dimensional, semisimple and either commutative or cocommutative. Let $A\#_{\sigma}H$ be a crossed product. Assume that R is H -prime. Then $R\#_{\sigma}H$ has only finite many minimal prime ideals and their intersection is zero.*

Index

- A^{op}, A^{cop} 7.3
 ad Ch.4
 $A \#_{\sigma} H, A_{\alpha, \sigma} \#^{\psi, Q} H$, Ch.3
 $A \bowtie H, A \overset{c}{\bowtie} H, A^{\phi} \bowtie^{\psi} H$ Ch.3
 $A \bowtie^R H$ 7.4
 $A \bowtie_r H$ 7.4
 $CYBE$ Ch.13
 C_0 9.1
 η , Ch.1
 d_V , Ch.1
 H_4 , 5.2
 $\int_H^l, \int_H^r$ Ch.1
 m Ch.1
 $M^{(\infty)}$ 9.1
 $O(H, \bar{\Delta})$, Ch.1
 $QYBE$ Ch.13
 S Ch.1
 $\rightharpoonup$, 8.1
 action , adjoint action, Ch.4
 algebra , Ch.1
 almost commutative, Ch.7
 anyonic Hopf algebra, 8.1
 bialgebra, Ch.1
 $A^*, A^{\hat{*}}$ 7.3, 8.1, Ch.1
 $A \# H$ 4.1
 $A^{\phi, P} \#_{\beta, \mu} H, A_{\alpha}^{\phi} \# H$, Ch.3
 $A_{\alpha}^{\phi} \bowtie_{\beta}^{\psi} H, A \overset{b}{\bowtie} H$ 3.1
 $A \bowtie_{\tau} H$ 7.3
 C, C^r, C^R , Ch.1
 $C(M)$ 9.1
 Δ, ϵ , Ch.1
 $f * g$, Ch.1
 b_V , Ch.1
 i_A, i_H , Ch.3
 rgD, wD, lpd , Ch.10
 M_0 , 9.1
 M^H, M^{coH} , Ch.1
 π_A, π_H , Ch.3
 $r_H, r_{Hb}, r_{bH}, r_{Hj}, r_{jH}$ 12.1, 12.2, 12.3
 $\Delta(h) = \sum (h_1 \otimes h_2)$ Ch.1
 I , Ch.1
 adjoint coaction, Ch.4
 almost cocommutative, Ch.7
 antipode, Ch.1
 anyonic algebra, 2.1

biproduct, bicrossproduct , 4.1
 braided group, *Ch.1*
 Braided tensor category , *Ch.1*
 classical Yang-Baxter equation, 260
 cobound Lie bialgebra, *Ch.13*
 coevaluation , *Ch.1*
 comodule, comultiplication , *Ch.1*
 component, 9.1
 coradical , 9.1
 double bicrossproduct, *Ch.3*
 Drinfeld double , 7.3
 ϵ Lie algebra , 2.1
 graded algebra, graded coalgebra, 2.2
 Hopf module , 2.1
 H -Hopf module, 2.1
 H - m -sequence, *Ch.11*
 H -radical, *Ch.10*,

 left dual, *Ch.1*
 Maschke's theorem, 2.1
 quantum Yang-Baxter equation , 7.1
 quasitriangular, *Ch.1*
 relative irreducible, 9.1
 restricted Lie algebra , 10.2
 strongly symmetric element , 13.2
 superspace, super-Hopf algebra , 2.1
 α, β - skew symmetric, 13.2
 skew group ring, *Ch.10*
 tensor category, *Ch.1*
 twisted H -module, *Ch.4*
 weak action, 79
 weak-closed, 9.1
 Yetter – Drinfeld module, *Ch.1*

 Braiding, *Ch.1*
 braided Hopf algebra, *Ch.1*
 braided group analogue, 5.1
 coalgebra , *Ch.1*
 2-cocycle, 2-cycle , *Ch.4*
 cocommutative, commutative , *Ch.4*
 convolution , *Ch.1*
 coquasitriangular , *Ch.1*
 counit , *Ch.1*
 cross (co)product, *Ch.3*
 evaluation, *Ch.1*
 full H -comodule, 9.1
 Hopf algebra , *Ch.1*
 Hopf module algebra , 11.1
 H -module (co)algebra, 11.1
 H - m -nilpotent , *Ch.11*
 inner action, *Ch.4*
 integral, *Ch.1*
 Lie colour algebra, Lie H-algebra, 2.1
 module algebra, module coalgebra, *Ch.1*
 quantum group, *Ch.1*
 quantum commutative , 4.3
 R -matrix, *Ch.7*
 smash coproduct, smash product, *Ch.4*
 symmetric tensor category , *Ch.1*
 α, β -symmetric, 13.4
 skew pairing, skew copairing, *Ch.7*
 triangular Hopf algebra, 2.1
 twist map, *Ch.1*
 wedge product, 9.1
 weak R -matrix, weak r -comatrix, *Ch.7*
 w-relational hereditary, 9.1

Bibliography

- [1] E. Abe. Hopf Algebra. Cambridge University Press, 1980.
- [2] F. W. Anderson and K. P. Fuller. Rings and Categories of Modules. Springer-Verlag. New York, 1974.
- [3] M. Auslander, I. Reiten and S.O. Smalø, Representation theory of Artin algebras, Cambridge University Press, 1995.
- [4] N. Andruskewisch and H.J.Schneider, Hopf algebras of order p^2 and braided Hopf algebras, J. Alg. **199** (1998), 430–454.
- [5] N. Andruskewitsch and H. J. Schneider, Lifting of quantum linear spaces and pointed Hopf algebras of order p^3 , J. Alg. **209** (1998), 645–691
- [6] N. Andruskewitsch and H. J. Schneider, Finite quantum groups and Cartan matrices, Adv. Math., **154** (2000), 1–45.
- [7] N. Andruskewisch and H.J.Schneider, Pointed Hopf algebras, new directions in Hopf algebras, edited by S. Montgomery and H.J. Schneider, Cambridge University Press, 2002.
- [8] M. Aslam and A. M. Zaidi, The matrix equation in radicals, Studia Sci.Math.Hungar., 28 (1993), 447–452.
- [9] R. J. Blattner, M. Cohen and S. Montgomery, Crossed products and inner actions of Hopf algebras, Transactions of the AMS., **298** (1986)2, 671–711.
- [10] A. A. Belavin and V. G. Drinfel'd. Solutions of the classical Yang–Baxter equations for simple Lie algebras. Functional Anal. Appl, **16** (1982)3, 159–180.
- [11] Y.Bespalov, B.Drabant, Hopf (bi-)modules and crossed modules in braided monoidal categories, J. Pure and Applied Algebra, 123(1998), 105-129.
- [12] Yuri Bespalov, Bernhard Drabant, Cross Product Bialgebras - Part I, J ALGEBRA, 219 (1999), 466-505.
- [13] Yu. Bespalov. Crossed Modules and quantum groups in braided categories. Applied categorical structures, **5**(1997), 155–204.

- [14] Y. Bahturin, D. Fischman and S. Montgomery. On the generalized Lie structure of associative algebras. Israel J. of Math., **96**(1996) , 27–48.
- [15] Y. Bahturin, D. Fischman and S. Montgomery, Bicharacter, twistings and Scheunert's theorem for Hopf algebra, J. Alg. **236** (2001), 246-276.
- [16] Y. Bahturin, D. Mikhalev, M. Zaicev and V. Petrogradsky, Infinite dimensional Lie superalgebras, Walter de Gruyter Publ. Berlin, New York, 1992.
- [17] E. Beggs and S. Majid. Matched pairs of topological Lie algebras corresponding to Lie bialgebra structures on $\operatorname{diff}(S^1)$ and $\operatorname{diff}(R)$. Ann. Inst. H. Poincare, Phys., Theor. **53**(1990)1, 15–34.
- [18] J. Bergen and S. Montgomery. Ideal and quotients in crossed products of Hopf algebras. J. Algebra **125** (1992), 374–396.
- [19] R. J. Blattner and S. Montgomery. Crossed products and Galois extensions of Hopf algebras. Pacific Journal of Math. **127** (1989), 27–55.
- [20] R. J. Blattner and S. Montgomery. A duality theorem for Hopf module algebras. J. algebra, **95** (1985), 153–172.
- [21] X. W. Chen, H. L. Huang, Y. Ye and P. Zhang, Monomial Hopf algebras, J. Alg. **275** (2004), 212–232.
- [22] Charles W. Curtis, Irving Reiner. Representation theory of finite group and associative algebras, New York:John Wiley&Sons, 1962.
- [23] William Chin. Crossed products of semisimple cocommutative of Hopf algebras. Proceeding of AMS, **116** (1992)2, 321–327.
- [24] William Chin. Crossed products and generalized inner actions of Hopf algebras. Pacific Journal of Math., **150** (1991)2, 241–259.
- [25] H.X.Chen, Quantum double in monoidal categories, Comm.Algebra, **28** (2000)5, 2303–2328.
- [26] Huixiang Chen. Quasitriangular structures of bicrossed coproducts. J. algebra, **204** (1998), 504–531.
- [27] W. Chin and S. Montgomery, Basic coalgebras, modular interfaces, AMS/IP Stud. Adv. Math., **4**, Amer. Math. Soc., Providence, RI, 1997, pp.41–47.
- [28] M. Cohen and S. Montgomery. Smash products and group acting. Trans. Amer. Math. Soc., **281** (1984)1, 237–258.
- [29] M. Cohen and S. Montgomery. Group-graded rings, smash products, and group actions. Trans. Amer. Math. Soc., **282** (1984)1, 237–258.

- [30] L. Corwin, Y. Ne'eman, and S. Sternberg. Graded Lie algebras in mathematics and physics (Bose-fermi symmetry). *Rev. Mod. Phys.*, **47** (1975), 573–604.
- [31] M. Cohen and S. Westreich. From supersymmetry to quantum commutativity. *J. algebra*, **168** (1994), 1–27.
- [32] Chuanren Cai and Jianhuan Sun. Hopf -Jacobson radical of Hopf module algebras. *Chinese Science Bulletin*, **41**(1996)41, 621–625.
- [33] H.X.Chen, Quantum double in monoidal categories, *Comm.Algebra*, **28** (2000)5, 2303–2328.
- [34] Weixin Chen and Shouchuan Zhang. The module theoretic characterization of special and supernilpotent radicals for Γ -rings. *Math.Japonic*, **38**(1993)3, 541–547.
- [35] Ru-Cheng Cao, *Combination Mathematics*, South China University of technology Press, 2002.
- [36] C. Cibils and M. Rosso, Hopf quivers, *J. Alg.*, **254** (2002), 241-251.
- [37] C. Cibils and M. Rosso, Algebres des chemins quantiques, *Adv. Math.*, **125** (1997), 171–199.
- [38] N. Divinsky. *Rings and Radicals*. Allen, London, 1965.
- [39] Y. Doi. Braided bialgebras and quadratic bialgebras. *Communications in algebra*, **5** (1993)21, 1731–1749.
- [40] S.Dascalescu, C.Nastasecu and S. Raianu, *Hopf algebras: an introduction*, Marcel Dekker Inc. , 2001.
- [41] V. G. Drinfel'd. On some unsolved problems in Quantum group theory. In “Proceedings, Leningrad 1990, p. 394”. *Lecture Notes in Mathematics*, Vol. 1510, Springer-Verlag. Berlin, 1992.
- [42] V. G. Drinfel'd. Quantum groups. In “Proceedings International Congress of Mathematicians, August 3-11, 1986, Berkeley, CA” pp. 798–820, Amer. Math. Soc., Providence, RI, 1987.
- [43] Y. Doi and M. Takeuchi. Multiplication algebra by two-cocycles –the quantum version–. *Communications in algebra*, **14**(1994)22, 5715–5731.
- [44] Pavel Etingof, Shlomo Gelaki, The classification of finite-dimensional triangular Hopf algebras over an algebraically closed field of characteristic 0, *Internat. Math. Res. Notices*, **5**(2000), 223-234.
- [45] C.Faith, *Algebra : Rings, modules and categories*, Springer-Verlag, 1973.

- [46] I. Frenkel and Y. Zhu, Vertex operator algebras associated to representations of affine and Virasoro algebras, *Duke Math. J.* , **66** (1992), 123-168.
- [47] T.Kerler, Bridged links and tangle presentations of cobordism categories, *Adv. Math.* **141** (1999), 207– 281.
- [48] J.R. Fisher. The Jacobson radicals for Hopf module algebras. *J. algebra*, **25**(1975), 217–221.
- [49] X. Gomez and S. Majid, Braided Lie algebras and bicovariant differential calculi over coquasitriangular Hopf algebras, *J. Alg.* **261**(2003), 334–388.
- [50] D. Gurevich, A. Radul and V. Rubtsov, Noncommutative differential geometry related to the Yang-Baxter equation, *Zap. Nauchn. Sem. S.-Peterburg Otdel. Mat. Inst. Steklov. (POMI)* **199** (1992); translation in *J. Math. Sci.* **77** (1995), 3051–3062.
- [51] D. I. Gurevich, The Yang-Baxter equation and the generalization of formal Lie theory, *Dokl. Akad. Nauk SSSR*, **288** (1986), 797–801.
- [52] J.Hietarinta. all solutions to the constant quantum Yang-Baxter equation in two dimensions. *Phy.Letters A*, **165** (1992), 245–251.
- [53] M. A. Hennings, Hopf algebras and regular isotopy invariants for link diagrams, *Proc. Cambridge Phil. Soc* **109** (1991), 59-77.
- [54] R. G. Heyneman and D. E. Radford. Reflexivity and coalgebras of finite type. *J. Algebra* **28** (1974), 215–246.
- [55] G.Q. Hu and Y. H. Xu. Quantum commutative algebras and dualities. *Science in China, ser A*, **26** (1996) 10, 892–900.
- [56] T. W. Hungerford, *Algebra*, GTM **73**, New York:Springer-Verlag, 1974.
- [57] Guan-Zhang Hu, *Applied Modern Algebras*, Qinghua University Press, Beijing, 2005.
- [58] L. Kauffman, invariants of links and 3-manifolds via Hopf algebras, in *Geometry and Physics*, Marcel Dekker Lecture Notes in Pure and Appl. Math. vol **184**, 1997, 471-479.
- [59] N. Jacobson. *Lie Algebras*. Interscience publishers a division of John Wiley and Sons , New York, 1962.
- [60] M. Jimbo. Introduction to the Yang-Baxter equation in: *Braid group, knot theory and statistical mechanics*. Eds. C. N. Yang and M. L. Ge (World scientific , Singapore, 1989).
- [61] A. Joyal and R. Street. Braided Monoidal Categories. *Math. reports* 86008, Macquaries university, 1986.
- [62] A. Joyal and R. Street. The geometry of tensor calculus I. *Adv. Math.* **88**(1991), 55–112.

- [63] A. Joyal and R. Street. Braided tensor categories. *Adv. Math.* **102**(1993), 20–78.
- [64] C. Kassel. *Quantum Groups*. Graduate Texts in Mathematics 155, Springer-Verlag, 1995.
- [65] V. G. Kac. Lie superalgebras. *Adv. in Math.*, **26**(1977) , 8–96.
- [66] V. K. Kharchenko, An existence condition for multilinear quantum operations, *J. Alg.* **217** (1999), 188–228.
- [67] R. G. Larson. characters of Hopf algebras. *J. algebra* **17** (1971), 352–368.
- [68] Shaoxue Liu. Baer radical and Levitzki radical for additive category. *J. Beijing Normal University* **4** (1987), 13–27(in Chinese).
- [69] Shaoxue Liu. *Rings and Algebras*. Science Press, 1983 (in Chinese).
- [70] R. G. Larson and D. E. Radford. Finite dimensional cosemisimple Hopf algebras in characteristic 0 are semisimple. *J. algebra* **117** (1988), 267–289.
- [71] R. G. Larson and D. E. Radford. Semisimple cosemisimple Hopf algebras. *Amer. J. Math.* **109** (1987), 187–195.
- [72] G. Lusztig, *Introduction to Quantum groups*, Progress Math. **110** , Berlin, 1993.
- [73] V. Lyubashenko. Tangles and Hopf algebras in braided tensor categories. *J.pure and applied algebra*, **98** (1995), 245–278.
- [74] S. Majid. Physics for algebraists: Non-commutative and non-cocommutative Hopf algebras by a bicrossproduct construction. *J.Algebra* **130** (1990), 17–64.
- [75] S. Majid. Quasitriangular Hopf algebras and Yang-Baxter equations. *Int. J. Mod. Phys. A*, **5** (1990), 1–91.
- [76] S. Majid. Beyond supersymmetry and quantum symmetry (an introduction to braided groups and braided matrices). (Proceeding of the 5th Nakai workshop, Tianjin, China), *Quantum Groups, Integrable Statistical Models and Knot Theory*, 1992, edited by M-L. Ge and H. J. de Vega, World Scientific, Singapore, 231–282, (1993).
- [77] S. Majid. Free braided differential calculus, braided binomial theorem, and the braided exponential map. *J. Math. Phys.*, **34**, 1993, 4843–4856 .
- [78] S. Majid. Braided groups. *J. Pure and Applied Algebra*, **86** (1993), 187–221.
- [79] S. Majid. Bicrossproduct structure of the quantum Weyl group. *J. algebra*, **163**(1994), 68–87.
- [80] S. Majid. Crossproducts by braided groups and bosonization. *J. algebra*, **165**(1994), 165–190.

- [81] S. Majid, Quantum and braided Lie algebras, J. Geom. Phys. **13** (1994), 307–356.
- [82] S. Majid. Algebras and Hopf algebras in braided categories. Lecture notes in pure and applied mathematics advances in Hopf algebras, Vol. 158, edited by J. Bergen and S. Montgomery, 1995.
- [83] S. Majid, Foundations of Quantum Group Theory, Cambridge University Press, 1995.
- [84] W. Michaelis. A class of infinite-dimensional Lie bialgebras containing the Virasoro algebra. Advances in mathematics, **107** (1994), 365–392.
- [85] W. Michaelis. Lie coalgebras. Advances in mathematics, **38** (1980), 1–54.
- [86] W. Michaelis. The dual Poincare-Birkhoff-Witt Theorem, Advances in mathematics. **57** (1985), 93–162.
- [87] S. Montgomery. Hopf algebras and their actions on rings. CBMS Number 82, Published by AMS, 1993.
- [88] S. Majid and M.J. Rodriguez Plaza. Random walk and the heat equation on superspace and anyspace. J. Math. Phys. **35**(1994) 7, 3753–3760.
- [89] J. C. McCommell and J. C. Robson. Noncommutative Noetherian Rings. John Wiley & Sons, New York, 1987.
- [90] S. Montgomery and H. J. Schneider. Hopf crossed products rings of quotients and prime ideals. Advances in Mathematics, **112** 1995, 1–55.
- [91] W. Nichols, Bialgebras of type one, Comm. Alg. **6** (1978), 1521–1552.
- [92] C. Nastasescu and F. van Oystaeyen. Graded Ring Theory. North-Holland Publishing Company, 1982.
- [93] W. D. Nichols and M. B. Zoeller. A Hopf algebras freeness theorem. Amer. J. Math., bf 111 (1989), 381–385.
- [94] F. Van Oystaeyen and P. Zhang, Quiver Hopf algebras, J. Alg., **280** (2004), 577–589.
- [95] D. S. Passman. The Algebraic Structure of Group Rings. John Wiley and Sons, New York, 1977.
- [96] G. K. Pedersen. C^* -Algebras and Their Automorphism Groups. Academic Press INC, (London), LTD, 1979.
- [97] H. C. Pop. Quantum group construction in a symmetric monoidal category. Communications in algebra, **25**, 1997, 117–158.
- [98] B. Pareigis, On Lie algebras in the category of Yetter-Drinfeld modules. Appl. Categ. Structures, **6** (1998), 151–175.

- [99] D.E. Radford. The structure of Hopf algebras with a projection. J. Alg., **92** , 1985, 322–347.
- [100] D. E. Radford. Minimal quasitriangular Hopf algebras. J. algebra **157** (1993), 281–315.
- [101] D. E. Radford. The order of antipode of a finite dimensional Hopf algebras is finite. Amer. J. Math. **98** (1976), 333–355.
- [102] D. E. Radford. On the quasitriangular structures of a semisimple Hopf algebra. J. algebra, **141** (1991), 354–358.
- [103] D. E. Radford, On Kauffman’s knot invariants arising from finite-dimensional Hopf algebras, in Advances in Hopf Algebras. Marcel Dekker Lecture Notes in Pure and Appl. Math, **158** (1994), 205–266.
- [104] D.E. Radford The trace function and Hopf algebras. J. algebra, **163** (1994), 583–622.
- [105] D. Robles-Llana and M. Rocek, Quivers, quotients, and duality, e-print hep-th/0405230.
- [106] J. J. Rotman. An introduction to homological algebras. Academic press, New York, 1979.
- [107] D.E.Radford, J.Towber, Yetter-Drinfeld categories associated to an arbitrary bialgebra, J. Pure and Applied Algebra, 87(1993) 259–279.
- [108] N. Yu. Reshetikhin and V.G. Turaev, Ribbon graphs and their invariants derived from quantum groups, Commun. Math. Phys., **127** (1990), 1–26.
- [109] A. D. Sands. Radicals and Morita contexts. J. algebra, **24**(1973), 335–345.
- [110] M. Scheunert. Generalized Lie algebras. J. Math. Phys., **20** (1979), 712–720.
- [111] Michio Suzuki, Group Theory I, New York:Springer-Verlag, 1978.
- [112] T. Shudo And H. Miyamoto. On the decomposition of coalgebras. Hiroshima Math. J., 8(1978), 499–504.
- [113] S. Stratila. Modular Theory in Operator Algebras. Editura Academiei and Abacus Press, 1981.
- [114] M. E. Sweedler. Hopf Algebras. Benjamin, New York, 1969.
- [115] M. E. Sweedler. Integrals for Hopf algebras. Ann. of Math. , **89** (1969), 323–335.
- [116] F. A. Szasz. Radicals of rings. John Wiley and Sons, New York, 1982.
- [117] E. J. Taft. Reflexivity of algebras and coalgebras. J. Amer. Math., **94** (1972)4, 1111–1130.
- [118] E. J. Taft. Witt and Virasoro algebras as Lie bialgebras. J.Pure Appl. algebra, **87** (1993), 301–312.
- [119] Wenting Tong. An Introduction to Homological Algebras. Chinese Education Press, 1998.

- [120] S. L. Woronowicz, Differential calculus on compactmatrix pseudogroups (quantum groups), Commun. Math. Phys., 122 (1989), 125-170.
- [121] Y. H. Xu and Y. Fong. On the decomposition of coalgebras. In “Proceedings International Colloquium on Words, Languages and Combinatorics- Kyoto, August 28–31, 1990”. pp. 504–522, World Scientific , Singapore, 1992.
- [122] Quan-Yan Xiang, Modern algebras, WuHan University Press, WuHan, 1995.
- [123] R. A. Brualdi, Introductory Combinatorics, Prentice Hall, 2004.
- [124] Y. H. Xu, K. P. Shum and Y. Fong. A decomposition Theory of Comodules. J. Algebra, 170(1994), 880–896.
- [125] C. N. Yang and M. L. Ge. Braid group, Knot theory and Statistical Mechanics. World scientific , Singapore, 1989.
- [126] Zhong Yi. Homological dimension of skew group rings and crossed products. Journal of algebra, **164** 1984, 101–123.
- [127] Shouchuan Zhang and Weixin Chen. The general theory of radicals and Baer radical for Γ -rings. J. Zhejiang University, **25** (1991), 719–724(in Chinese).
- [128] Shouchuan Zhang. Relation between decomposition of comodules and coalgebras. Science in China, **40** (1997), 25–29.
- [129] Shouchuan Zhang. The radicals of Hopf module algebras. Chinese Ann. Mathematics, Ser B, 18(1997)4, 495–502.
- [130] Shouchuan Zhang. The Baer and Jacobson radicals of crossed products. Acta math. Hungar., 78(1998), 11–24.
- [131] Shouchuan Zhang. Classical Yang-Baxter equation and low dimensional triangular Lie bialgebras. Physics Letters A, **246** (1998) , 71–81.
- [132] Zhang, Shouchuan, Chen, Hui-Xiang: The double bicrossproducts in braided tensor categories, Communications in Algebra **29**(2001)(1), 31–66.
- [133] Shouchuan Zhang, Duality theorem and Drinfeld double in braided tensor categories, Algebra colloq. **10** (2003)2, 127-134.
- [134] Shouchuan Zhang. The radicals of Hopf module algebras. Chinese Ann. Mathematics, Ser B, **18**(1997)4, 495–502.
- [135] Shouchuan Zhang. The H -von Neumann regular radical and the Jacobson radical of twisted graded algebra. Acta Math. Hungar., Acta Math. Hungar., 86(2000)4, 319-333.
- [136] Shouchuan Zhang. The strongly symmetric elements and Yang-Baxter equations. Physics Letters A, 261(1999)5-6, 275-283.

- [137] Shouchuan Zhang. The Homological dimension of crossed products. Chinese Math. Ann. 22A(2001), 767–772.
- [138] S.C. Zhang, The Baer radical of generalized matrix rings, in Proc. of the Sixth SIAM Conf. on Parallel Processing for Scientific Computing, pp.546–551, Norfolk, Virginia, 1993. Eds: R.F. Sincovec, D.E. Keyes, M.R. Leuze, L.R. Petzold, D.A. Reed. Also in math.RA/0403280.
- [139] Zhang S C, Zhang Y Z. Braided m -Lie algebras. Letters in Mathematical Physics, 2004, 70: 155-167. Also in math.RA/0308095.
- [140] Shouchuan Zhang, Y-Z Zhang, H.X. Chen, Classification of PM Quiver Hopf Algebras, Journal of Algebra and Its Applications, 6(2007)4, 1-32
- [141] Yuan-Da Zhang, The Structure of Finite Groups (I), Science Press, Beijing, 1982.
- [142] W. Zhao, S. H. Wang and Z. Jiao. On the Hopf algebra structure over double crossproduct. Communications in algebra, **26**(1998)2, 467–476.
- [143] X. G. Zou and Y. H. Xu. The decomposition of comodules. Science in China (Series A), 37(1994)8, 946–953.
- [144] X. Zhu, Finite representations of a quiver arising from string theory, e-print math.AG/0507316.